

CONGENITAL CARDIOLOGY TODAY

Timely News and Information for BC/BE Congenital/Structural Cardiologists and Surgeons

September 2015; Volume 13; Issue 9
North American Edition

IN THIS ISSUE

Rare Association of Retroaortic Innominate Vein and Coarctation of Aorta with Hypoplastic Left Aortic Arch: Embryology Review of Retroaortic Innominate Vein

Siddharth Patel, MD; Pooja Makhija, MD; Steffan Sernich, MD

~Page 1

CHIP Network Update

By Gary Webb, MD

~Page 8

A Case Series of Unfortunate Gerbode Defects

By Lucky R. Cuenza, MD; Anna Christina Adora, MD; Thessie Minelli O. Valdez, MD; Cheryl P. Fomaneg, MD; Efren R. Vicaldo, MD

~Page 11

Medical Meeting Focus

~Page 15

Medical News, Products & Information

~Page 18

Upcoming Medical Meetings

2nd Annual Basic & Advanced Fetal Cardiac Symposium and Workshop
Sep. 10-12, 2015; Chicago, IL USA
www.FetalCardiacSymposium.com

PICS-AICS

Sep. 18-21, 2015; Las Vegas, NV USA
www.picsymposium.com

34th Annual Echocardiography Symposium

Sep. 25-26, 2015; Miami, FL USA
MiamiEcho.Baptisthealth.net

48th Annual Southeast Pediatric Cardiology Society Conference
Sep. 25-26, 2015; Birmingham, AL USA
www.ChildrensAL.org/SPCS2015

CONGENITAL CARDIOLOGY TODAY

Editorial and Subscription Offices

16 Cove Rd, Ste. 200
Westerly, RI 02891 USA

www.CongenitalCardiologyToday.com

© 2015 by Congenital Cardiology Today
Published monthly. All rights reserved.

Recruitment advertisements on
pages: 3, 4, 7, 10, 12, 23

Rare Association of Retroaortic Innominate Vein and Coarctation of Aorta with Hypoplastic Left Aortic Arch: Embryology Review of Retroaortic Innominate Vein

Siddharth Patel, MD; Pooja Makhija, MD;
Steffan Sernich, MD

Introduction

The retroaortic course of the left innominate vein is a rare entity which can be misinterpreted during echocardiography for other abnormal vascular structures under the aortic arch. We reported the case of a 2-week-old female infant whose echocardiogram showed coarctation of the aorta and 2 vascular structures beneath the hypoplastic left aortic arch, one of which was traced to be a retroaortic left innominate vein. Right aortic arch is a common association.¹ More than 80% of the patients with anomalous retroaortic left innominate vein have obstruction of the right ventricular outflow tract (Tetralogy of Fallot).² However, our patient has a left sided aortic arch with hypoplastic transverse arch and juxtaductal coarctation of aorta without any right ventricular outflow tract obstruction.

Case

A newborn girl with prenatal diagnosis of hypoplastic aortic valve and hypoplastic aortic arch on fetal echocardiogram was delivered at a referring hospital. Her initial postnatal echocardiogram was read as normal for age with a patent ductus arteriosus; the patient was discharged home with a cardiology follow-up appointment. However, after missing her

appointment, she presented to the Emergency Department in cardiogenic shock at 2 weeks of age. She was hypotensive, tachycardic and tachypneic with cold extremities on presentation. Pulses were diminished in lower extremities with poor perfusion. She was intubated and started on mechanical ventilation due to worsening respiratory distress. Coarctation of aorta was suspected based on physical exam. Her initial echocardiography showed juxtaductal Coarctation of the aorta, tiny Patent Ductus Arteriosus (PDA) and mildly impaired left ventricular function. The patient was immediately started on intravenous prostaglandin E1, inotropic support, and metabolic acidosis was corrected.

Once patient was stabilized, a repeat complete echocardiogram demonstrated bicuspid aortic valve, juxtaductal coarctation of the aorta, patent ductus arteriosus, hypoplastic left aortic arch and two retroaortic vessels; the upper vessel being smaller in size was identified as the left innominate vein (Figure 1). Typically, there is only one vessel under the aortic arch, the right pulmonary artery. The differential diagnosis of an additional vessel in the suprasternal windows include an aorto-pulmonary collateral, a PDA arising from aortic arch, persistent fifth aortic arch, the vertical vein in supracardiac-type of Total Anomalous Pulmonary Venous Connection (TAPVC), and venous structures such as a retro-aortic innominate vein. Doppler interrogation

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share?

Submit your manuscript to: RichardK@CCT.bz

NIT-OCCLUD[®]

COIL SYSTEM FOR PDA CLOSURE



A SAFER, EASIER WAY TO CLOSE

The Nit-Occlud[®] coil system for PDA closure is designed to match individual morphologies and sizes

The delivery system facilitates optimal device positioning

The Nit-Occlud[®] PDA coil is repositionable and retrievable prior to release

Tight and compact windings ensure efficient occlusion

The Nit-Occlud[®] PDA coil was determined to be MRI conditional

Radiopaque

For more information or to place an order, contact your B. Braun Interventional Systems Inc. representative or call 1-877-VENA-CAV (836-2228)

pfmmedical

Manufacturer:
pfm medical, ag
Wankelstraße 60
50996 Köln, Germany
T +49 (0)2236 9641-10
F +49 (0)2236 9641-20



B | BRAUN

Interventional
Systems

Distributed by:
B. Braun Interventional Systems Inc.
824 Twelfth Avenue
Bethlehem, PA 18018 USA
Tel: 1-877-VENA CAV (836-2228) (USA)
Fax: 1-610-849-1334
www.bisusa.org

Indication: The Nit-Occlud[®] PDA coil is a permanently implanted prosthesis indicated for percutaneous, transcatheter closure of small to moderate size patent ductus arteriosus with a minimum angiographic diameter less than 4mm. Nit-Occlud Brief Statement: Do not implant the Nit-Occlud PDA into patients who have endocarditis, endarteritis, active infection, pulmonary hypertension (calculated PVR greater than 5 Wood Units), thrombus in a blood vessel through which access to the PDA must be obtained, thrombus in the vicinity of the implantation site at the time of the implantation or patients with a body weight < 11 lbs (5 kg). An angiogram must be performed prior to implantation for measuring length and diameter of the PDA. Only the pfm medical implantation delivery catheter should be used to implant the device. Administration of 50 units of heparin per kg body weight should be injected after femoral sheaths are placed. Antibiotics should be given before (1 dose) and after implantation (2 doses) in order to prevent infection during the implant procedure. Do not implant the Nit-Occlud PDA in an MR environment. Do not pull the Nit-Occlud coil through heart valves or ventricular chambers. Contrast media should not be injected through the implantation catheter. The catheter must not be connected to high pressure injectors. Patients may have an allergic response to this device due to small amounts of nickel that has been shown to be released from the device in very small amounts. If the patient experiences allergic symptoms, such as difficulty in breathing or swelling of the face or throat, he/she should be instructed to seek medical assistance immediately. Antibiotic prophylaxis should be performed to prevent infective endocarditis during first 6 months after coil implantation. Potential Adverse Events: Air embolism, Allergic reaction to drug/contrast, Apnea, Arrhythmia requiring medical treatment or pacing, Arteriovenous fistula, Bacterial endocarditis, Blood loss requiring transfusion, Chest pain, Damage to the tricuspid or pulmonary valves, Death, Embolization of the occluder, requiring percutaneous or surgical intervention, Endarteritis, False aneurysm of the femoral artery, Fever/Headache/migraine, Heart failure, Hemolysis after implantation of the occluder, Hypertension, Hypotension or shock, Infection, Myocardial infarction, Occluder fracture or damage, Perforation of the heart or blood vessels, Stenosis of the left pulmonary artery or descending thoracic aorta, Stroke/TIA, Thromboembolism (cerebral or pulmonary), Valvular Regurgitation, Vessel damage at the site of groin puncture (loss of pulse, hematoma etc.)




Figure 1. Suprasternal view shows retroaortic left innominate vein, hypoplastic arch, coarctation of aorta and patent ductus arteriosus.

demonstrated continuous low velocity pattern suggestive of venous structure. The probe was tilted leftward and anteriorly to trace the vessel which revealed venous flow from the left side of the neck passing below the aorta, directed to the right side and draining into the superior vena cava (Figure 2), confirming the diagnosis of retro-aortic left innominate vein.

The patient underwent a computerized topographic angiogram to assess the aortic arch which showed diffuse left aortic arch hypoplasia (3 mm) beyond the origin of the innominate artery with mild focal discrete juxtaductal coarctation of the aorta (2.5 mm) just distal to the left subclavian artery. Also, poststenotic dilation of the descending thoracic aorta (7 mm) and low origin of the left subclavian artery was noted (Figure 3).

The patient underwent coarctation of the aorta and hypoplastic aortic arch repair with bovine pericardium via a left lateral thoracotomy. The patient had an uneventful postoperative course.

Discussion

The left innominate vein normally courses obliquely downward to the right passing superoanterior to the aortic arch, and in front of its branches to drain into superior vena cava. Retro-aortic innominate vein is a rare entity where the innominate vein takes an anomalous course below the aortic arch with an incidence of 0.2-1.0% among congenital cardiac anomalies.¹ Right aortic arch is also a common association.¹ More than 80% of the patients with anomalous retroaortic left innominate vein have obstruction of the right ventricular outflow tract (Tetralogy of Fallot).² However, our patient has a left-sided aortic arch with a hypoplastic transverse arch and juxtaductal coarctation of the aorta, without any right ventricular outflow tract obstruction. One such case is described in literature so far.⁶

Embryologically, as most of the left cardinal veins disappear, the venous drainage from the left side of the head and neck and the left arm is directed into the right anterior cardinal vein by 2 transverse capillary

Pediatric Electrophysiologist

The Department of Pediatrics of Washington University School of Medicine in St. Louis, is seeking a pediatric cardiologist to join our team of 18 clinicians and basic scientists, based at St. Louis Children's Hospital.

We seek to recruit a **Pediatric Electrophysiologist** to join two full time faculty members in our arrhythmia program, which is directed by Jennifer N. Avari Silva, MD. The program offers the full range of diagnostic and interventional electrophysiology procedures, in a biplane cineangiography laboratory outfitted with Nav-X. This year, the program will do approximately 140 ablations, 40 pacemakers/ICDs, and 40 implantable loop recorders. We also offer a multidisciplinary inherited arrhythmia clinic, and Washington University offers a clinical genomic program which we access for genotyping. Opportunities for basic, translational and clinical research are robust, and we have close connections with researchers in the Department of Biomedical Engineering, the Department of Cell Biology and Physiology, and the Department of Developmental Biology at Washington University.

The ideal candidate must be eligible for licensure in Missouri, be board certified (or eligible) in pediatric cardiology, and have had advanced training in pediatric cardiac electrophysiology.

Washington University School of Medicine is consistently ranked as one of the best medical schools in the country, and is a longstanding leader in funding for pediatric research. St. Louis Children's Hospital is a 250 bed free-standing children's hospital established in 1879. The St. Louis Children's and Washington University Heart Center includes an active surgical program, a 16-bed Cardiac Intensive Care Unit, and one of the nation's largest pediatric heart failure and heart transplantation programs.

Interested candidates should provide a curriculum vitae and contact:

George F. Van Hare, M.D., F.H.R.S., F.A.C.C.
Director, Pediatric Cardiology
1 Children's Place
St. Louis, MO 63110
e-mail: vanhare@kids.wustl.edu
Phone: 314-454-4217

Washington University is an Equal Opportunity Affirmative Action Employer. We welcome CVs and resumes from women and from minority candidates.

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share?
Submit your manuscript to: RichardK@CCT.bz



PROVEN PERFORMANCE. SIMPLY DELIVERED.

MELODY TPV
DELAYS CONDUIT
REPLACEMENT

98%

Patients at 2 years

91%

Patients at 5 years

Data from US IDE Study



Melody[®]
Transcatheter Pulmonary Valve Therapy

Melody-TPV.com
Medtronic | Minneapolis, MN 55432-5604
Toll-free: 1 (800) 328-2518

UC201601525a EN ©2015 Medtronic.
All rights reserved 08/2015

Medtronic
Further. Together

Important Labeling Information for United States

Indications: The Melody TPV is indicated for use as an adjunct to surgery in the management of pediatric and adult patients with the following clinical conditions:

- Existence of a full (circumferential) RVOT conduit that was equal to or greater than 16 mm in diameter when originally implanted AND
- Dysfunctional RVOT conduits with a clinical indication for intervention, AND:
 - regurgitation: \geq moderate regurgitation, AND/OR
 - stenosis: mean RVOT gradient \geq 35 mm Hg

Contraindications: None known.

Warnings/Precautions/Side Effects

- DO NOT implant in the aortic or mitral position. Preclinical bench testing of the Melody valve suggests that valve function and durability will be extremely limited when used in these locations.
- DO NOT use if patient's anatomy precludes introduction of the valve, if the venous anatomy cannot accommodate a 22-Fr size introducer, or if there is significant obstruction of the central veins.
- DO NOT use if there are clinical or biological signs of infection including active endocarditis. Standard medical and surgical care should be strongly considered in these circumstances.
- Assessment of the coronary artery anatomy for the risk of coronary artery compression should be performed in all patients prior to deployment of the TPV.
- To minimize the risk of conduit rupture, do not use a balloon with a diameter greater than 110% of the nominal diameter (original implant size) of the conduit for pre-dilation of the intended site of deployment, or for deployment of the TPV.
- The potential for stent fracture should be considered in all patients who undergo TPV placement. Radiographic assessment of the stent with chest radiography or fluoroscopy should be included in the routine postoperative evaluation of patients who receive a TPV.
- If a stent fracture is detected, continued monitoring of the stent should be performed in conjunction with clinically appropriate hemodynamic assessment. In patients with stent fracture and significant associated RVOT obstruction or regurgitation, reintervention should be considered in accordance with usual clinical practice.

Potential procedural complications that may result from implantation of the Melody device include the following: rupture of the RVOT conduit, compression of a coronary artery, perforation of a major blood vessel, embolization or migration of the device, perforation of a heart chamber, arrhythmias, allergic reaction to contrast media, cerebrovascular events (TIA, CVA), infection/sepsis, fever, hematoma, radiation-induced erythema, blistering, or peeling of skin, pain, swelling, or bruising at the catheterization site.

Potential device-related adverse events that may occur following device implantation include the following: stent fracture, * stent fracture resulting in recurrent obstruction, endocarditis, embolization or migration of the device, valvular dysfunction (stenosis or regurgitation), paravalvular leak, valvular thrombosis, pulmonary thromboembolism, hemolysis.

**The term "stent fracture" refers to the fracturing of the Melody TPV. However, in subjects with multiple stents in the RVOT it is difficult to definitively attribute stent fractures to the Melody frame versus another stent.*

For additional information, please refer to the Instructions For Use provided with the product.

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician.

Melody is a registered trademark of Medtronic.

Medtronic
Further, Together



Medical Director and Staff Level Pediatric Cardiovascular Critical Care Physicians General Pediatric Cardiologist Pediatric Cardiac Interventionalist Geneticist

Medical City Children's Hospital has an unwavering focus on patient care and offers world-renowned excellence in comprehensive pediatric services. Since 1996, our specialists haven't let anything distract them from serving children. As a result, we've helped thousands of children from more than 75 countries. We are a comprehensive children's hospital with specialists in virtually every pediatric subspecialty. Medical City is the only facility in north Texas where fetal diagnosis, maternal, neonatal and pediatric transport, high risk delivery stabilization in the NICU, corrective surgery, state of the art postoperative monitoring and care and long term follow-up of patients with complex congenital heart disease can all be delivered under one roof.

The Congenital Heart Surgery Unit (CHSU) accommodates around 400 children annually who undergo heart operations performed by Dr. Eric Mendeloff. 30% of our cases are neonates and 58% are under the age of 2 years. Cases range in complexity from palliation of hypoplastic left heart syndrome to closure of atrial and ventricular septal defects. Highly specialized care in the CHSU is provided by subspecialty-trained physicians and an excellent group of long term nurses and respiratory therapists. This focus on pediatric cardiac critical care has resulted in superlative patient outcomes that exceed national norms. The heart program's success has attracted referrals from across the country. With the addition of a second Congenital Heart Surgeon to our already robust program, we anticipate growth that will require a sixth member for our CICU team in addition to our need for a Medical Director of the Unit. Preferred candidate for the director level position will possess leadership attributes with evidenced experience, along with a strong clinical skill set.

All candidates are preferred to be BC/BE in Pediatric Cardiology and Pediatric Critical Care or boarded in one of these with additional training in Pediatric Cardiac Critical Care. Those with certification in one discipline and solid experience in the alternate subspecialty should also apply. Positions are employed and offer a competitive salary and excellent benefits packet.

Our hospital has immense current capabilities and is positioned to grow.

Kathy Kyer
National Director of Pediatric Subspecialty Recruitment
Kathleen.Kyer@HCAHealthcare.com
937.235.5890

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share? Submit your manuscript to:
RichardK@CCT.bz



Figure 2. Suprasternal view showing retroaortic left innominate vein venous flow from the left side of the neck passing below the aorta to the right side.

plexi above and below the fourth aortic arch. Normally, the aortic arch shortens during the embryological development and occupies the space of the inferior transverse capillary plexus, thus causing its regression. The rest of the venous blood shunts into the superior transverse capillary plexus and forms the normal supraaortic course of the left innominate vein.² In contrast, reduced shortening of the aortic arch (right-sided or high aortic arch) may cause regression of the superior capillary plexus and results in preservation of the inferior capillary plexus and formation of the retroaortic vein (Figure 4). Abnormal development of the pulmonary arteries (pulmonary atresia or pulmonary stenosis) encourages the sparing of the lower dorsal plexus, leading to formation of an anomalous innominate vein.¹

The retroaortic innominate vein can usually be traced by echocardiography in the suprasternal long axis view to the left side of neck when the transducer is tilted leftwards and anteriorly, Doppler color flow and Doppler spectral analysis shows low velocity venous flow into the superior vena cava.³ An aorto-pulmonary collateral and PDA arising from the aortic arch show a high velocity continuous flow.

Usually, the retroaortic innominate vein in isolation has no clinical importance. The anomalous innominate vein may cause technical difficulties during pacemaker insertion or central venous line placement




Figure 3. CT angiogram, Figure 3a-lateral view and Figure 3b-3D reconstruction showing hypoplastic left aortic arch, juxtaductal coarctation of aorta, patent ductus arteriosus and low origin of left subclavian artery just proximal to the coarctation site.

through the left arm approach. For patients undergoing cardiac surgery, the superior vena caval cannulation for cardiopulmonary bypass has to be done more caudally than usual to avoid obstruction of the retroaortic innominate vein.¹ The anomaly may complicate exposure of the pulmonary arteries while creating systemic vein to

pulmonary artery anastomosis during Glenn shunt. Also, it may obscure the surgical field in the construction of a subclavian to pulmonary artery shunt and the ligation of a patent ductus arteriosus.⁴ Retroaortic innominate vein has been used in pulmonary artery reconstruction by creating a wide side-to-side anastomosis.⁵



Volunteer / Get Involved
www.chimsupport.com

HOW WE OPERATE

The team involved at C.H.I.M.S. is largely a volunteering group of physicians, nurses and technicians who are involved in caring for children with congenital heart disease.

The concept is straightforward. We are asking all interested catheter laboratories to register and donate surplus inventory which we will ship to help support CHD mission trips to developing countries.




Figure 4. Schematic diagrams illustrating the development of retroaortic left innominate vein (RLIV) (a) In the sixth week of gestation: 1 - right anterior cardinal vein; 2 - left anterior cardinal vein; 3 - primitive aorta; 4 - superior transverse capillary plexus; 5 - inferior transverse capillary plexus; 6 - common cardinal veins; 7 - right posterior cardinal vein; 8 - left posterior cardinal vein; 9 - sinus venosus; 10 - vitelline vein; 11 - umbilical vein; 12 - developing inferior vena cava. (b) Retroaortic left innominate vein: 1 - innominate vein; 2 - azygos vein; 3 - superior vena cava; 4 - right atrium; 5 - inferior vena cava; 6 - coronary sinus; 7 - ascending aorta.

References

1. Chen SJ, Liu KL, Chen HY, Chiu IS, Lee WJ, Wu MH, et al. Anomalous brachiocephalic vein: CT, embryology and clinical implications. *Am J Roentgenol* 2005;184:1235-40.
2. Kulkarni S, Jain S, Kasar P, Garekar S, Joshi S. Retroaortic left innominate vein-Incidence, association with congenital heart defects, embryology, and clinical significance. *Ann Pediatr Card* 2008;1:139-41 and Sukulal K, Bijulal S, Tharakan JA. Differential diagnosis of vascular structures in relation to upper ascending aorta: The retro-aortic innominate vein. *Ann Pediatr Card* 2013;6:97-8.
3. Choi JY, Jung MJ, Kim YH, Noh CI, Yun YS. Anomalous subaortic position of the brachiocephalic vein (innominate vein): An echocardiographic study. *Br Heart J* 1990;64:385-7.
4. Gerlis LM, Ho SY. Anomalous subaortic position of the brachiocephalic (innominate) vein: A review of published reports and report of three new cases. *Br Heart J* 1989;61:540-5.
5. Agrawal R, Krishnan GS, Kulkarni S, Bhatt K, Cherian KM. Novel use of a retroaortic innominate vein in cavopulmonary anastomosis. *J Thorac Cardiovasc Surg* 2006;132:166-7.
6. Konstantinov IE, Van Arsdell GS, O'Blenes S, Roy N, Campbell A. Retroaortic

innominate vein with coarctation of the aorta:surgical repair and embryology review. *Ann Thorac Surg*.2003 Mar;75(3):1014-6.

CCT

Corresponding Author



Siddharth Patel, MD
Pediatric Cardiology Fellow
Department of Pediatrics
Division of Cardiology
Louisiana State University School of Medicine
Children's Hospital of New Orleans, Louisiana
Phone: 567.225.2082; Fax: 504.896-3952
spat16@lsuhsc.edu

mailing address:
5315 Citrus Blvd, Apt #D323
River Ridge, LA 70123

Steffan Sernich, MD
Associate Professor of Pediatrics
Department of Pediatrics
Division of Cardiology
Louisiana State University School of Medicine
Children's Hospital of New Orleans, Louisiana

Pooja Makhija, MD
Staff Physician Pediatric ER
Department of Pediatrics
Division of Pediatric ER
Children's Hospital of New Orleans, Louisiana



Pediatric Cardiologist

The Department of Pediatrics of the University of Texas Health Science Center at San Antonio is seeking a pediatric cardiologist with advanced training and expertise in interventional cardiology to join the faculty of the Division of Pediatric Cardiology. The University of Texas Health Science Center at San Antonio partners with University Health System to provide comprehensive care for children throughout South Texas.

Our congenital heart program includes a dedicated pediatric catheterization laboratory, a full-service pediatric echocardiography laboratory and a twelve bed variable acuity pediatric cardiac inpatient unit.

The candidate must be fellowship trained, board certified/board eligible in pediatric cardiology, and either possess or be able to easily obtain an unrestricted Texas medical license. The candidate will join an established academic clinical practice with 4 pediatric cardiologists and 2 congenital heart surgeons.

The School of Medicine has 230 medical students at each level. Cardiology faculty are engaged in the training of these medical students and 35 pediatric residents.

Please submit a letter of interest, curriculum vitae, and three letters of recommendation to:

Steven R. Neish, M.D., S.M.,
Professor and Vice Chairman,
Department of Pediatrics,
Chief, Pediatric Cardiology,
The University of Texas Health Science
Center at San Antonio,
7703 Floyd Curl Drive
San Antonio, Texas 78229-3900.

The University of Texas Health Science Center at San Antonio is an Equal Employment Opportunity/Affirmative Action Employer including protected veterans and persons with disabilities. All faculty appointments are designated as security sensitive positions.

CHiP Network Update

By Gary Webb, MD

The basic idea of the CHiP Network (the Congenital Heart Professionals Network) is that congenital heart and pediatric cardiac professionals can subscribe to the service, indicating the type of information they would like to receive. This builds a list that is global, and that can be used by approved/appropriate organizations to communicate with professional members of this community. As a result, these organizations will be able to communicate with a larger readership. There is no charge for the service.

The CHiP Network has the capacity to send information only to certain types of professionals, in certain countries or cities, and to subscribers requesting messages in languages other than English. The potential of the system is still being explored. The CHiP Network is all-inclusive and is comprised of everyone who considers themselves a congenital heart or pediatric cardiac professional or administrator, including:

- Pediatric cardiologists
- ACHD cardiologists
- Cardiac surgeons
- RNs and APNs
- Physician assistants
- Cardiac care associates
- Trainees/fellows
- Administrators
- Social workers
- Psychologists and mental health professionals
- Researchers/scientists
- Transition medicine specialists
- Intensivists
- Anesthetists
- and Industry representatives.

This will make it possible for these various groups to communicate with people who identify themselves as being in the same profession. Of course, individuals in those groups will need to initiate the communications and consider how best to do this.

This will enable the circulation of various types of information:

- pending events
- *Journal Watch*
- recruitment opportunities

Geographic Distribution by Continent		
Continent	#	%
North America	1720	60%
Europe	420	15%
Asia-Pacific	335	12%
Middle East	171	6%
Central & South America	127	4%
Africa	106	4%

Profession Distribution		
Work Name Type	#	%
Cardiologist	1420	58%
Cardiac Care Associate / PA / RN	308	13%
Cardiac Surgeon	231	9%
Cardiac Anesthetist	120	5%
Nurse Practitioner	102	4%
Cardiac Intensivist	95	2%
Echo Technician	56	2%
Researcher / Scientist	54	2%
Administrator	49	2%
Industry Representative	28	1%

- research opportunities
- educational resource announcements
- news stories
- and professional society news reports and membership recruitment.

Undoubtedly, other applications will be developed.

To this point, the CHiP Network has circulated a monthly newsletter and a monthly *Journal Watch* service to its subscribers. The *Journal Watch* includes abstracts from the previous month for six different subspecialty areas: pediatric cardiology; interventional CHD; EP CHD; CHD surgery; fetal cardiology; and ACHD cardiology. The section editors identify the most important abstracts for the month, and offer a featured commentary on them. *Journal Watch* is housed on the ACHD Learning Center, so users can examine abstracts on a monthly basis going back 2-3 years.

CHiP's challenge, and the challenge of congenital heart professionals with an interest in the success of this resource, is to grow the network of subscribers and to

“The CHiP Network has the capacity to send information only to certain types of professionals, in certain countries or cities, and to subscribers requesting messages in languages other than English. The potential of the system is still being explored.”



Volunteer / Get Involved
www.chimsupport.com

HOW WE OPERATE

The team involved at C.H.I.M.S. is largely a volunteering group of physicians nurses and technicians who are involved in caring for children with congenital heart disease.

The concept is straightforward. We are asking all interested catheter laboratories to register and donate surplus inventory which we will ship to help support CHD mission trips to developing countries.

Table 3. Geographic Distribution by County Distribution

Countries		Countries		Countries		Countries	
United States	1485	Malaysia	16	Greece	4	Zambia	2
Canada	240	Belgium	15	Poland	4	Libyan Arab Jamahiriya	2
United Kingdom	137	Denmark	14	Slovakia	4	Cameroon	2
Australia	94	China	14	Slovenia	4	Netherland Antilles	2
Saudi Arabia	58	Israel	13	Sudan	4	Bolivia	1
South Africa	50	Korea, Republic of	13	Ghana	4	Cuba	1
Brazil	46	Turkey	12	Ethiopia	4	El Salvador	1
India	46	Venezuela	10	Morocco	4	Panama	1
Netherlands	37	Colombia	10	Bangladesh	4	Guatemala	1
Iran, Islamic Republic of	28	United Arab Emirates	10	Unspecified	4	Ecuador	1
Sweden	28	France	10	Costa Rica	3	Syrian Arab Republic	1
Argentina	24	Pakistan	10	Trinidad and Tobago	3	Iceland	1
Egypt	24	Philippines	10	Iraq	3	Albania	1
Switzerland	24	Qatar	9	Oman	3	Malta	1
Japan	24	Hungary	9	Bulgaria	3	Croatia	1
Germany	21	Russian Federation	9	Finland	3	Senegal	1
New Zealand	21	Austria	7	Tanzania, United Republic of	3	Zimbabwe	1
Thailand	21	Chile	6	Singapore	3	Mozambique	1
Italy	20	Kenya	6	Nepal	3	Georgia	1
Norway	20	Lithuania	5	Mongolia	3	Uzbekistan	1
Indonesia	19	Ukraine	5	Jamaica	2	Azerbaijan	1
Nigeria	17	Portugal	5	Bahrain	2	French Polynesia	1
Taiwan	17	Uganda	5	Jordan	2	Brunei Darussalam	1
Mexico	16	Hong Kong	5	Ireland	2	Bhutan	1
Spain	16	Kuwait	4	Cyprus	2		

“CHiP’s challenge, and the challenge of congenital heart professionals with an interest in the success of this resource, is to grow the network of subscribers and to enhance the value of the service.”

enhance the value of the service. CHiP has over 25 partner organizations. Each of these is asked to invite its members to join CHiP, and to show informational material about CHiP at their meetings.

To this point, we are grateful that financial support has been provided by Cincinnati Children’s Hospital Heart Institute. We are also introducing a new category of partners, Institutional Partners. The first of these to be identified is Toronto’s Hospital for Sick Children, with 58 subscribers to date.

Join CHiP - membership is free - www.chipnetwork.org

CCT



*Gary Webb, MD
Cincinnati Adult Congenital Heart Program
The Heart Institute at Cincinnati Children’s Hospital Medical Center
Ph: 513-803-4244; Cell: 215-313-8058
Fax: 513-803-0079
gary.webb@cchmc.org*



The Anthony Bates Foundation

www.AnthonyBates.org

Anthony Bates Foundation

428 E. Thunderbird Rd., # 633
Phoenix, AZ 85022
Tel: 602-482-5606

Sharon Bates
sharon@anthonybates.org

Education and awareness of the leading killer of young people, HCM, Hypertrophic Cardiomyopathy, through organizing and training in cardiac screening programs for young people everywhere.



PEDIATRIC ECHOCARDIOGRAPHY

OCHSNER HEALTH SYSTEM in **NEW ORLEANS** is searching for a **BC/BE PEDIATRIC CARDIOLOGIST** to join the Section of Pediatric Cardiology. The section is recruiting a **Pediatric Cardiologist specializing in non-invasive imaging with expertise in fetal echo, TEE, MRI, and/or Cardiac CT** to join 3 other imaging physicians in a group of 9 pediatric cardiologists. Completion of an advanced non-invasive imaging fellowship is preferred. Salary with be commensurate with training and experience.

The Department of Pediatrics and our pediatric subspecialists (collectively, Ochsner for Children™) represent over 110 physicians. Our pediatric subspecialties include: Allergy, Cardiology, CV Surgery, Critical Care Medicine, Dermatology, ENT, Endocrinology, Emergency Medicine, Gastroenterology, Hematology/Oncology, Hospital Medicine, Infectious Diseases, Medical Genetics, Neonatology, Nephrology/Hypertension, Neurology, Neurosurgery, Ophthalmology/Optomety, Orthopedics, Psychiatry/Sports Medicine, Plastic and Reconstructive Surgery, Psychiatry, Pulmonology, General Surgery, Transplant, and Urology. We also have a Pediatric Anesthesiology Team, Pediatric CV Surgery Anesthesia Team and Pediatric Radiology support. We support a pediatric heart failure program, including left ventricular assist devices and transplantation. We also have a pediatric transcatheter valve program. We utilize a 54 bed Level III-C NICU and 14 bed PICU as well as a dedicated, full-time, region-wide in-house pediatric transport team.

Ochsner is the region's leading integrated provider of multi-specialty care for infants, children and adolescents. Our physicians practice in multiple sites across Louisiana, including a large, state-of-the-art dedicated pediatric ambulatory facility on our main campus. We have an outstanding primary care pediatric network across the region with over 40 general pediatricians, in addition to a large outside referral base. Ochsner pediatric physicians care for over 55,000 unique pediatric patients annually, and Ochsner hospitals have more than 4,000 deliveries per year.

Ochsner Health System is Louisiana's largest non-profit, academic, healthcare system. Driven by a mission to Serve, Heal, Lead, Educate and Innovate, coordinated clinical and hospital patient care is provided across the region by Ochsner's 25 owned, managed and affiliated hospitals and more than 50 health centers. Ochsner is the only Louisiana hospital recognized by U.S. News & World Report as a "Best Hospital" across six specialty categories caring for patients from all 50 states and more than 90 countries worldwide each year. Ochsner employs more than 17,000 employees, nearly 1,000 physicians in over 90 medical specialties and subspecialties, and conducts over 750 clinical research studies. For more information, please visit www.ochsner.org and follow us on Twitter and Facebook.

Please email curriculum vitae to:
profrecruiting@ochsner.org

Ref # PECHO4,
or call (800) 488-2240 for more information.

Sorry, no opportunities for J-1 applications exist at present.

Ochsner is an equal opportunity employer and all qualified applicants will receive consideration for employment without regard to race, color, religion, sex, national origin, sexual orientation, disability status, protected veteran status, or any other characteristic protected under law.



PEDIATRIC CARDIOLOGIST

OCHSNER HEALTH SYSTEM is recruiting a **BC/BE GENERAL PEDIATRIC CARDIOLOGIST** to join the Section of Pediatric Cardiology at our newest multi-specialty location in **WEST MONROE, LA**. The successful candidate will practice in West Monroe as a member of a local group of 2 pediatric cardiologists, closely affiliated with the senior cardiologist on-site and with the Ochsner Section of Pediatric Cardiology, currently comprised of 9 pediatric cardiologists. Salary is commensurate with training and experience.

The Department of Pediatrics and pediatric subspecialists (collectively, Ochsner for Children™) is comprised of 110 physicians with specialties including: General Pediatrics, Allergy, Cardiology, CV Surgery, Critical Care Medicine, Dermatology, ENT, Endocrinology, Emergency Medicine, Gastroenterology, Hematology/Oncology, Hospital Medicine, Infectious Diseases, Medical Genetics, Neonatology, Nephrology/Hypertension, Neurology, Neurosurgery, Ophthalmology/Optomety, Orthopedics, Psychiatry/Sports Medicine, Plastic and Reconstructive Surgery, Psychiatry, Pulmonology, General Surgery, Transplant, and Urology.

Congenital Cardiovascular Program support includes a pediatric CV Surgery Anesthesia Team, perfusion and pediatric radiology. We support comprehensive catheter intervention and EP services including a pediatric transcatheter valve program and a pediatric heart failure program that includes percutaneous and surgical circulatory assist devices and transplantation. We utilize a 54 bed Level III-C NICU and 14 bed PICU as well as a dedicated, full-time, region-wide in-house pediatric transport team.

Ochsner is the region's leading integrated provider of multi-specialty care for infants, children and adolescents. Our physicians practice in multiple sites across Louisiana, including a large, state-of-the-art dedicated pediatric ambulatory facility on our main campus. We have an outstanding primary care pediatric network across the region with over 40 general pediatricians, in addition to a large outside referral base. Ochsner pediatric physicians care for over 55,000 unique pediatric patients annually, and Ochsner hospitals have more than 4,000 deliveries per year.

Ochsner Health System is Louisiana's largest non-profit, academic, healthcare system. Driven by a mission to Serve, Heal, Lead, Educate and Innovate, coordinated clinical and hospital patient care is provided across the region by Ochsner's 25 owned, managed and affiliated hospitals and more than 50 health centers. Ochsner is the only Louisiana hospital recognized by U.S. News & World Report as a "Best Hospital" across six specialty categories caring for patients from all 50 states and more than 90 countries worldwide each year. Ochsner employs more than 17,000 employees, nearly 1,000 physicians in over 90 medical specialties and subspecialties, and conducts over 750 clinical research studies. For more information, please visit www.ochsner.org and follow us on Twitter and Facebook.

Please send curriculum vitae to:
profrecruiting@ochsner.org,
Ref. # PCARD01,
or call (800) 488-2240 for more information.

Sorry, no opportunities for J-1 applications exist at present.

Ochsner is an equal opportunity employer and all qualified applicants will receive consideration for employment without regard to race, color, religion, sex, national origin, sexual orientation, disability status, protected veteran status, or any other characteristic protected by law.

A Case Series of Unfortunate Gerbode Defects

By Lucky R. Cuenza, MD; Anna Christina Adora, MD; Thessie Minelli O. Valdez, MD; Cheryl P. Fomaneg, MD; Efren R. Vicaldo, MD

Introduction

Gerbode Defect is a very rare congenital anomaly described as a communication between the left ventricle and the right atrium. It was named after Frank Gerbode, a surgeon from Stanford University who described his surgical findings in 1958. He reported two types (direct and indirect) depending if the defect lies above or below the tricuspid valve along with the presence of a true ventriculoatrial communication.¹ The pathology may be due to a congenital defect, or can result from trauma, myocardial infarction, endocarditis or even aortic valve replacement.² It can appear as an isolated lesion, but it has been reported to accompany other congenital lesions. We describe 3 cases of patients who had Gerbode Defect along with other findings noted on echocardiography also with other findings not previously reported.

Case 1

A 31-year-old female came in due to frequent symptoms of palpitations of five months duration accompanied by dyspnea on exertion. Physical examination showed a blood pressure of 100/60 with a heart rate of 82 beats per minute. Auscultation of the chest revealed a 4/6 holosystolic murmur at the 4th intercostal space left sternal border with a thrill. Doppler studies showed a mosaic color flow display going from the left ventricle to the left atrium (Figure 1). This was confirmed by transesophageal echocardiogram (Figure 2). Peak systolic gradient between the left ventricle and the right atrium was measured at 89mmHg. There was no chamber dilatation noted. The patient was started on diuretics and eventually underwent surgical closure of the defect. She was discharged stable and improved.

Case 2

A 25-year-old female was referred to our institution for difficulty of breathing, exertional dyspnea and bipedal edema. Physical examination revealed a 4/6 holosystolic murmur at the 3rd left intercostal space. Doppler color flow on transthoracic echocardiogram showed an abnormal color flow display across the perimembranous portion of the interventricular septum during systole from the left ventricle to the right atrium. The defect was noted above the tricuspid valve (Figure 3). There was dilatation of the right atrium and the right

pulmonary artery with moderate tricuspid regurgitation. There was also note of Patent Ductus Arteriosus (PDA) with a bidirectional shunt (Figure 4). She was started on Furosemide 40mg once a day which relieved her symptoms. She was initially scheduled for transesophageal echo and further evaluation, but was lost to follow-up.

Case 3

A 29-year-old female diagnosed with Congenital Heart Disease since 9-years-old complained of palpitations and easy fatigability a week postpartum. Physical examination showed a blood pressure of 130/80 and tachypnea at 32 cycles per minute with an oxygen saturation of 95% at




Figure 1. Transthoracic echocardiography apical four chamber view showing color flow directed towards the right atrium from the left ventricle (left and middle). Short axis view at the level of the aortic valve shows the jet situated above the tricuspid valve.




Figure 2. Transesophageal echocardiography mid esophageal short axis view showing an echo dropout at the ventriculoatrial septum which measures around 0.45cm (left). The flow is directed towards the right atrium from the left ventricle (right).




Figure 3. Transthoracic echocardiogram at the short axis view shows the ventriculoatrial communication on Color flow Doppler. Apical four chamber view shows a mosaic color flow from the left ventricle to the right atrium situated above the valve.

room air. She was tachypneic at 32 cycles per minute with an oxygen saturation of 95% at room air. She was tachycardic, but with regular rhythm, with right ventricular heave, single S2, with 4/6 holosystolic murmur at the 2nd left intercostal space, (+) thrill, a 4/6 holosystolic murmur at the 4th left intercostal space and positive for Carvallo's sign. There was electrocardiographic and x-ray evidence of right atrial enlargement and right ventricular hypertrophy. There was a suspicious mosaic color flow during systole from the left ventricle to the right atrium on transthoracic echocardiogram. Two dimensional and 3D transesophageal echocardiogram confirmed the findings of a direct Gerbode Defect above the tricuspid valve (Figure 5). Transthoracic echocardiogram showed a Patent Ductus Arteriosus (Figure 6) with severe pulmonary hypertension. The left atrium, right atrium, main pulmonary artery and right ventricular had signs of volume and pressure overload and there was mitral valve prolapsed with moderate regurgitation. Additional findings included a mild valvar pulmonic stenosis (Figure 7) with a peak gradient of 20 mmHg. A flap-like structure was likewise noted at the interatrial septum suggestive of Patent Foramen Ovale (PFO). Contrast study demonstrated passage of microbubbles from the RA to the LV and LA and from the PA to the descending aorta indicative of the presence of Gerbode Defect, PFO and PDA, respectively.

Discussion

Gerbode defects are very rare abnormalities with an incidence of 0.08% of congenital defects.³ It is primarily caused by an anatomic deficiency of the membranous septum which can be either congenital or acquired. Riemenschneider and Moss devised a classification⁴ based on the anatomical relationship of the left ventricle to the right atrium shunt with the tricuspid valve. This anatomical arrangement allows the septal leaflet to divide the membranous ventricular septum into two portions: a supravulvar portion, and an infravalvular portion. A defect in the supravulvar portion results in a direct left ventricle to right atrium communication and a defect below the tricuspid valve represents a perimembranous ventricular septal defect and would normally result in a communication between the left ventricle and the right ventricle then through a defective tricuspid valve into the right atrium. This is termed as an indirect Gerbode Defect. The true (or direct) Gerbode Defect is a true left ventricular to right atrial communication and is rarer than the indirect type.⁵ All the patients in our series had the true or direct form. In both defects left ventricular to right atrium communication allows shunting of blood to the right atrium during systole. If the communication is large enough, the patient may become symptomatic, and may present with signs of volume overload.⁶

Typical findings are similar to that of a ventricular septal defect, with a harsh systolic murmur accompanied by a thrill. A disproportionate enlargement of the right atrium may be found on radiograph along with features of right atrial abnormality on electrocardiogram.⁷ However, definitive diagnosis may not be possible based on these methods only. Two of the cases also presented with other shunt abnormalities, as well as varying degrees of pulmonary hypertension, making the diagnosis clinically challenging.

Echocardiography identifies the location of the defect and color flow imaging can identify the flow pattern from the left ventricle to the right atrium. It is also used to locate the relationship of the defect with respect to the tricuspid valve. Transesophageal echocardiography can reliably distinguish the two types and was



Florida Hospital for Children

Pediatric Inpatient Cardiologist Orlando, Florida

The Walt Disney Pavilion at Florida Hospital for Children is looking for a Pediatric Cardiologist with primary inpatient care focus to help further develop our open heart program in Orlando. This physician will lead the cardiology portion of the pediatric cardiac intensive care team at Florida Hospital in Orlando as part of the multi-site John's Hopkins/All Children's Heart Surgery program.

Responsibilities of the new physician will include: cardiology management of all patients in our 10 bed pediatric CICU, hospital inpatient and NICU consults, post-op and post procedure management, performing TEEs and reading echocardiograms. Some outpatient clinic time is also possible depending on physician's level of interest. We have an active Extracorporeal Membrane Oxygenation (ECMO) Program, a pediatric residency program, and clinical research opportunities.

Working with an expanding and committed children's hospital, a collaborative relationship with Johns Hopkins and All Children's, an established and successful pediatric cardiology group with a strong referral base, a dedicated inpatient unit, a family-centered practice atmosphere, the ability to build and shape a program, and a competitive benefits / compensation package.

Requirements of the position:

- Diplomate (or board eligible) of the American Board of Pediatric Cardiology, extra training in cardiac intensive care is preferred.
- Experience and proficiency with TEE's and ECHO's.
- Strong leadership and communication skills.
- "Team player" mentality.

Contact:

Jason Junker
Director, Physician Recruitment
Phone: (407) 200-2538
Jason.Junker@FLHosp.org

<https://www.floridahospital.com/children>

MD 1 World

CONNECTING THE GLOBAL MEDICAL WORLD

Collaborate, Learn & Engage

We Empower Doctors From Around the Globe to Save Lives

WWW.MD1WORLD.COM



**We are committed to the
lifetime management of
congenital heart disease.**

Transcatheter and Surgical
Heart Valves

RVOT Conduits

Ablation Technologies

ICDs

Oxygenators and Filters

Cannulae

Pacemakers

Pulse Oximetry Monitoring
for CCHD Screening

3rd Generation PFO, ASD,
and PDA Occluders*

Cerebral/Somatic Monitoring

*These products are not
available in the US.

Melody-TPV.com

Medtronic | Minneapolis, MN 55432-5604

Toll-free: 1 (800) 328-2518

UC201601683 EN ©2015 Medtronic.

All rights reserved. 08/2015

INNOVATIVE TECHNOLOGIES. EVERY STEP OF THE WAY.

Medtronic
Further, Together




Figure 4. Transthoracic echocardiogram short axis view at the level of the pulmonary artery bifurcation tract showing a connection between the pulmonary artery and the aorta.




Figure 5. Transesophageal Echocardiography mid-esophageal short axis view showing an echo drop out at the ventriculoatrial septum between the LVOT and right atrium measuring 0.46 cm in diameter (left) with mosaic color flow display across the ventriculoatrial septum and tricuspid valve. 3D TEE mid-esophageal short axis view showing an echo drop out at the ventriculoatrial septum between the LVOT and right atrium with mosaic color flow display across the ventriculoatrial septum and tricuspid valve.




Figure 6. Transthoracic Echocardiography showing the pulmonic valve in parasternal short axis view with systolic doming motion (left) and continuous flow from the aorta to the pulmonary artery (right).




Figure 7. Transesophageal Echocardiography showing the pulmonic valve in mid-esophageal long axis view during diastole (left) and mosaic color flow display across the pulmonic valve during systole (middle). 3D TEE showing the pulmonic valve in mid-esophageal short axis view during systole (right).

also found to impact operative repair.⁸ One of the hallmarks of Gerbode ventriculoatrial defect is a high Doppler gradient between the left ventricle and the right atrium accompanied by right atrial dilation. The echocardiographer must be vigilant as the jet from the Gerbode Defect may be confused with a tricuspid regurgitant jet and can be

CHIP NETWORK

CONGENITAL HEART PROFESSIONALS

WHAT IS THE CHIP NETWORK? - The CHIP Network, the Congenital Heart Professionals Network, is designed to provide a single global list of all CHD-interested professionals in order to:

- Connect pediatric and adult CHD-interested professionals to events, conferences, research opportunities and employment
- Keep members up with the literature through the monthly *Journal Watch* service
- Increase education and provider awareness of new developments
- Bring the pediatric and adult congenital heart communities into closer contact
- Offer a communication tool for critical issues

WHO SHOULD PARTICIPATE? - The CHIP Network is all inclusive and is comprised of everyone who considers themselves a congenital heart professional or administrator, including: Pediatric cardiologists, ACHD cardiologists, RNs and APNs, Cardiac surgeons, Cardiac care associates, Trainees/fellows, Administrators, Psychologists and Mental health professionals, Researchers/scientists, Intensivists, Anesthetists, Industry representatives

OUR SUPPORTING PARTNERS:

- Adult Congenital Heart Association
- Asia Pacific Society for ACHD
- Children's Hospital of Philadelphia Cardiology meeting
- Cincinnati Children's Hospital
- Congenital Cardiology Today (official publication of the CHIP Network)
- Congenital Heart Surgeons Society
- ISACHD
- Japanese Society of ACHD
- Johns Hopkins All Children's Heart Institute
- North American ACHD program
- Paediatric Cardiac Society of South Africa
- Pan Arab Congenital Heart Disease Association
- PCICS
- PICS
- Specialty Review in Pediatric Cardiology
- World Congress of Pediatric Cardiology and Cardiac Surgery

JOIN US - Membership is Free!

The CHIP Network management committee invites the participation of other organizations who want to communicate with all or some of the congenital heart professionals on this list. Please contact Dr. Gary Webb (gary.webb@cchmc.org) to ask that your organization's or institution's name be added to the list of partner organizations.

Register at: www.chipnetwork.org.



Funded by Cincinnati Children's Heart Institute

mistakenly associated with pulmonary hypertension.⁹ The presence of normal diastolic pulmonary arterial pressure using pulmonic regurgitation jet is very useful to distinguish the true pulmonary arterial hypertension from high velocity jet in the right atrium caused by Gerbode-type defect. Our second patient had a bidirectional shunt with severe pulmonary hypertension and the tricuspid regurgitation jet was delineated properly from the jet coming from the Gerbode Defect. Our third patient had multiple congenital heart defects. The Gerbode Defect was only suspected on transthoracic echo, while integrating clinical findings of right-sided volume overload. Transesophageal echocardiogram revealed the Gerbode Defect, to go along with the patent ductus arteriosus, a patent foramen ovale and severe pulmonary hypertension. Other imaging modalities such as 3D transesophageal echocardiogram may add incremental diagnostic value by delineating the morphologic nature of the ventricular septal defect and its shape, and showing other structures in their realistic spatial distribution.¹⁰ This may have implications in surgical planning.

Surgical repair is indicated for symptomatic patients with significant-sized shunts and is usually associated with favorable outcomes.¹¹ Gerbode Defect may or may not be associated with other cardiac abnormalities. In the study by Anderson et al they noted concomitant congenital anomalies in six children. One patient had a right aortic arch, one patient had a patent ductus arteriosus, three patients had a left superior vena cava, two patients had an anomalous left hepatic vein.¹² In this series, the first patient had an isolated Gerbode Defect with good functional capacity. The second patient similarly had a patent ductus arteriosus with a bidirectional shunt as well as moderate tricuspid regurgitation and severe pulmonary hypertension. Our third patient had multiple congenital defects. Aside from the Gerbode Defect, she also had a Patent Ductus Arteriosus with a PFO, with severe tricuspid regurgitation and mild supralvalvar pulmonary stenosis. The findings from the latter two patients have not been previously described in literature. Mechanistically the shunt from the Gerbode Defect may contribute to more right-sided volume overload and pulmonary hypertension. The study by Anderson et al was done in children, and had no reported operative or late mortality.¹² The outcomes in patients with multiple and advanced concomitant congenital heart disease is uncertain. We recommended surgical closure for our first patient and further work up

(hemodynamic studies) on the latter two cases to determine operative prognosis.

Conclusion

Gerbode's defect is a very rare congenital anomaly which can occur in isolation or may be associated with other congenital heart defects. It may complicate these conditions and even lead to misinterpreted diagnosis. We reported three cases in our institution, which had a direct type of ventriculoatrial communication. These cases underlie the importance of integrating sound clinical assessment, as well as meticulous echocardiographic investigation in order to properly diagnose this unusual anomaly, facilitating surgical planning and determination of optimum management.

References

1. F. Gerbode, H. Hultgren, D. Melrose, and J. Osborn, Syndrome of left ventricular-right atrial shunt. Successful repair of defect in five cases, with observation of bradycardia on closure, *Annals of Surgery*, vol. 148, pp. 433–446, 1958.
2. Pursnani AK, Tabaksblat M, Saric M, Perk G, Loulmet D, Kronzon I. Acquired Gerbode defect after aortic valve replacement, *J Am Coll Cardiol* 2010;55(25):e145.
3. Silbiger JJ, Kamran M, Handwerker S, et al: The Gerbode defect: Left ventricular to right atrial communication-anatomic,

“These cases underlie the importance of integrating sound clinical assessment, as well as meticulous echocardiographic investigation in order to properly diagnose this unusual anomaly, facilitating surgical planning and determination of optimum management.”

MEDICAL MEETING FOCUS

2nd Annual Basic & Advanced Fetal Cardiac Symposium and Workshop
Sep. 10-12, 2015; Chicago, IL
www.FetalCardiacSymposium.com

This symposium will discuss various concepts and common CHD malformations encountered in the fetus, with emphasis on imaging and practical tips for diagnosis. In particular, it will include sessions focusing on the following areas:

- Fetal Cardiovascular Physiology and Pathophysiology
- Latest Guidelines in Prenatal Screening and Red Flags in the Diagnosis of Fetal CHD
- Live Demonstration on Performing a Fetal Echocardiogram
- Practical Tips in the Diagnosis of Commonly Encountered Cardiac Defects
- Evaluation of Fetuses with Borderline Cardiac Findings
- Diagnosis and Management of Fetal Arrhythmia
- Updates on Fetal Genetic Evaluation

The symposium will provide the audience with the unique experience of hands-on and live demonstration sessions, as well as opportunities for the audience to discuss interesting cases and interact with experts in the field. Registrants will have the chance to submit interesting cases to be presented during the course of the symposium.

Hands-on Scanning Workshop: There will be two separate sessions focusing on scanning the fetal heart under the supervision of experienced faculty and using various up-to-date machines.

Participants become familiar with: Identifying Situs; Obtain the 4-chamber View; Demonstrate Pulmonary Venous Flow; Obtain the Right and Left Ventricular Arterial Outflow Tracts View; Obtain the 3-vessel View; Obtain the Arch Views.

Sponsored for CME credit by Rush University Medical Center which designates this live activity for a maximum of 19.25 AMA PRA Category 1 Credit(s)[™] & 19.25 Continuing Nursing Education credit(s). This activity is also approved by the Society for Diagnostic Medical Sonography (SDMS), and is eligible for up to 19.25 SDMS credits.

Course Director: Karim A. Diab, MD

Invited and Local Faculty: Ra-id Abdulla, MD; Alfred Abuhamad, MD; Emerio T. Alboliras, MD; Sawzan M. Awad, MD, MSC; John Bokowski, PhD, RDCS, FASE; Massimo Caputo, MD, Mch, FRCS; Lisa Hornberger, MD; Edgar Jaeggi, MD; Carolyn Jones, MD, PhD; Michelle Rexilius, RDCS; Mark Sklansky, MD; Wayne Tworetzky, MD; Xavier Pombar, DO; Luciana T. Young, MD



Archiving Working Group
International Society for Nomenclature of
Paediatric and Congenital Heart Disease
ipccc-awg.net

Save
the Date

Evolution or Revolution?

Disruptive Thinking, Technology and Innovation In the Treatment of Pediatric Cardiac Disease



Biltmore Hotel, Coral Gables, FL

LIVE MEETING

December 3-6, 2015

Virtual Meeting using iLearn®
with Global Transmission Featuring
Simultaneous Spanish Translation

Faculty:

Redmond Burke, MD
Robert Campbell, MD
Aldo R. Castañeda, MD, PhD
Paul Checchia, MD
Mitchell Cohen, MD
John Costello, MD, MPH
Martin Elliott, MD
Jon Epstein, MD
J. William Gaynor, MD
Rani Gereige, MD
Jeff Jacobs, MD
Tom Karl, MD
John Kheir, MD
Terry King, MD
Narendra Kini, MD
Lisa Kohr, MSN, CPNP

Carole Lannon, MD, MPH
Stacey Lihn
Leo Lopez, MD
Brad Marino, MD, MSCE
Juan-Carlos Muñoz, MD
Richard Ohye, MD
Sara Pasquali, MD
Dan Penny, MD, PhD
Andrew Redington, MD
John Rhodes, MD
Anthony Rossi, MD
Jack Rychik, MD
Steve Schwartz, MD
Girish Shirali, MD
Edward Walsh, MD
Gil Wernovsky, MD

SPONSORED BY:



Nicklaus
Children's
Hospital

MIAMI CHILDREN'S HEALTH SYSTEM

Heart Program




For more information or to register, please visit: nicklauschildrens.org/heartinnovation

Accreditation Statement: The Miami Children's Health System is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.
Designation Statement: The Miami Children's Health System designates this live activity for a maximum of 17 AMA PRA Category 1 Credit(s)™. Physicians should only claim credit commensurate with the extent of their participation in the activity.
Faculty/Planner Disclosure Policy: In accordance with the Accreditation Council for Continuing Medical Education's Standards for Commercial Support, all planners, teachers and authors involved in the development of CME content are required to disclose to the accredited provider their relevant financial relationships. Relevant financial relationships will be disclosed to the activity audience.

- hemodynamic, and echocardiographic features. *Echocardiography* 2009;26:993-98.
4. T. A. Riemenschneider and A. J. Moss, Left ventricular-Right atrial communication, *The American Journal of Cardiology*, vol. 19, no. 5, pp. 710-718, 1967.
 5. Jacobs JP, Burke RP, Quintessenza JA, Mavroudis C. Congenital Heart Surgery Nomenclature and Database Project: ventricular septal defect. *Ann Thorac Surg* 2000;69(Suppl):S25-35.
 6. O. M. Cheema, A. A. Patel, M. C. Su, and D. J. Shah, Gerbode ventricular septal defect diagnosed at cardiac MR imaging: case report, *Radiology*, vol. 252, no. 1, pp. 50-52, 2009.
 7. Perloff JK. Ventricular Septal Defect. In: *The Clinical Recognition of Congenital Heart Disease*, 4th ed, W.B. Saunders Company, Philadelphia 1994. p.396.
 8. Eltzschig HK, Rosenberger P, Löffler M, et al: Impact of Intraoperative Transesophageal Echocardiography on Surgical Decisions in 12,566 Patients undergoing Cardiac Surgery. *Ann Thorac Surg* 2008 Mar;85(3):845-52.
 9. Gosciniak, et al, The Gerbode defect — a frequent echocardiographic pitfall, *Kardiologia Polska* 2012; 70, 11: 1191-1193.
 10. Acar, P., Séguela, P.-E. and Hascoet, S. (2011), The Gerbode Defect or Left Ventricular to Right Atrial Shunt Assessed by Transthoracic 3D Echocardiography. *Echocardiography*, 28: E140-E142. doi: 10.1111/j.1540-8175.2011.01396.
 11. Kelle et al, The Gerbode defect: the significance of a left ventricular to right atrial shunt, *Cardiol Young*. 2009 Nov;19 Suppl 2:96-9.
 12. Anderson, et al, The Gerbode Defect: A Ventriculo-Atrial Defect (CTSNet) available from <http://www.ctsnet.org/sections/clinicalresources/congenital/article-24>.

CCT

Corresponding Author



*Lucky R. Cuenza, MD
Clinical Research Fellow, Section of
Cardiac Rehabilitation
Philippine Heart Center
East Avenue
Quezon City, Philippines, 0850
Tel: +63 02 9252450;
Fax: +63 02 9220551
lrcuenza@gmail.com*

*Anna Christina Adora, MD
Fellow in Training, Department of Adult
Cardiology
Philippine Heart Center
Quezon City, Philippines, 0850*

*Thessie Minelli O. Valdez, MD
Research Fellow in Peripheral Vascular
Medicine
Philippine Heart Center
Quezon City, Philippines, 0850*

*Cheryl P. Fomaneg, MD
Consultant in Echocardiography, Non
Invasive Section
Philippine Heart Center
Quezon City, Philippines, 0850*

*Efren R. Vicaldo, MD
Congenital Heart Disease Specialist,
Philippine Heart Center
Quezon City, Philippines, 0850*

Letters to the Editor

Congenital Cardiology Today welcomes and encourages Letters to the Editor. If you have comments or topics you would like to address, please send an email to: LTE@CCT.bz, and let us know if you would like your comment published or not.

DIRECTOR ADULT CONGENITAL HEART DISEASE Chicago and Suburbs

Advocate Children's Hospital/Advocate Medical Group seeks a BC/BE **Director of our Adult Congenital Heart Disease** program. Our goal is to develop a top tier, nationally recognized full service ACHD program. ACH's Heart Institute has been a major center for pediatric cardiology and cardiovascular surgery for 25 + years includes 21 + pediatric cardiologists, & 3 congenital CV surgeons. The adult cardiology practice has 100's of Advocate affiliated cardiologists. Both the pediatric and adult cardiology services have several inpatient and many outpatient sites throughout the metro Chicago region. The candidate would be responsible for organizing and directing the Adult Congenital Heart Disease Program throughout the Advocate Health Care system. Our goal is to develop a top tier nationally recognized high quality full service ACHD program.

The focus of this new position includes: Clinical management of complex ACHD population, development and growth of the program, the interdisciplinary team and demonstrated leadership, clinical and teaching experience. Research interest and options for academic affiliation are in place.

We require 5+ yrs. experience, unrestricted MD/DO degree, eligibility for Illinois licensure, BC in either pediatric or adult cardiology, fellowship trained or demonstrated experience in ACHD clinical competency, leadership and organizational skills. BC/BE in ACHD.

AMG is a 1,300+ physician led multi-specialty group affiliated with AHC, the largest Illinois healthcare provider. ACH is one top 10 largest national network providers, recognized by *U.S. News & World Report* as a national leader in Pediatric Cardiology, Cardiovascular Surgery and Neonatology.



Please send detailed cover letter and CV to:
Nancy Mathieu - Nancy.mathieu@advocatehealth.com.

www.advocatehealth.com | www.advocatechildrenshospital.com/ach/ | www.amgdoctors.com/about-us.com

Medical News, Products & Information

Compiled and Reviewed by Tony Carlson, Senior Editor

The William Novick Global Cardiac Alliance

Announcing the William Novick Global Cardiac Alliance - a new non-profit charity launched in November 2014 by 18 professional colleagues formerly of the International Children's Heart Foundation (ICHF).

Professor William Novick is a Pediatric Cardiothoracic Surgeon and Paul Nemir Professor of International Child Health at the University of Tennessee Health Science Center.

Professor Novick and colleagues left the ICHF, and launched the Novick Cardiac Alliance (NCA) as a new, completely separate entity after a 22 year experience operating on over 7500 children, and developing over 25 independent pediatric cardiac surgical programs in 5 continents. The NCA was founded with the continued goals of collaboratively operating on children with heart disease, educating teams in their own environment, innovating new solutions to complex problems and fostering sustainable solutions to pediatric cardiac care.

With this announcement, the NCA is extending an invitation to professionals in pediatric congenital cardiac care to volunteer for our 2 week cardiac teams. We need Cardiac Surgeons, Cardiologists (ECHO and Cath lab), Anesthesiologists, Perfusionists, Intensivists, PICU nurses and Respiratory Therapists. Volunteers are not required to pay for accommodation or travel. The NCA has 30 trips scheduled for 2015.

The NCA is by design non-sectarian and provides comprehensive care to all children with congenital or acquired heart disease regardless of gender, ethnicity, religion, political ideation, genetic factors or economic means.

Founding Board of the William Novick Global Cardiac Alliance are: William Novick, MD, MS; Brian Forsberg, MPH CCP; David Wieduwilt, BME CE; Elizabeth Novick, BSN/RN; Eugene Suslin, MD; Farzana Shah, RN; Frank Molloy, RN MSc; Humberto Rodriguez, MD; Igor Polivenok, MD, PhD; Jean Towne; Karen Bowtell, RN, PGDip; Kathleen Fenton, MD; Marcelo Cardarelli, MD, MPH; Martina Pavanic, BSN/RN; Pavel Shauchenka, MD; Sjarhei Liauchonak, MD; Sri Rao, MD; and Stacey Marr, RN, MSc.

For more information got to: facebook.com/cardiacalliance or www.cardiac-alliance.org

Acute Kidney Injury can be a Serious Complication Following Heart Surgery

Newswise — The anti-inflammatory drug dexamethasone helps prevent serious kidney complications that can arise following

heart surgery, according to the results of a randomized clinical trial. The findings, which appeared in an upcoming issue of the *Journal of the American Society of Nephrology* (JASN), could lead to a change in care for patients during cardiac operations.

Acute Kidney Injury (AKI) is one of the most devastating complications following cardiac surgery. Approximately 1% of patients undergoing cardiac surgery require dialysis to treat severe AKI that arises after surgery, and the incidence is higher among patients with pre-operative chronic kidney disease. These patients experience strikingly high death rates while in the hospital that exceed 40%. "One percent sounds like a small percentage; however, given the fact that each year, over half a million people undergo heart surgery in the United States alone; this means that an estimated 5000 patients develop renal failure, and of those about 2500 die as a result of this complication," said Kirolos Jacob, MD (University Medical Center, Utrecht, The Netherlands). He noted that these figures are rising due to the aging population.

Because heart surgery initiates an inflammatory reaction in the body that can have negative effects on the kidneys, Dr. Jacob and his colleagues wondered whether giving patients dexamethasone, an anti-inflammatory drug, could decrease the risk of severe AKI following cardiac surgery. The team analyzed the results of a large randomized controlled trial called the Dutch Dexamethasone for Cardiac Surgery (DECS) trial, which included 4465 patients undergoing cardiac surgery who were randomized to receive placebo or dexamethasone during surgery. The original trial tested whether dexamethasone could reduce the risk of a variety of major postoperative complications. In this analysis, the investigators specifically examined kidney failure and focused on the most severe form: AKI requiring dialysis.

Dexamethasone appeared to protect against the development of severe AKI. Patients who received the drug had about a 2.5-times lower risk of developing AKI requiring dialysis compared with those receiving a placebo.

"The beneficial effects of dexamethasone were particularly present in those who already had pre-existing kidney disease before heart surgery," said Dr. Jacob. "This reinforces the fact that this drug could be of major importance for the increasing elderly population with pre-existing kidney disease undergoing a heart operation."

The study is the largest randomized, placebo-controlled trial showing a potential benefit of any therapeutic drug for the prevention of severe kidney injury following heart surgery. A single dose of dexamethasone during a heart operation is inexpensive, straightforward, painless, and safe for patients. "These advantages make the intervention very accessible and cost-effective, especially since the costs for dialysis are very high," said Dr. Jacob.

2nd Annual Basic & Advanced Fetal Cardiac Symposium and Workshop

Rush University Medical Center

Sep 10-12, 2015; Chicago, IL USA

www.FetalCardiacSymposium.com



Study co-authors include David Leaf, MD, MMSc, Jan Dieleman, MD, Diederik van Dijk, MD, PhD, Arno Nierich, MD, PhD, Peter Rosseel, MD, PhD, Joost van der Maaten, MD, PhD, Jan Hofland, MD, PhD, Jan Diephuis, MD, Fellery de Lange, MD, PhD, Christine Boer, PhD, Jolanda Kluin, MD, PhD, Sushrut Waikar, MD, MPH, for the Dexamethasone for Cardiac Surgery (DECS) Study Group.

Disclosures: The DECS study was sponsored by grants from the Netherlands Organisation for Health Research and Development (ZonMw) and the Dutch Heart Foundation.

The article, entitled "Intraoperative High-Dose Dexamethasone and Severe Acute Kidney Injury after Cardiac Surgery," appeared online at <http://jasn.asnjournals.org/> on May 7, 2015.

Founded in 1966, and with more than 15,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.

Hospital to Create Emergency Experiences Using Virtual Reality: Walk Around Inside Crisis Situations, Educate Medical Professionals

Next Galaxy Corp (OTC: NXGA) recently announced the signing of an agreement with Miami Children's Hospital. Next Galaxy will develop immersive Virtual Reality medical instructional content for patient and medical professional education using the Company's VR Model. Per the multi-year agreement, Next Galaxy and Miami Children's Hospital are jointly creating VR Instructionals on cardiopulmonary resuscitation (CPR) and other lifesaving procedures, which will be released as an application for smartphones.

Incorporating eye gaze control, gestures, and voice commands while "walking around" inside an emergency medical experience or crisis, Next Galaxy's Virtual Reality Model engages participants far beyond today's methodology of passively watching video and taking written tests.

"Assessments are incorporated directly into the medical VR models. We will design situations where participants are required to make the appropriate decisions about proper techniques. The Virtual CPR instructional will measure metrics and provide real-time feedback ensuring participants accurately perform CPR techniques. Further, the instructional will explain any mistake and prompt users to try again when errors are made. Supportive messages are delivered upon success," states Mary Spio CEO, Next Galaxy Corp.

The medical VR models will be viewable through smartphones and desktops as 3D, and via VR devices such as Google Cardboard, VRONE and Oculus Rift.

For further information, visit www.nextgalaxycorp.com.



PICS-AICS
Pediatric and Adult Interventional Cardiac Symposium

SAVE THE DATE
SEPT. 18-21, 2015

ARIA • LAS VEGAS

WWW.PICSYMPOSIUM.COM



LIVE CASE DEMONSTRATIONS • ABSTRACT SESSIONS • "MY NIGHTMARE CASE IN THE CATH LAB" • HOT DEBATES • WORKSHOPS • SMALLER BREAKOUT SESSIONS •



SPONSORED FOR CME CREDIT BY
RUSH UNIVERSITY MEDICAL CENTER

Master Class in Congenital Cardiac Morphology

With world renowned cardiac pathologist Professor Robert Anderson, MD, FRCPath

Sept. 30 to Oct. 2, 2015 • Children's Hospital of Pittsburgh of UPMC • Pittsburgh, PA

This activity has been approved for AMA PRA Category 1 Credits™ • The University of Pittsburgh is an affirmative action, equal opportunity institution.

Learn more at www.chp.edu/masterclass



Higher Activity Levels Increase Survivability of ICD Patients

Patients who had higher activity levels following ICD implantation had better survival, according to research in the *Journal of the American Heart Association*. The research will be simultaneously presented at the Heart Rhythm Society 2015 Scientific Sessions.

An implantable cardioverter-defibrillator (ICD) is a battery-powered device that typically combines a "generator" placed under the skin near the shoulder with a wire that is inserted into the heart through the vein beneath the collarbone. ICDs are very effective at recognizing fast, potentially life-threatening heart rhythm disorders and providing timely shocks that restore a normal rhythm. However, many patients who receive ICDs may still be at risk of dying from progression of their underlying heart disease or other problems.

In the largest study on the relationship between activity and survival in ICD patients, researchers analyzed how active participants were in the first 30-60 days after implantation and then over time up to four years. ICDs collect patients' daily activity automatically using sensors embedded in the device itself, which determines whether patients are active or not on a minute-by-minute basis. "Active" measured in this way means approximately a walking speed of 2 miles an hour.

Researchers found that patients in the most active group after getting an ICD had a 40% absolute lower risk of death at four years compared to patients who had engaged in the least activity. Regardless of age, gender or device type, lower average activity during the first 30-60 days was independently associated with a 44% higher risk of death over time. A similar analysis looking at activity over several years demonstrated a similar risk of death for each 30 minutes' difference in activity. "We expected to see a difference, but we were struck by the magnitude of these results," said Daniel B. Kramer, MD, MPH, lead author of the study and Assistant Professor of Medicine at Harvard Medical School in Boston, MA.

"Patients' functional status clearly predicts survival. Our hope would be to use activity as a factor in not just predicting outcomes but also to guide strategies that may improve outcomes. But that is much further down the line."

Researchers studied the ALTITUDE registry, a nationwide database that involved 98,437 patients enrolled in a remote monitoring program. About 57% of the patients had received a new or replacement ICD and 43% had received cardiac resynchronization therapy (CRT-D) devices in 2008-12. CRT-D therapy combines an ICD with cardiac resynchronization therapy. Patients were followed for a median 2.2 years.

"What is intriguing about our results is that looking at just one piece of information collected automatically after a device is implanted provides very powerful prognostic information about how they are likely to do over the next several years," Kramer said.

The major limitation of the analysis, researchers noted, is that this study design only supports describing an association between activity and survival. Further studies are needed to test strategies aimed at promoting activity or using this information to change patients' treatment.

Co-authors are: Susan L. Mitchell, MD, MPH; Joao Monteiro, PhD; Paul W. Jones, MS; Sharon-Lise Normand, PhD; David L. Hayes, MD; and Matthew R. Reynolds, MD, M.Sc. Author disclosures and funding information are on the manuscript. The ALTITUDE study is supported by Boston Scientific.

Tiny Heart, Big Promise - Understanding How Cells Become Coronary Vessels May Lead to Advances in Repairing Heart Damage

Newsweek — The heart has its own dedicated blood supply, with coronary arteries that supply oxygen-rich blood to the heart and cardiac veins that remove deoxygenated blood. This system of vessels nourishes the heart, enabling it to pump blood to all the other organs and tissues of the body. Yet despite their critical importance, the process and molecules required for coronary vessel development have not been fully determined.

Studying zebrafish, investigators at The Saban Research Institute and the Heart Institute of Children's Hospital Los Angeles discovered a new source for cells that can develop into coronary vessels and have identified the signaling protein, a chemokine called CXCL12, which guides this process. Results of the study was published online May 26th by the journal *Developmental Cell*.

Zebrafish have emerged as an important vertebrate model for cardiovascular research for a number of reasons, including the ability to regenerate its heart if damaged, and because the transparency of the embryos allows easy observation of internal processes like blood vessel development. Using confocal and time-lapse imaging, the investigators were able to visualize coronary vessels developing from the endocardium, or the inner lining of the heart – specifically from the atrioventricular canal, the structure that divides the heart into compartments.

"This furthers our efforts into heart regeneration to repair human hearts," said Ching-Ling (Ellen) Lien, PhD, principal investigator at The Saban Research Institute of CHLA and senior author on the paper. "We have now found a novel source of cells that can differentiate into coronary vessels and have identified the factors required."

Lien and her team observed that zebrafish with a mutation at the CXCR4 receptor survive, but are not able to form coronary vessels or undergo heart regeneration following injury. Since fish without this mutation are able to do both, the investigators concluded that an interaction between CXCR4 receptors on endothelial cells and the CXCL12b protein expressed by the myocardium regulate the process. In addition to providing basic information about the developing heart, this finding may also have clinical relevance.



Educating, operating, and supporting pediatric cardiac care around the world

www.cardiac-alliance.org/volunteer



An anatomical illustration of the human aorta. A triaxial catheter with a blue handle and a silver wire is shown inserted into the aorta. A mesh stent is mounted on the catheter, positioned to treat a narrowing (coarctation) in the aortic wall. The stent has a 'zig' pattern design. The catheter is covered by a sheath.

NuDEL™

Triaxial BIB Catheter
with Sheath & Covered
CP Stent for the Aorta

The All-In-One Aortic Stent System

The NuDEL™ Stent Delivery System is designed for the efficient and effective treatment of Coarctation of the Aorta.

The NuDEL includes a triaxial balloon in balloon designed catheter with a Covered Mounted CP Stent™, which is then covered by a sheath as an all-in-one system. Combining the proven technologies of our NuMED BIB® balloon catheter and our Covered CP Stent™, the NuDEL System employs both our compact delivery method and the “zig” pattern stent design.

The NuDEL System is available for immediate purchase in the EU. Contact us or your local distributor to place an order.

NuMED

World Leader in Pediatric Cardiology

NuMED, Inc. | 2880 Main Street | Hopkinton, NY 12965 USA
Tel: 315.328.4491 | Fax: 315.328.4941 | www.numedforchildren.com

CE 0120

"Children or young adults may not be aware of having abnormal coronary vessels because their circulation is adequate until the heart is stressed by increased demands, for instance when participating in strenuous sports," explains Lien, who is also an assistant professor at the Keck School of Medicine and an investigator at the Cardiovascular Thoracic Institute, both at the University of Southern California. "Then suddenly, an apparently healthy, young person dies. Alternatively, a person with abnormal coronary vessels might have higher risk of experiencing heart attacks later on in life. Our findings will guide future study toward understanding these devastating conditions in order to be better able to diagnose them and develop interventional strategies."

The first author, Dr. Michael R.M. Harrison is a CIRM scholar and Saban RCDF fellow. Additional contributors include: Ying Huang and Arthela Osorio, The Saban Research Institute of CHLA; Jeroen Bussmann and Arndt F. Siekmann, Max Planck Institute for Molecular Biomedicine, Muenster, Germany; Long Zhao, C. Geoffrey Burns and Caroline E. Burns, Harvard Medical School; and Henry M. Sucov, Broad Center for Regenerative Medicine and Stem Cell Research at USC.

The study was supported in part by the National Heart, Lung and Blood Institute R01HL096121, The Saban Research Institute Career Development Award, and a California Institute for Regenerative Medicine (CIRM) postdoctoral fellowship TG2-01168.

For more information, visit CHLA.org.

Letters to the Editor

Congenital Cardiology Today welcomes and encourages Letters to the Editor. If you have comments or topics you would like to address, please send an email to: LTE@CCT.bz, and let us know if you would like your comment published or not.



CONGENITAL CARDIOLOGY TODAY

Can Help You Recruit:

- Pediatric Cardiologists
- Pediatric Interventional Cardiologist
- Adult Cardiologist focused on CHD
- Congenital/Structural Heart Surgeons
- Echocardiographers, EPs
- Pediatric Transplant Cardiologist

Reach over 6,000 BC/BE Cardiologists focused on CHD worldwide:

- Recruitment ads include color!
- Issue email blast will include your recruitment ad!
- We can create the advertisement for you at no extra charge!

Contact:

Tony Carlson

+1.301.279.2005 or

tcarlsonmd@gmail.com

CONGENITAL CARDIOLOGY TODAY

© 2015 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499-online). Published monthly. All rights reserved.

www.CongenitalCardiologyToday.com

8100 Leaward Way, PO Box 444,
Manzanita, OR 97130 USA
Tel: +1.301.279.2005; Fax: +1.240.465.0692

Publishing Management:

- Tony Carlson, Founder, President & Sr. Editor - TCarlsonmd@gmail.com
- Richard Koulbanis, Group Publisher & Editor-in-Chief - RichardK@CCT.bz
- John W. Moore, MD, MPH, Group Medical Editor - JMoore@RCHSD.org
- Virginia Dematatis, Assistant Editor
- Caryl Cornell, Assistant Editor
- Loraine Watts, Assistant Editor
- Chris Carlson, Web Manager
- Rob Hudgins, Designer/Special Projects

Editorial Board: Teiji Akagi, MD; Zohair Al Halees, MD; Mazeni Alwi, MD; Felix Berger, MD; Fadi Bitar, MD; Jacek Bialkowski, MD; Mario Carminati, MD; Anthony C. Chang, MD, MBA; John P. Cheatham, MD; Bharat Dalvi, MD, MBBS, DM; Horacio Faella, MD; Yun-Ching Fu, MD; Felipe Heusser, MD; Ziyad M. Hijazi, MD, MPH; Ralf Holzer, MD; Marshall Jacobs, MD; R. Krishna Kumar, MD, DM, MBBS; John Lamberti, MD; Gerald Ross Marx, MD; Tarek S. Momenah, MBBS, DCH; Toshio Nakanishi, MD, PhD; Carlos A. C. Pedra, MD; Daniel Penny, MD, PhD; James C. Perry, MD; P. Syamasundar Rao, MD; Shakeel A. Qureshi, MD; Andrew Redington, MD; Carlos E. Ruiz, MD, PhD; Girish S. Shirali, MD; Horst Sievert, MD; Hideshi Tomita, MD; Gil Wernovsky, MD; Zhuoming Xu, MD, PhD; William C. L. Yip, MD; Carlos Zabal, MD

Free Subscription to Qualified Professionals:

Send your name, title(s), hospital or practice name, work address and url, phone, fax and email to: sub@cct.bz.

Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors, and are not necessarily the views of Congenital Cardiology Today.



Barth Syndrome Foundation

Symptoms:

Cardiomyopathy, Neutropenia, Muscle Weakness,
Exercise Intolerance, Growth Delay, Cardiolipin Abnormalities

www.barthsyndrome.org



Fetal Cardiology Opportunity

Health care is transforming. We're transforming health care.
Join Us

If you are an experienced fetal cardiologist interested in joining a progressive practice, consider Arizona Pediatric Cardiology (APC). APC, in affiliation with Phoenix Children's Hospital (PCH), provides state-of-the-art, comprehensive inpatient and outpatient pediatric cardiology services throughout the state of Arizona. Phoenix Children's Hospital, ranked 16th by the U.S. News and World Report, has 24 CVICU beds and two cardiothoracic surgeons who perform over 500 surgeries annually.

Help grow our already large fetal heart program and collaborate with maternal-fetal medicine and neonatology practices in the Phoenix metro area. As a leader in fetal cardiology, our practice sponsors the Annual Phoenix Fetal Cardiology Symposium and is a participating site in multiple fetal cardiology multi-center trials.

This is an excellent opportunity to be a part of a team of echocardiographers who perform and interpret transthoracic and transesophageal echocardiograms in an IAC-accredited lab. Clinical responsibilities also include inpatient and outpatient general pediatric cardiology care.

In addition to fetal cardiology, APC provides a wide range of services and programs including interventional and diagnostic catheterization, electrophysiology/ablations, cardiac imaging (MRI), adult congenital heart disease program, anti-coagulation program, preventive cardiology, heart transplant/heart failure program and a metabolic stress lab.

About Phoenix

- ★ Wide variety of cultural and sporting events
- ★ California coast within a day's drive
- ★ Diverse geography offers desert, lake and mountain recreational activities
- ★ 310 days of sunshine per year
- ★ Family-oriented community
- ★ Excellent school districts
- ★ Unlimited activities for children

Candidate requirements:

- ✓ A minimum of three years' experience
- ✓ A fourth year of fellowship in advanced imaging training, including fetal echocardiography, preferred

Strength in numbers

25 pediatric cardiologists, including
5 fetal cardiologists

5 fetal sonographers

2 advanced practice fetal
cardiac nurse coordinators

5 office locations

More than **1,500** fetal echocardiograms
performed last year

LEARN MORE

www.pediatrix.com/clinicalcareers
800.243.3839, ext. 5589



Toshiba **gives you a voice.**



At Toshiba, we go beyond listening and give you a voice. Because understanding your business is our business, your challenges become our challenges, and together, we find creative ways to solve them. With decades of experience and groundbreaking technology built into all of our products, you can see you're actually being heard. It's not just lip service.

COMPUTED TOMOGRAPHY

MAGNETIC RESONANCE

PET/CT

ULTRASOUND

VASCULAR X-RAY

X-RAY