

CONGENITAL CARDIOLOGY TODAY

News and Information for Pediatric and Congenital Cardiovascular Physicians and Surgeons

Vol. 5 / Issue 6
June 2007
North American Edition

WWW.CONGENITALCARDIOLOGYTODAY.COM

INSIDE THIS ISSUE

Noninvasive Monitoring of Cardiac Output: Benefits of NIRS Technology

by George M. Hoffman, MD and Nancy S. Ghanayem, MD
-Page 1

SPECIAL COMMENTARY

Allen Tower: 'The Case is Altered'

by Michael J. Tynan and Shakeel A. Qureshi, MD
-Page 2

Highlights from Cardiology 2007 – 10th Annual Update on Pediatric Cardiovascular Disease

by Gil Wernovsky
-Page 8

DEPARTMENTS

Medical News, Products and Information

-Page 18

CONGENITAL CARDIOLOGY TODAY

9008 Copenhaver Drive, Ste. M
Potomac, MD 20854 USA
Tel: +1.301.279.2005
Fax: +1.240.465.0692

Editorial and Subscription Offices:

16 Cove Road, Ste. 200
Westerly, RI 02891 USA
www.CongenitalCardiologyToday.com
www.CHDvideo.com

© 2007 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499-online). Published monthly. All rights reserved. Congenital Cardiology provides timely news and information for pediatric and congenital cardiologists. Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors, and are not necessarily the views of Congenital Cardiology Today.

See Recruitment Ads on pages: 4, 5, 7, 9, 10, 11, 12, 16, 19 and 20

NONINVASIVE MONITORING OF CARDIAC OUTPUT: BENEFITS OF NIRS TECHNOLOGY

By George M. Hoffman, MD and Nancy S. Ghanayem, MD

While adequate organ perfusion is a primary goal of supportive and therapeutic intensive care, numerous studies show that cellular oxygen deficit (dysoxia or shock) is under-recognized, under-diagnosed and under-treated. Significantly impaired organ blood flow frequently occurs in patients who are normotensive, but who have reduced or maldistributed cardiac output. Indicators of oxygen supply-demand relationships such as venous oximetry (SvO₂) can significantly improve the diagnosis of shock states, and can reduce the incidence and severity of multi-organ system dysfunction and mortality when utilized to guide resuscitation and support.

There is a narrow temporal window for diagnosis and intervention in shock states before organ injury occurs. Multiple studies suggest early detection of inadequate oxygen delivery and appropriate goal-directed interventions can reduce complications of shock, while delayed diagnosis and treatment is ineffective. According to research presented in Pediatric Cardiac Surgery Annals,[1] authors indicate that current clinical descriptors of cardiac output and circulatory function that lead to shock are inadequate, and provide incomplete and sometimes


misleading information, especially in children. Without a clear picture of the cardiovascular status, the physician's ability to assess systemic and regional tissue oxygenation, to intervene early in shock, and to limit resulting complications is impaired.

Noninvasive monitors are becoming increasingly useful in children for the assessment of cerebral and somatic (i.e., organ bed, skeletal muscle tissue) tissue oxygenation, providing a more comprehensive clinical picture that allows early recognition and treatment of circulatory abnormalities that lead to multiorgan dysfunction. One of the most exciting developments is a novel multi-site, non-invasive monitor for the brain and body using near-infrared spectroscopy, or NIRS.

Limitations of Conventional Monitoring Techniques

A closer look at the mechanism of shock sheds some light on the limitations of conventional monitoring techniques. The body reflexively responds to shock by redistributing blood flow to preserve the brain and heart, leaving other tissue beds – typically kidneys, liver, and intestines - at increased risk for silent ischemia.[2-4] In pediatric patients, shock is typically characterized by low cardiac output and very high systemic vascular resistance.[5] If regional

Continued on Page 3



SPECIAL COMMENTARY

ALLEN TOWER: 'THE CASE IS ALTERED'

By Michael J. Tynan and Shakeel A. Qureshi, MD

All interventionists involved with children know the NuMED name and the company's products. Most of us will know the name of the company President, Allen J. Tower, and many will have met him. However, those involved in other fields of the speciality may know little about NuMED and Allen Tower and know less about how important the products are to paediatric cardiology.

At present Allen, and thus the NuMED company, is in dispute with the United States Government and it appears that there is the possibility that Allen could be sentenced to a term in prison. This would be a disaster for children with congenital heart disease. NuMED is the only manufacturer specialising in equipment for catheter interventions for babies and children.

Allen Tower has always been an innovator in our field. He was the manufacturer who collaborated with Dr. Berman in the development of the balloon angiographic catheter. This catheter, now manufactured by another company, is for many, the mainstay of diagnostic angiocardiology.

When, in the mid 1980s he started NuMED Inc, in his home territory of upstate New York, he turned to paediatric cardiologists and asked what it was that we needed. That has been his philosophy throughout. Anyone with a need for new or modified catheters has been listened to and where ever possible he has tried to satisfy that need.

Usually his interpretation of our ideas has resulted in a superior product than we had



Allen J. Tower

"The esteem in which Allen is held by we interventionists was shown when, in 2003, he was the recipient of the 'Lifetime Achievement Award' at the 7th Pediatric Interventional Cardiac Symposium."

envisaged. First of the NuMED 'inventions' was the co-axial angioplasty and valvuloplasty balloon catheter, the TyShak. This allowed the production of very low profile balloons with very rapid inflation and deflation times. These are beneficial for all children but are ideal when it comes to interventions in newborn babies and small infants. The ease of introduction of relatively large balloon catheters in small arteries and veins and the short time that the balloon was obstructing blood flow means shorter procedure times, less circulatory disturbance and less vascular damage. These catheters are used by most paediatric interventionists worldwide, so any interruption in their supply would have a negative impact on the outcome for babies and children in the USA but its effect would be worldwide.

His other innovations are too numerous to detail, but some should be mentioned. Together with Philipp Bonhoeffer he developed both the 'Multitrack' catheter system used by many interventionists. They have also produced a catheter implantable pulmonary valve which is under study in several countries. An integral component of this system is another NuMED innovation, a stent for children developed with John Cheatham, the CP stent with a special balloon for its stable implantation, the BiB balloon. A development of the stent is the Covered CP stent which has been life saving for a number of patients. All this equipment has approval in Europe but not in the USA. And it is the supply of the BiB balloon, the CP and covered CP stents, on a compassionate basis, which lit the fuse for the initial FDA investigation.

The esteem in which Allen is held by we interventionists was shown when, in 2003,

he was the recipient of the 'Lifetime Achievement Award' at the 7th Pediatric Interventional Cardiac Symposium. To date, he is the only recipient of this award who does not have a medical qualification.

Allen has been unstinting with his time and resources for the improvement of our equipment and thus our ability to do a better job for our patients. He has also been very generous in his support of paediatric cardiology meetings and has made many donations to his local community.

Allen has built up a team of highly skilled and very dedicated workers in Hopkinton, but there is no doubt that he is the driving force behind NuMED and the whole paediatric community, workers, patients and parents will suffer if he is incarcerated. In England during the reign of Queen Elizabeth I, a gentleman was accused of holding a Catholic Mass, which was illegal. His lawyer Edmund Plowden (1518-1585) proved that it had been the work of a provocateur, not a priest. Plowden said 'The case is altered, no priest, no Mass.' Now, for Allen 'The case is altered,' but with no easy way out.

We do not have detailed knowledge of the full legal case but it appears that his problems may be, to a great extent, due to his willingness to help physicians and their patients. Perhaps the bureaucratic errors, for that is what they appear to be from this side of the Atlantic, were as much to do with healthcare professionals such as us, as with Allen. In which case we have a duty to support him. We urge you to go to <http://www.thepetitionsite.com/takeaction/250006502> and sign the petition in his support.

~CCT~

Michael J. Tynan, MD, FRCP
Emeritus Professor of Pediatric
Cardiology at King's College,
London, UK

Shakeel A. Qureshi, MB, ChB, FRCP
Head of Service for Paediatric
Cardiology
Consultant Paediatric Cardiologist /
Senior Lecturer
Evelina Children's Hospital, Guy's &
St. Thomas' Hospital Foundation
Trust, London, UK

Continued from Page 1

vascular resistance is sufficiently high, ischemic damage may occur to specific organ beds, even in the presence of normal global oxygen economy and blood pressure.

Conventional noninvasive assessment tools reliably measure heart rate, blood pressure, arterial oxygen saturation, and that data can be used to approximate some of the determinants of cardiac output and oxygen delivery. But because these are systemic metrics, they may be late indicators of shock. Other techniques such as indirect calorimetry, alveolar gas analysis, bioimpedance, and pulse contour and Doppler studies add more information about cardiac output and systemic oxygenation, but can still leave clinicians uncertain about the adequacy of regional perfusion. They are also more prone to influencing factors (e.g. physiology, environment, staff skills), making the data less reliable. The result may be an inaccurate picture of the clinical status of critically ill infants and children, since even experienced physicians may not recognize patterns of clinical parameters indicative of the early stages of shock.

Venous oximetry monitoring is a more accurate reflection of oxygen metabolism[6] in neonates than conventional monitoring tools, but poses a risk of injury and/or infection during placement and removal of fiberoptic indwelling catheters. Further, since SvO₂ is a global measure, it does not reliably detect maldistribution of blood flow or regional tissue dysoxia.

The Mechanics and Use of NIRS Technology

NIRS technology eliminates the technical challenges and risks associated with invasive SvO₂ monitoring, and identifies regional perfusion abnormalities. Noninvasive light-emitting diodes (LEDs), that transmit near-infrared light to the tissue bed at a depth of about 2.5-3cm, and corresponding photodetectors are placed on the area of the brain or body to be assessed.

Like pulse oximeters, NIRS monitors utilizes the differential absorption of two or more wavelengths of light by hemoglobin as it associates and disassociates with oxygen. Unlike pulse oximetry, NIRS techniques measure the average hemoglobin saturation in a field of tissue, with arterial, capillary, and venous components. Over a wide range of conditions, the amount of red and infrared absorption by hemoglobin in a field of tissue is most closely related to the regional venous blood saturation. NIRS devices which provide spatial resolution and algorithms calibrated from a 25%;75% arterial:venous blood ratio thus can provide a continuous, non-invasive approximation of tissue oxyhemoglobin saturation in a field of tissue 2-3cm beneath the skin. These measures allow early and rapid identification of inadequate tissue oxygenation, and allow guided interventions before

hemodynamic changes further destabilize critically ill pediatric patients.

When used on two or more sites such as over the brain and a peripheral organ bed, NIRS (through cerebral/somatic oximetry) provides a real-time window on oxygen economy in two distinct tissue beds and a comparative model of regional tissue oxyhemoglobin saturation.[7-8] An accurate assessment of cerebral and somatic tissue oxygenation in critically ill patients enables timely intervention to prevent or minimize the often irreversible damage caused by dysoxia. Studies which capture continuous, real-time monitoring of cerebral and regional tissue oxyhemoglobin saturation support the advantages of NIRS in reducing post-operative morbidities.

Multi-Site rSO₂ Monitoring

Our institution has evaluated a novel monitoring strategy that employs NIRS to measure regional oxygen saturation (rSO₂). Initially designed as a cerebral monitoring tool—and used successfully for that purpose in published studies of adults undergoing cardiac surgery for early detection of cerebral blood flow derangements--the INVOS System (Somanetics Corporation, Troy, MI) comprises a multi-detector system (up to four probes) that enables concurrent and continuous cerebral and regional oxygen saturation monitoring.

Somanetics' NIRS device received U.S. FDA approval for trends monitoring of rSO₂ in the brain and skeletal muscle tissue. Clinical data in both children and adults support a link between cerebral rSO₂ values less than 40% to 50%, or a deviation from baseline of more than 20%, and hypoxic-ischemic neural injury. [9-12]

Our hospital initially used the Somanetics device, which originally came with two probes, to monitor pediatric patients for cerebral oxygen derangements. Since infants and neonates have limited geography on their foreheads, we chose to place the second probe over the flank. Our intent was to assess blood flow and regional blood oxygenation in a tissue bed with circulatory controls very different from the brain, such as the kidney. We found this multi-site approach provided us with a unique window on blood flow throughout the body, and we've integrated it into our standard care protocol.

Autoregulatory mechanisms help maintain cerebral blood flow and brain oxygenation at relatively constant levels over a wide range of physiologic conditions. The kidney, on the other hand, is a high-flow, low-extraction organ in which renovascular resistance is under strong sympathetic control. The sympathetic stress response to shock triggers a dramatic increase in vascular



For information, please call 1-800-BRAUN2 (227-2862)

www.bbraunusa.com



Working Together to Develop a Better Tomorrow

Pediatric Cardiology

We are looking for a non-invasive Pediatric Cardiologist for the Orlando Florida area. Established in 1994, we serve the Central Florida area with outpatient clinics, inpatient consults, fetal echo's and cardiac catheterizations.

Competitive salary and benefits, including paid health, disability, malpractice insurance and CME allowances.

Come enjoy Florida with its beautiful beaches, lakes, great outdoor activities, and many cultural and sporting events. The Orlando area boasts 'A' schools and a prominent local University.

For a glimpse into our practice visit our website at:
www.carson-appleton.com

Please contact: Thomas Carson, M.D. for more details on this great opportunity via email at: redvet@aol.com or our office at 407-902-2866.

Submit your CV via fax to 407-902-2585.




Figure 1. SvO₂ parallels changes in cerebral and somatic rSO₂ in postoperative neonates.




Figure 2. A linear combination of cerebral and somatic rSO₂ provided a better approximation of SvO₂ than either alone.



PICS 2007 JULY 22-25, 2007
 Bellagio, Las Vegas

www.picsymposium.com

resistance within the kidney, with accompanying circulatory impairment and heightened risk of ischemic organ damage. Direct, real-time, dual-site monitoring of cerebral and peri-renal perfusion clearly demonstrates the profound differences in the oxygenation of these organ beds as the patient's circulatory status changes.

Based on our experiences with the cerebral-somatic oximeter at Children's Hospital of Wisconsin, we find it reliable in showing the adequacy of site-specific perfusion and the maldistribution of blood flow. As such, we believe it's today's best non-invasive method for detection of shock; providing unique information for a more complete patient assessment. It is appropriate for use on any patient at risk for shock, and we presently use it on patients undergoing operative procedures or cardiac catheterization, as well as patients admitted to the neonatal and pediatric intensive care units. Because it is entirely non-invasive, NIRS has also been useful during triage assessment within the emergency room and for assessment of inpatients not admitted to critical care units.

Important Findings

The ability to monitor regional oxygen saturation at multiple sites is an important strength of the cerebral/somatic oximeter using NIRS. Trending correlation studies of cerebral and somatic oximetry performed in neonates after the Norwood procedure—the first stage surgical palliation of Hypoplastic Left Heart Syndrome—have demonstrated parallel changes in tissue oxyhemoglobin saturation (rSO_2) and SvO_2 (Figure 1).[13] Prolonged low rSO_2 is more predictive of ischemic neurologic damage that traditional assessments used currently during surgery and in the ICU.[14]

Dual-site monitoring of rSO_2 enables an accurate prediction of the mixed SvO_2 . In

fact, calculation of the SvO_2 based on the linear combination of cerebral and somatic NIRS monitoring yields a closer approximation of the SvO_2 than does either regional measure alone (Figure 2).[15] Furthermore, although inter-patient variability is high, within-patient trends of two-site cerebral/somatic oximeter correlate closely with SvO_2 values (Figure 3).[15]

Figure 4 depicts the output of two-site cerebral/somatic oximeter using NIRS in a patient undergoing resuscitation from hypovolemic/septic shock.[1] Data trends underscore the correlation between somatic and cerebral rSO_2 and SvO_2 modeling. Early detection of shock allows for aggressive resuscitative measures with fluid and epinephrine, restoring rSO_2 values to normal levels.

Conclusion

Multi-site NIRS is feasible technology that plays an important role in monitoring critically ill patients at risk for circulatory derangements. Cerebral-somatic oximetry with NIRS should not replace conventional monitoring, but rather serve as an important adjunct that allows for continuous, earlier assessment of perfusion abnormalities. This permits therapeutic interventions to maintain adequate systemic perfusion and oxygen economy, and thereby reduce ischemia-related complications with associated organ failure and mortality.

References

1. Hoffman GM, Ghanayem NS, Tweddell JS. Noninvasive assessment of cardiac output. *Ped Cardiac Surg Ann.* 2005;8:12-21.
2. Haljamae H. Pathophysiology of shock. *Acta Anesthesiol Scand.* 1993;98:3-6.
3. Dyess DL, Powell RW, Swafford AN. Redistribution of organ blood flow after hemorrhage and resuscitation in full-term piglets. *J Pediatr Surg.* 1994;29:1097-1102.



Asheville Cardiology Associates Recruiting BC/BE Pediatric Non-Invasive Cardiologist

Known for clinical excellence, Asheville Cardiology Associates is seeking a third pediatric cardiologist with advanced skills in both inpatient and outpatient arenas. Those with expertise in new modality imaging, adult-congenital cardiology, or interest in exercise physiology or electrophysiology, are encouraged to contact us. Development of new programs within the practice is encouraged.

The Peds Division has a complex patient base, with active fetal and adult-congenital programs. Outpatient practice encounters ~3,000 clinic visits, ~2,000 echoes per year, with a growth rate of 5-7% annually. Inpatient is performed at Mission Hospitals, consistently rated in Top 100 Heart Hospitals in last six years. Its new children's outpatient facility is well represented by subspecialists and ancillary services.

Asheville is a beautiful city of 70,000 (county has 200,000 plus MSA of 391,000). Located in the Blue Ridge Mts. of North Carolina, it offers excellent public and private schools, universities and colleges. Amenities such as great restaurants, arts, music, & theater put it on the list of best places to live in many publications. A mild four season climate gives opportunities for mountain biking, hiking, camping, fishing, golf, whitewater activities and skiing.

Visit our website, www.avlcard.com
for more information.

All inquiries remain confidential.
Please send CV to:

James J. McGovern, MD, FACC
Fax: (828) 277-6350
Email: jimm@avlcard.com



CONTEMPORARY QUESTIONS
IN CONGENITAL HEART DISEASE
THE LEFT HEART

Four Seasons Hotel

June 10 -12, 2007; Toronto, Canada

www.sickkids.ca/cardiacsymposium




Figure 3. The two-site NIRS model shows very good prediction of SvO₂ for within-patient trends.




Figure 4. Two-site NIRS trends from a patient undergoing resuscitation from hypovolemic/septic shock. Early aggressive resuscitation with fluid and epinephrine to normal regional rSO₂ values restored urine output. The effect of changes in pCO₂ on cerebral blood flow are evident at 0700. The mirror changes in cerebral and somatic rSO₂ suggest that total cardiac output was relatively limited but that the distribution changed.

4. Dantzker DR. Monitoring tissue oxygenation: the search for the graal. *Chest*. 1997;111:12-13.
5. Parker MM, Hazelzet JA, Carcillo JA. Pediatric considerations. *Crit Care med*. 2004;32 (suppl):S591-S594.
6. Kremzar B, Spec-Marn A, Kompan L, et al. Normal values of SvO₂ as therapeutic goal in patients with multiple injuries. *Intensive Care Med*. 1997;23:65-70.
7. Boushel R, Langberg H, Olesen J, et al. Monitoring tissue oxygen availability with near infrared spectroscopy (NIRS) in health and disease. *Scand J Med Sci Sports*. 2001;11:213-222.
8. Fadel PJ, Keller DM, Watanabe H, et al. Noninvasive assessment of sympathetic vasoconstriction in human and rodent skeletal muscle using near-infrared spectroscopy and Doppler ultrasound. *J Appl Physiol*. 2004;96:1323-1330.
9. Levy WJ, Levin S, Chance B: Near-infrared measurement of cerebral oxygenation: Correlation with electroencephalographic ischemia during ventricular fibrillation. *Anesthesiology* 83:738-746, 1995.
10. Austin EH III, Edmonds HL Jr, Auden SM, et al: Benefit of neurophysiologic monitoring for pediatric cardiac surgery. *J Thorac Cardiovasc Surg* 114:707-715, 717, 1997.
11. Nollert G, Jonas RA, Reichart B: Optimizing cerebral oxygenation during cardiac surgery: A review of experimental and clinical investigations with near infrared spectrophotometry. *Thorac Cardiovasc Surg* 48:247-253, 2000.
12. Hoffman GM, Stuth EA, Jaquiss RD, et al: Changes in cerebral and somatic oxygenation during stage 1 palliation of hypoplastic left heart syndrome using continuous regional cerebral perfusion. *J Thorac Cardiovasc Surg* 127:223-233, 2004.
13. Hoffman G, Robertson FA, Berens R, et al. Relationship of cerebral and somatic oxygenation by two-site near infrared spectroscopy and venous saturation in neonates following the Norwood procedure. Presented at the World Congress of Anesthesiology.



Healing hearts. Training minds. Bringing hope.

5075 Arcadia Avenue Minneapolis, MN 55436 U.S.A.
Toll Free: 888.928.6678; Phone: 952.928.4860; Fax: 952.928.4859

www.childrensheartlink.org

14. Dent CL. Cerebral oximetry—preventing brain injury after the Norwood procedure. *Neonatal Intensive Care*. 2006;19(6):45-47.

15. Hoffman GM, Stuth EA, Berens RJ, et al. Two-site near-infrared transcutaneous oximetry as a non-invasive indicator of mixed venous oxygen saturation in cardiac neonates. *Anesthesiology*. 2003;97:A-139.

Special Note: Images used in this article are reprinted from Seminars in Thoracic & Cardiovascular Surgery: Pediatric Cardiac Surgery Annual, V8, George M. Hoffman, Nancy S. Ghanayem, and James S. Tweddell, Non-invasive Assessment of Cardiac Output, 12-21, 2005, with permission from Elsevier.

~CCT~






Corresponding Author:



George M. Hoffman, MD
 Medical Director & Chief of Pediatric Anesthesiology; Associate Director, Pediatric Intensive Care Unit; Professor, Anesthesiology & Critical Care
 Children's Hospital of Wisconsin
 Medical College of Wisconsin
 9000 W Wisconsin Ave.
 Milwaukee, WI 53226 USA
 Office Phone: 414-266-3560

Nancy S. Ghanayem, MD
 Associate Professor, Pediatric Critical Care, Medical College of Wisconsin, and Medical Director, Cardiac Critical Care, Children's Hospital of Wisconsin, Medical College of Wisconsin

PEDIATRIC CARDIOLOGIST

To enhance an expanding program, the Division of Pediatric Cardiology at the Medical College of Wisconsin and Children's Hospital of Wisconsin is recruiting board-certified/board eligible pediatric cardiologists to join a dynamic practice of **12 pediatric cardiologists and 3 cardiovascular surgeons**. Children's Hospital of Wisconsin is in the process of a major facility expansion, and pediatric cardiology and critical care fellowship training programs are currently in place. Academic cardiologists are being recruited at the assistant or associate level in the following roles:



The **Interventional candidate** will serve in the largest congenital program in Wisconsin performing more than 450 catheterizations per year with two additional catheterization labs planned. Expertise in hemodynamics, angiography, and skills and experience in interventional techniques required.

The **intensive care candidate** will assist with management of more than 650 operative procedures annually. A new 24-bed pediatric cardiac ICU is under construction. Expertise in preoperative and postoperative hemodynamics, pharmacology with skills in intensive care technical procedures and assist device management required.

The **candidate for electrophysiology** should have expertise in invasive and non-invasive electrophysiology and skills and expertise in diagnosis and management of complex arrhythmias including catheter-based intervention/ablation.

The **candidate for non-invasive imaging** will join a busy clinical practice with opportunities for research and career development; the department has developed a NIH-funded pediatric echo research lab with 2 research staff and 2 dedicated myocardial deformation/2-D and 3-D imaging review stations. Expertise in transthoracic and transesophageal echocardiography; interest in fetal echocardiography and cardiac MRI and CT desirable.

The Medical College of Wisconsin offers a competitive compensation/benefits package. Interested applicants should fax or E-mail CVs to:
Stuart Berger, M.D.
Director, Division of Pediatric Cardiology
Medical College of Wisconsin
sberger@chw.org
Fax# 414-266-3261

Children's Specialty Group*



Evolving Concepts in Management of Complex Congenital Heart Disease

October 5-6, 2007 San Diego, CA USA

(858) 966-8587 FAX; (888) 829-9249 TEL

www.rchd.org

HIGHLIGHTS FROM CARDIOLOGY 2007 – 10TH ANNUAL UPDATE ON PEDIATRIC CARDIOVASCULAR DISEASE

By Gil Wernovsky, MD



The Cardiac Center at The Children's Hospital of Philadelphia hosted our 10th Annual Postgraduate Course at Disney's Yacht and Beach Club Resorts in sunny Orlando, Florida from February 21st-25th, 2007. Over 750 professionals and exhibitors gathered from around the globe to hear late breaking research, discuss controversial topics, review current practices and enjoy each other's company and insight. I was privileged to learn from such a distinguished faculty and insightful group of course attendees, and we look forward to planning our 11th meeting next year in Scottsdale, Arizona.

What distinguishes this postgraduate from other excellent subspecialty meetings is the multidisciplinary group of course attendees. Physicians made up approximately half of the attendees, and included representation from all disciplines involved in the care of children



with heart disease including: cardiologists, intensivists, surgeons, anesthesiologists, neonatologists and maternal fetal specialists. Approximately 50 physicians were fellows, residents or medical students. The remaining attendees included: advanced practice, operating room, catheterization lab and bedside nurses; sonographers; physicians assistants; respiratory therapists; and the largest group of perfusionists ever present at our meeting (48). Finally, a growing number of administrators continue to partner with front-line care givers in sessions devoted to delivery of care, fiscal responsibility and patient safety.

Featured Research, Posters and Abstracts

From over 100 initially submitted, 65 abstracts were presented in two separate poster sessions from 40 institutions and 16 countries. Those that have not been previously published as abstracts are included later in this electronic supplement. The top three abstracts were presented as oral presentations by Kirsten Odegard of Children's Hospital, Boston ("Prospective longitudinal study of coagulation profiles in children with hypoplastic left heart syndrome from stage I through Fontan completion"), John Costello, also of Children's Hospital, Boston ("A systematic initiative to reduce blood stream infections in a pediatric cardiac intensive care unit") and Jon Kaufman from Children's Hospital of Denver/University of Colorado Health Science Center ("Correlation of abdominal oximetry with gastric tonometry: measurement of splanchnic oxygenation in neonates and infants with congenital heart disease"). In a very close vote by the entire faculty, the 4th Annual Outstanding Investigator Award was given to Dr. Kaufman and his colleagues.

Surgical innovations and practices in complex congenital heart disease were the focus of many of the posters. William Douglas (Lexington) presented information on an adjustable flow device for

systemic artery to pulmonary artery shunts; Daniel Nento (Cleveland) presented improved outcomes in side-to-side (as opposed to end-to-side) technique in augmenting the ascending aorta in patients with hypoplastic left heart syndrome and Martin Zahorec (Bratislava) discussed the role for arterial duct closure in patients undergoing modified Blalock-Taussig shunt placement. The most challenging aspects of hypoplastic left heart syndrome and other univentricular heart disease were tackled with researchers showing several "outcome challenges." Areas covered included results of the Fontan palliation in children with trisomy 21 (Francis Moga, Minneapolis); catheter intervention for hypoplastic left heart syndrome with intact atrial septum (Chris Petit – Philadelphia), neurodevelopmental abnormalities overall (Anke Furck, Kiel), and the role that inherent coagulation abnormalities may play – even before any surgery – in patients with a univentricular heart (Nina Hakáčová, Bratislava). Lindsay Ryerson from the University of Michigan discussed the role of heparin therapy in protein losing enteropathy. Wasim Khan from Rainbow Babies described the physiological responses to exercise in children after the Fontan Operation, while the group from The Children's Hospital of Philadelphia utilized magnetic resonance imaging to study the effects of exercise on power loss after Fontan palliation. Collaboration between institutions was evident in Duke's efforts (presented by Maura Catherine Baldwin) to validate the follow-up pro-




gram reported by The Children's Hospital of Wisconsin.

High fidelity simulators were shown to significantly improve cardiac center nurses' confidence in critical scenarios – an important support in times when rapid education and skills development is an issue for many centers (Roberta Hales, Philadelphia). Simulators were also shown to improve time for cardiac care teams to initiate extracorporeal membrane oxygenation (Cecilia St. George-Hyslop, Toronto). Resuscitation scientists also showed improvements in cardiopulmonary resuscitation techniques with use of a Voice Advisory Manikin system (Robert Sutton, Philadelphia).

Administrators and others taking a "50,000 foot view" of our care systems were benefited by a cost analysis of various mechanical support techniques as a bridge to heart transplantation (Andrei Morgan, Newcastle-Upon-Tyne), and the role that virtual regionalization models (Shabib Alhadheri, Syracuse) can play in improving outcomes when providers can be mobile but care can't be given under one roof. Sonographers and their colleagues had their interests focused on the roles for the discharge echocardiogram in improving outcomes (Christa Barlow, Cincinnati), and on the correlation between pulmonary vein Doppler signals and left atrial pressures (Sarah Gelehrter, Ann Arbor).

The field of electrophysiology was investigated in five abstracts. Aimee Malpass and colleagues from the University of Mississippi showed that amiodarone is both safe and effective in the acute management of supraventricular tachycardia in neonates. The clinical and electrophysiologic characteristics of both left and right ventricular tachycardia in children with structurally normal hearts were presented by Nelangi Pinto from The Children's Hospital of Philadelphia. Armando Alfaro and colleagues from Costa Rica showed that ventricular single lead pacing is an effective and safe method to maintain atrioventricular synchrony in third degree atrioventricular block.

Fetal cardiology was also strongly represented. Paul Brooks studied the effects of antenatal diagnosis of pulmonary atresia with an intact ventricular septum and found that while there was an improvement in systemic oxygenation prior to the initiation of prostaglandin E1, there was no overall improvement in mortality. Kirsten Dummer reported from the Boston experience that prenatal diagnosis of tetralogy of Fallot is highly accurate and that it is unusual to see progression to pulmonary atresia. Anita Szwast from The Children's Hospital of Philadelphia demonstrated that maternal hyperoxygenation in the fetus with hypoplastic left heart syndrome is a safe and useful



PEDIATRIC CARDIOLOGIST CORPUS CHRISTI, TX


Driscoll Children's Hospital is actively recruiting for Board Certified/Eligible Pediatric Cardiologists with expertise in Echocardiography, Perinatal Echocardiography, cardiac MRI/CT and Electrophysiology. The center has a new catheterization laboratory, ten satellite clinics, two cardiothoracic surgeons, and six pediatric cardiologists.

Driscoll Children's Hospital is a not-for-profit two hundred bed tertiary care hospital serving over 33 counties of South Texas. The hospital shares an affiliation with the Texas A&M University Health Science Center and maintains a large Pediatric residency. The Cardiology service actively participates in a variety of teaching activities within the training Program.

Driscoll Children's Hospital is located in Corpus Christi, which is located on the Texas Gulf Coast. Corpus Christi enjoys some of the finest weather and outdoor activities in the United States and is the 2nd most popular vacation destination in the state of Texas. The city is nestled behind the barrier islands of Mustang and Padre and is the gateway to Padre Island National Seashore, a short 15-minute drive from downtown.

Please contact:

Tim Hanners
 Tel: (361) 694-4759
 email: tim.hanners@dchstx.org
 Driscoll Children's Hospital
 3533 S Alameda Street
 Corpus Christi, TX 78411
www.driscollchildrens.org




Watch LIVE CASES from Congenital Cardiology Today
www.CHDVideo.com
 Presented courtesy of:
 The Pediatric and Adult Interventional Therapies for Congenital and Valvular Disease (PICS)
 and
 The 6th International Workshop on Interventional Pediatric Cardiology (Workshop IPC)
 Watch a Percutaneous Valve Implantation, a Perimembranous VSD Closure or both



Pediatric Cardiologist

FLORIDA - The Department of Pediatrics at the University of Florida, College of Medicine-Jacksonville AND Wolfson Children's Hospital in Jacksonville are recruiting a full-time faculty member to the Division of Pediatric Cardiology. Candidates require a MD degree and Florida medical license eligibility. Applicants must be BE/BC in pediatric cardiology and have a primary focus on non-invasive imaging. This individual should have expertise in all aspects of echocardiography, including TEE, fetal, three-dimensional and is expected to direct the non-invasive imaging program of the University of Florida, College of Medicine-Jacksonville. The appointment will be at the non-tenure accruing level of Assistant Professor/Associate Professor/Professor depending upon experience.

Deadline for accepting applications will be August 1, 2007 with an anticipated negotiable start date of September 1, 2007. Academic rank will be commensurate with qualifications and salary is negotiable.

Forward letter of intent, curriculum vitae, and three letters of recommendation to:

Mobeen H. Rathore, MD
 Professor and Assistant Chairman
 Department of Pediatrics
 653-1 West Eighth Street
 Jacksonville, FL 32209
 Tel: 904-244-3050, and/or fax 904-244-3028
 e-mail: mobeen.rathore@jax.ufl.edu

Visit our website at www.hscj.ufl.edu/peds


The University of Florida is an Equal Opportunity Institution.



procedure to assess the pulmonary vasculature. Lisa Wise-Faberowski from the University of Colorado presented two abstracts utilizing a chronic hypoxia model in the developing brain to describe the use of magnetic resonance spectroscopy, as well as the effects of isoflurane.

Several abstracts focused on patient care from the nursing perspective, and included a report from Stanford that emphasized the importance of continued nursing educational sessions as a mechanism to improve professional autonomy and enhances patient safety, a similar point made by the nursing staff at The Children's Hospital of Philadelphia who routinely utilize the benefits of an educational resource nurse in the training of new cardiac intensive care nurses. Nursing and safety interventions were the highlight of many posters. Pharmacist-led pediatric medication safety teams (Jennifer Costello, Newark – winner of the outstanding nursing poster award), patient safety checklists at handovers (Deanna Edwards, Little Rock), and targeted mining of administrative databases for cardiovascular medication errors (Diana Alexander, Baltimore) were all shown to play roles in improving outcomes and safety for our most vulnerable patients. Sternotomy pain control augmentation with continuous local anesthetic infusions (Marilyn Torres, Miami), and nurse coordinated programs to improve enteral feeding success (Britt Elin Fredriksen, Oslo) drew comments and crowds in the busy session.

In the featured research session, Andrew Redington (Toronto) 'teased' the audience with the study design of a randomized trial of aspirin and warfarin for patients with a Fontan circulation, and then told us that the results were still being analyzed....undoubtedly to garner an invitation to speak next year!



kids play.

Camp Odayin provides residential, day and family camp experiences for children with heart disease from all around the world. The camp offers the opportunity to strengthen self-confidence, gain independence, develop life skills, and meet other young people with similar health, emotional and social concerns.

worries rest.

fun happens!

www.campodayin.org

Scott Bradley and colleagues from the Medical University of South Carolina performed a prospective cohort study on neonates with midline arch reconstruction (both for Hypoplastic Left Heart Syndrome and biventricular repairs), and reported a surprisingly high incidence of swallowing dysfunction, aspiration, and vocal cord injury. Barry Byrne from the University of Florida in Gainesville reported on exciting work into both the mechanism and treatment for inherited cardiomyopathies. Finally, Lynn Mahoney updated the group on the exciting work being conducted by the Pediatric Heart Network, sponsored by the National Heart, Lung and Blood Institute. See www.pediatricheartnetwork.com for more details.

Plenary Sessions

The opening session on cardiopulmonary resuscitation featured experts in the fields of resuscitation science, nursing, cardiac intensive care and pediatric cardiac anesthesia. Vinay Nadkarni (Philadelphia) and Mary Fran Hazinski (Nashville) reviewed the current state of the art and the science behind the American Heart Association guidelines for pediatric resuscitation. Sarah Tabbutt (Philadelphia) gave excellent examples of why these guidelines are not always relevant to postoperative patients with congenital cardiac disease, and Ravi Thiagarajan (Boston) showed us some practical and convincing work on the improvement gained with a simulation program for staff in an intensive



care unit. George Hoffman (Milwaukee) gave a comprehensive review of the literature on post-resuscitation brain protection, and speculated on the relevance to children following cardiac surgery. Finally, the day ended with a spirited debate between Tom Spray (Philadelphia) and Jim Tweddell (Milwaukee) on the risks



**PEDIATRIC
CARDIAC INTENSIVIST
PEDIATRIC CARDIOLOGISTS**

HCA is the largest healthcare company in the US, with over 170 hospitals in 21 states. Whether you are looking for a place to start, or somewhere to complete a successful career in pediatric cardiac care, chances are we can help you find it.

We have opportunities in big cities and rural areas, near sandy beaches and skyward reaching mountains. Wherever your internal compass points, it's likely that we can help guide you to a destination that will match your personal preferences while fulfilling your career dreams.

As our co-founder, Dr. Thomas Frist, Sr. said, "Bettering the human condition is the greatest good any individual can achieve." You are doing your part by taking care of our most precious commodity, the children who will be our future. Now let us do our part by helping you find your dream job.

*Call or
e-mail
me today
for more
information
on opportunities
in your
sub-specialty.*



If you are a fellow in training, please be sure to ask about our excellent stipend program!

For more information on available positions contact Kathleen L. Kyer, Manager, National Pediatric Subspecialty Recruitment at:
888-933-1433 or
kathleen.kyer@hcahealthcare.com



For information, please call 1-800-BRAUN2 (227-2862)

www.bbraunusa.com



Working Together to Develop a Better Tomorrow



Pediatric Cardiologist

The Division of Pediatric Cardiology of Sanger Clinic and Carolinas Physicians Network in Charlotte, NC is recruiting a board certified/eligible pediatric cardiologist to join a rapidly growing practice of six pediatric cardiologists and two cardiothoracic surgeons. We are searching for a cardiologist with excellent interpersonal skills to build relationships with patients, families and primary care physicians in a predominantly out-patient setting.

The candidate should have expertise in echocardiography and ideally an interest in adult congenital heart disease. Skills in interventional catheter procedures for back-up purposes would also be attractive but certainly not mandatory. The Sanger Clinic is the largest congenital heart center in North Carolina performing three hundred surgeries and three hundred catheterizations per year.

The practice is affiliated with Carolinas Medical Center but operates much like a private practice with very competitive compensation and benefits. The hospital authority is one year away from opening Levine Children's Hospital, an eighty five million dollar state-of-the-art facility. Interested applicants should fax or e-mail their CV's to me.

Herbert J. Stern, MD, FACC
Director, Division of Pediatric Cardiology
Sanger Clinic and Carolinas Heart Institute

hsstern@sanger-clinic.com

Fax# 704-543-0018

and benefits of open cardiac massage during resuscitation. Dr. Spray's video comparing the size of his hand to that of a typical neonatal heart was quite visually "revealing" of the difficulties of direct massage in small neonates.

Two plenary sessions were given back to back on the second day of the conference, covering coronary artery disease in children and double outlet right ventricle. Jack Rome (Philadelphia) moderated the session on coronary artery disease in children, which began with Alan Friedman (New Haven) reviewing normal coronary anatomy followed by a discussion on the anatomy, hemodynamics, and clinical characteristics of abnormalities of coronary origin (including anomalous coronary artery from the pulmonary trunk and intramural coronary arteries), as well coronary fistulae, and Kawasaki Disease. Michele Frommelt (Milwaukee) discussed how to best image the coronary arteries by echocardiography which was followed by Paul Weinberg (Philadelphia) showing superb anatomic specimens of the previously discussed anomalies. Mark Fogel and Jeff Hellinger (Philadelphia) subsequently advocated use of cardiac magnetic imaging and computerized tomographic imaging to delineate these anomalies showing beautiful images, including the "fly-thru" technique of three dimensional imaging. David Nykanen (Orlando) illustrated the continued necessity for coronary angiography as an additional imaging modality, especially in evaluation and potential intervention of several coronary anomalies. Paul Stephens (Philadelphia) discussed the determinants of coronary flow, including how best to assess myocardial perfusion abnormalities. The session concluded with the debate on whether children with an anomalous origin of a coronary artery from the 'wrong' sinus of Valsalva should have surgery. Tim Feltes (Columbus) argued that only those who have evidence of ischemia should have surgery, as the true risk of a sudden catastrophic event is unknown; Jim Tweddell countered that surgery should occur irrespective of ischemia, since there is an increased risk of ischemia and sudden death. By a show of hands, the winner by a small margin was Tim Feltes.

The plenary session discussing double outlet right ventricle was moderated by Jack Rychik (Philadelphia). Professor Robert Anderson (London) convinced the audience that double outlet right ventricle is not a single isolated defect, such as tetralogy of Fallot, but rather is a series of different relationships between the great arteries, where both are supported by the anatomic right ventricle, the ventricular septal defect being the most important feature. Paul Weinberg followed showing beautiful specimens, including those with mitral atresia and straddling of the mitral valve, as well as various relationships of the ventricular septal defect to the great vessels. Meryl Cohen (Philadelphia) reviewed

Do You Want to Recruit a Pediatric Cardiologist?

Advertise in the only monthly publication totally dedicated to pediatric and congenital cardiology. For more information: call +1.301.279.2005, or send an email to: TCarlsonmd@gmail.com

the key goals of the echocardiographic evaluation: the relationship of the ventricular septum to the great vessels, the number of ventricular septal defects, the relationship of the great arteries to each other; the relative size of the ventricles, and the potential pathway from the left ventricle to great vessels through the ventricular septal defect. This point was re-emphasized by Pedro del Nido (Boston) showing that the creation of a three dimensional baffle from the left ventricle to the aorta or the pulmonary artery is the greatest surgical challenge in these patients, and advocated for three dimensional imaging as a routine part of the preoperative assessment.

For a change of pace, a 4th plenary session - "Improving Safety for Cardiovascular Patients" - proved to be an exciting testament to the work being done by so many colleagues in our field to improve the safety of our care delivery systems. Peter Laussen (Boston) led off the session with a wonderful overview entitled "Changing culture for continuous quality improvement." Defining moments were described that required novel efforts to improve governance, management, and effect culture change. Dr. Laussen showed the central role of nursing leadership in promoting culture change and in leading to improved consistency in patient care. Adequate support for culture change can involve significant financial resources and attending physician accountability was emphasized. We heard about Children's Hospital in Boston's Program For Patient Safety And Quality, the importance of audits to measuring and improving performance, and the imperative for leadership to promote a balance of work and personal goals.

Karen Harrington (Montreal) gave an absolutely outstanding presentation "How may we improve communication during patient handover?" The audience was treated to a broad review with ideas of safety as a "dynamic non-event", communication breakdowns as a major cause of sentinel events, and that a knock-on effect of working hour restrictions is an increased need for handovers and an imperative to do them well. Dr. Harrington implored our field to recognize handovers as both a transfer of responsi-

bility and accountability as well as information, and that improvements should be driven by local gradual change.

Troy Dominguez (Philadelphia) reviewed the risk factors for nosocomial infections, with an emphasis on central line associated blood stream infection, ventilator associated pneumonia, surgical site infections, the particular vulnerability of patients with congenital cardiac disease given their high number of risk factors, and the crucial role of adopting a team approach for better control and prevention.

Richard Ohye (Ann Arbor), in an effort towards "Ensuring patient safety during clinical trials, the ethics of human subjects' research," gave surely the most humorous presentation of the session. The information given, though, was absolutely critical and gave the audience plenty to sink its teeth into. Dr. Ohye dared us to ask "Do trials help patients and are they relevant?" He reviewed definition of various clinical trials, problems with historically controlled trials, the benefits of randomization, and the clever way in which, given the inexorable progress in medicine, trials using historical controls always tend to favor the current intervention at question. Dr. Ohye implored us to base our practice of medicine in research and to promote, as well with its attendant requirements for respect for persons, beneficence, justice, and equipoise. John Charpie (Ann Arbor) followed with a presentation on "Developing and maintaining an effective quality assurance and improvement program." We learned the multiple aspects of quality (safe, effective, patient-centered, timely, etc.) and the details and rationale behind the "Plan-Do-Check-Act" cycle for improvement.

Patricia Hickey (Boston) riveted us with her bold opening that mandated nurse staffing ratios are simply not the answer as they can exacerbate staffing- and work-related stress for staff nurses. Her presentation, "Rational work-hour assignments for bedside nurses: what's the rationale?" described that best practice in addressing nurse staffing depends on a multifactorial approach. The best models allow staffing to patient demand without having a negative impact

on staff; nurses should maintain control and choice over their own schedules. Introduced to Florence Nightingale's edict to "put patient in best condition for nature to act," the audience was then led through Ms. Hickey's team's "Nightingale metrics" and her novel "continuity of care index."

The session closed with a terrific debate between Martin Elliott (London) and Alan Friedman on the pros and cons of restricted work hours for physicians. To most accounts, despite excellent arguments by both combatants, the contest was a draw (perhaps because all of the junior doctors had reached their work quota and were at the pool!). Nonetheless, patient safety remains a crucial element in improving outcomes for our patients. Look forward to upcoming meetings from Children's Hospital, Boston (www.childrenshospital.org/clinicalservices/Site457/mainpageS457P15sub-level119Flevel120.html) the Hospital for Sick Children at Great Ormond Street in London, and The Children's Hospital of Philadelphia (www.chop.edu/cardiology2008) to promote cardiovascular patient safety.

The closing plenary session on Heterotaxy Syndrome/Isomerism was perhaps the most well attended Sunday morning session of the past 10 years. The session was dedicated to the work and in honor of Stella Van Praagh who was lauded by all of the speakers for her insight into complex congenital cardiac disease, her compassion, her love of teaching her patients and her family. Professors Anderson and Weinberg concisely reviewed a very complex topic where nomenclature is perhaps the most controversial. For those of us (like myself) who were taught and 'grew up' thinking of these diseases as "polysplenia" and "asplenia," Professor Anderson made a very convincing argument to stick to nomenclature based upon the anatomy of the atrial appendages. Paul Weinberg compared Heterotaxy syndrome to "Murphy's Law:" if something can go wrong, it will! Peter Gruber (Philadelphia) showed some new and exciting information regarding the genetic basis of abnormalities of sidedness being a 'proximal' event in cardiogenesis, with

absolutely fascinating microscopic images of cilia directing cardiac cells to one side or another. Leo Lopez (Miami) reviewed the anomalies as shown by echo, Jack Rome (Philadelphia) did the same with angiography, and Mitchell Cohen (Phoenix) showed some very elegant examples of the abnormalities of the conduction system. The session closed with newly released data from the Pediatric Heart Network presented by Andy Atz (Charleston) from the recently completed Fontan cross-sectional study. Interestingly, despite having a higher complexity of disease (and most likely a higher initial mortality risk), the current survivors appear to be no different than other patients with a Fontan circulation in terms of exercise performance, laboratory assessments of ventricular function, and importantly, patient and parent reported quality of life.

The featured lectures this year were given by Martha Curley (Philadelphia), Philipp Bonhoeffer (London), Pedro del Nido (Boston) and Gavin Kerr (Philadelphia). Dr. Curley gave an inspiring overview of what it takes to be a mentor, what (and how) we learn from them, and how they start a generational process of learning, teaching and compassion. Dr. Bonhoeffer reviewed percutaneous pulmonary valve therapy from its humble beginning in the cow fields of France (collaborating with butchers), through stent design (collaborating with jewelers) through implementation in patients (collaborating with industry). His internationally known work was interspersed with funny anecdotes, and wonderful photos from his work in Africa. Dr. del Nido reviewed the pioneering work being done in Boston to 'recruit' two ventricles in the subgroup of pa-

tients with Hypoplastic Left Heart Syndrome and aortic and mitral stenosis. His presentation covered surgical technique, fetal intervention and the underpinnings of basic science suggesting abnormalities of cell migration being the common denominator of such diverse diseases as pulmonary vein stenosis, mitral valve pathology and endocardial fibroelastosis. Finally, Gavin Kerr gave the first Keynote Lecture in Cardiovascular Administration, highlighting the partnership necessary between front-line care givers and hospital administration.

Breakout Sessions

Echocardiography breakout sessions were well attended throughout the meeting, including overviews of three dimensional imaging (Girish Shirali, Charleston), fetal imaging (Jack Rychik, Philadelphia) and intracardiac imaging (Mike Brook, San Francisco). Difficult cases were reviewed by the faculty in an informal session, and Leo Lopez, Michele Frommelt and Norman Silverman (San Francisco) gave wonderful overviews of specific cardiac lesions.

Similarly, the inpatient breakout sessions for intraoperative care, surgery, perfusion and intensive care were comprehensive and insightful. New strategies in monitoring using near infrared spectroscopy are becoming more ubiquitous in our field, and front line experts from Houston, Milwaukee and Charleston updated the group on current strategies and results. A particular focus of all of the breakout sessions was on reducing morbidity, particularly to the central nervous system. Dean Andropoulos and Chuck Frasier (Houston) presented their current "best-practice" model to intraoperative perfusion and monitoring, followed by informal comments from practitioners in Philadelphia, Boston, Ann Arbor and Milwaukee. Lara Shekerdemian (Melbourne) then updated the audience on their prospective studies following neonatal surgery in Australia. Andrew Redington once again showed some seminal work being done on remote ischemic preconditioning, perhaps one of the best examples of translational research we have seen in our field.

The outpatient cardiology breakout sessions were also comprehensive and well attended, including topics such as pulmonary hypertension, routine screening for cardiovascular disease, heart transplantation, Barth's Syndrome, Marfan Syndrome, exercise testing and outpatient decision analysis, to name just a few. Smaller subspecialty breakouts were also held in developmental cardiology, electrophysiology, interventional catheterization and cardiovascular administration.

Nursing has traditionally been a particularly strong component of this meeting, especially in terms of the science pre-



sented by the faculty and the number of attendees from all disciplines in nursing. This year was no exception with a featured nursing plenary session, a hands-on research planning session and nearly 20 abstracts submitted by nurses from around the globe. Barbara Medoff-Cooper (Philadelphia) and Mary Fran Hazinski moderated the first nursing plenary session which focused on the care of neonates with hypoplastic left heart syndrome Eliot May (Milwaukee) led a thorough discussion of low cardiac output in the early postoperative period and the usefulness of continuous monitoring of mixed venous oxygen saturation and near infrared spectroscopy in early detection and early intervention. Erika Wintering (Phoenix) followed with





an excellent review of preoperative and postoperative ventilation strategies, using examples and research to highlight key points. The remainder of the session addressed long term morbidities. Barbara Medoff-Cooper presented late breaking results from her National Institutes of Health funded study of feeding patterns in newborns with congenital cardiac disease, and focused on some of the concerning results in this population - reduced length, weight, and head circumference at 3 and 6 months of age. Jo Ann Nieves (Miami) reviewed her international survey results on practice variability, and Nancy Rudd (Milwaukee) reviewed the results of their formalized home-monitoring program, which has resulted in a dramatic reduction of inter-stage mortality at that center. Kathy Mussatto (Milwaukee) closed the session with an insightful look at factors affecting quality of life for our patients and their families.

The second nursing plenary session focused on low cardiac output syndrome. Jo Ann Nieves and Erika Wintering reviewed several strategies for recognition and management of low cardiac output, which was followed by a comprehensive review of mechanical support of the failing circulation by Lisa Moore (St. Petersburg). A special sunrise session was held with Barbara Medoff-Cooper and Kathy Mussatto on how to get nursing research off the ground, funded and published, followed by presentation of the top three nursing abstracts.



For the first time at this meeting, a full half-day session was devoted to Career Planning for Trainees and Junior Faculty, including rarely-discussed topics at scientific meetings such as: job interview skills, negotiating salary and benefits, comparisons of academic and private practice, work-life balance, managing funding and choosing an academic career and mentor. They graded very favorably by the trainees and junior faculty alike, and will most likely become a routine part of future meetings. An opening plenary session featured a wonderful talk on why clinical pediatric cardiology practitioners need to know about basic science from Jon Epstein (Philadelphia), a self-flagellating talk by Martin Elliott on what he endured during his surgical training, and commentary by yours truly on how the theory of cognitive dissonance in social psychology pertains to the way we read (and sometimes dismiss) the literature and respond to new and/or conflicting data.

Two pre-conferences were also held one on cardiac anatomy and imaging and one

on resuscitation. The anatomy/imaging pre-conference was attended by nearly 150 physicians, nurses and sonographers, and featured anatomic specimens from the Cardiac Registry in Philadelphia and the Van Mierop collection in Gainesville. Wonderful correlations between the specimens and imaging examples brought it all together for the attendees.

The simultaneous session on 'mock codes' was coordinated by Stacie Peddy (Philadelphia) the team from the Center for Simulation, Advanced Education and Innovation at The Children's Hospital of Philadelphia. Small group didactics were held, but most of the experience was 'hands-on' with real life cardiac intensive care scenarios and the use of the SimBaby from Laerdal. Working in small groups, the participants were taken through both predictable (low cardiac output, tachyarrhythmias) and sudden, unanticipated events (respiratory failure, cardiac tamponade and shunt thrombosis) that characterize the 24-48 hours after cardiopulmonary bypass. In addition, pre-operative scenarios





mous audience responses to various clinical scenarios. In the physician session, a number of cases were shown similar to most center's 'surgical conference', once again highlighting areas that require further investigation, such as optimal management strategies for common disease (for example, small ventricular septal defect with aortic regurgitation, management of atrial septal defect) and consensus guidelines for outpatient follow-up. In the afternoon session, nearly 500 respondents were led by Geoff Bird (Philadelphia) in a series of clinical scenarios which highlighted the incredible diversity of management styles around the globe, including management of hyperglycemia (as many respondents used insulin protocols as those who offered no treatment), types of inotropic support and the routine use of delayed sternal closure.

Mike Brook starred as Alex Trebek in Cardiology Jeopardy, simulating the popular game show. Three teams competed on topics from geography, radiographs, boils and sores and historical facts. The 'home' team from The Children's Hospital of Philadelphia (Tom Spray, Jack Rychik and Gil Wernovsky) was pitted against a United States team (Norm Silverman, Tim Feltes and Jim Tweddell), and an All-World team (Andrew Redington, Martin Elliott and Dan Penny). Despite cries of 'foul' and controversial answers, the United States team won in a close battle. Jeopardy is sure to become a standard offering in future meetings.

Finally, there were a number of networking receptions for nurses, trainees, staff physicians and perfusionists, and the Gala Reception was followed by the 1st annual Cardiology Ensemble, featuring musical numbers from the faculty and attendees.

Future Directions

Cardiology 2008 is planned for February 6-10, 2008 in Scottsdale, Arizona. High-

lights from this meeting, as well as those from the 7th Annual International Symposium on Congenital Heart Disease (St. Petersburg, Florida) will be published as a supplement to Cardiology in the Young in the Fall of 2007.



Special thanks to the Staff of the Cardiac Center for their contributions to this report.

~CCT~



*Gil Wernovsky, MD, FACC, FAAP
Staff Cardiologist, Cardiac Intensive Care Unit
Director, Program Development
The Cardiac Center
Professor of Pediatrics
Divisions of Cardiology, Anesthesia and Critical Care Medicine
The Children's Hospital of Philadelphia
University of Pennsylvania
School of Medicine*

*Mailing Address:
Pediatric Cardiology, 34th St and Civic Center Blvd
Philadelphia PA, 19104
215-590-6067 (office);
215-590-5825 (fax)*

wernovsky@email.chop.edu



6th International Pediatric Cardiovascular Symposium

Novel Medical Interventions and Their Role in Children with Heart Disease

June 28-July 1, 2007 ♦ The Ritz-Carlton, Amelia Island, FL

www.choa.org/cme

or call Kathy Murphy 404-785-6480

MEDICAL NEWS, NEW PRODUCTS AND INFORMATION

SickKids Opens The Mitchell Goldhar Cardiac Diagnostic & Interventional Unit at The Hospital for Sick Children, a State-of-the-Art Cardiac Diagnostic and Interventional Unit Specially Designed for Children

The newly integrated unit combines the latest technology to diagnose and treat children with congenital heart disease using minimally invasive procedures that can reduce or eliminate the need for open-heart surgery. The 18,000 sq. ft. unit includes two new catheterization labs with state-of-the-art equipment, an integrated magnetic resonance imaging (MRI) machine and a post-anaesthetic recovery area for patients. Designed with children in mind, the unit's outer space theme provides a fun and stress-free environment in waiting areas, corridors and treatment rooms. Built-in lava lamps, an interactive "asteroid," as well as colorful lights and wall decorations are designed to distract patients from upcoming procedures.

"The new unit allows us to explore new minimally invasive treatments that put less strain on children with cardiac conditions," says Dr. Lee Benson, director of The Mitchell Goldhar Cardiac Diagnostic & Interventional Unit. "Ultimately, we want to improve the quality of care and life for children with heart disease and The Mitchell Goldhar Cardiac Diagnostic & Interventional Unit will help us achieve our goal."

The establishment of The Mitchell Goldhar Cardiac Diagnostic & Interventional Unit was made possible with \$23 million in combined funding from the Ministry of Health and Long-Term Care and from a committed group of donors to SickKids Foundation. The Ministry of Health and Long-Term Care provided \$7.2 million to fund the unit.

Every detail in the unit has been designed to provide a distracting and entertaining environment for children. The entrance to the unit sets the stage for a child's journey through outer space. Patients will first encounter a large "asteroid" that symbolizes a real meteorite that landed on earth over 5,800 years ago. Funded by the Women's Auxiliary of The Hospital for Sick Children, the asteroid has many hidden features and stories that children will want to explore on each visit to the unit. The patient reception area also features a virtual interactive game, computer kiosks and a wall of lava lamps that will entertain both children and their families.

The Labatt Family Heart Centre at SickKids is already home to the largest paediatric interventional catheterization program in Canada and continues to be a leader in Ontario for innovative procedures to treat complex cardiac conditions in children. It conducts approximately 90% of cardiac catheterization procedures for Ontario children, with 1,000 catheterizations performed last year. Approximately half of these procedures are interventional, while the remaining half include diagnostic, electrophysiological, ablation and pacemaker-related procedures.

The Hospital for Sick Children (SickKids), affiliated with the University of Toronto, is Canada's most research-intensive hospital and the largest centre dedicated to improving children's health in the country. For more information, please visit www.sickkids.ca.

Philips' Save an Athlete Campaign Provides Simple Tests to Find Cardiac Conditions in Student Athletes

Royal Philips Electronics (NYSE: PHG, AEX: PHI) announced its continued support of Team of Physicians for Students (TOPS) annual sports physicals event in Phoenix, AZ to help uncover heart conditions that may lead to sudden cardiac death (SCD). Philips' *Save an Athlete* program aims to improve health and wellness by educating student athletes, their families, doctors, athletic directors and coaches about preventing sudden death through early cardiac testing.

Each year, SCD affects approximately 5,000-7,000 young people in the United States. (School Health Corporation. www.schoolhealth.com/shop/mc_archive_berger.asp. Accessed April 16, 2007).

SCD in student athletes is sometimes caused by congenital cardiovascular conditions, such as Long QT Syndrome or Hypertrophic Cardiomyopathy (HCM), which can be identified using easy but sophisticated diagnostic tests, like an electrocardiogram (ECG) or echocardiogram (echo or cardiac ultrasound). Preventative cardiac testing helps detect these types of conditions. Most athletes, their friends and family are unaware of these conditions unless SCD strikes. According to the Cleveland Clinic, HCM is a condition that causes excessive thickening of the heart muscle and affects as many as 1.5 million Americans, making it the most common cause of SCD



In support of infants, children and teens with pediatric cardiomyopathy

CHILDREN'S CARDIOMYOPATHY FOUNDATION

P.O. Box 547, Tenafly NJ 07670

Tel: 201-227-8852 info@childrenscardiomyopathy.org www.childrenscardiomyopathy.org

"A Cause For Today.... A Cure For Tomorrow"

in people under age 30. 9The Cleveland Clinic: Hypertrophic Cardiomyopathy. Accessed on April 16, 2007. www.clevelandclinic.org/heartcenter/pub/guide/disease/hcm/default.htm)

In fact, an ECG exam can help detect heart conditions that account for up to 60% of SCDs in young athletes. (Dominico Corrado et. al., "Cardiovascular Pre-Participation Screening of Young Competitive Athletes for Prevention of Sudden Death: Proposal for a Common European Protocol," European Heart Journal (2005) 26, 516-524, February 2, 2005 - notebook 1).

"Too many student athletes are dying from cardiac conditions that are detectable and treatable," said Paul Steingard, DO and founder of TOPS. "TOPS is thrilled to join forces once again with Philips to help reduce the risk of sudden death in young athletes and to raise awareness around this significant and sometimes overlooked issue."

A leader in the diagnostic health market, Philips started the *Save an Athlete* campaign in 2006 to help support testing programs across the country, such as the TOPS event. Unlike most sports physicals, TOPS events are free, include a family health history review and non-invasive cardiac testing which utilizes ECGs and an echo exam, an ultrasound of the heart.

Last year's TOPS event conducted cardiac testing on over 2,500 student athletes, identifying 385 students that required follow-up examinations. From those, 114 students were not cleared to participate in sports due to potentially fatal heart conditions first detected by abnormal ECGs and echo exams.

"The use of cardiac testing with ECG and echo exams in sports physicals can help identify heart conditions that could trigger sudden death that would not be otherwise identified through an ordinary

Pediatric Cardiologist

BC/BE Pediatric Cardiologist needed to join established Pediatric Cardiology Private Practice Group with 5 offices located in the suburbs of a major metropolitan area in the Midwest with access and interaction with two large university based Pediatric Cardiology Centers.

This position offers the opportunity to achieve financial compensation above the national average.

If you are interested, please email your CV to:

PC-1 @ CCT.bz

© 2007 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499-online) Published monthly. All rights reserved

Publishing Management

Tony Carlson, Founder & Editor
TCarlsonmd@gmail.com
 Richard Koulbanis, Publisher & Editor-in-Chief
RichardK@CCT.bz
 John W. Moore, MD, MPH, Medical Editor/
 Editorial Board
JMoore@CHSD.org
 Jeffrey Green, Contributing Editor

Editorial Board

Teiji Akagi, MD
 Zohair Al Halees, MD
 Mazeni Alwi, MD
 Felix Berger, MD
 Fadi Bitar, MD
 Jacek Bialkowski, MD
 Philipp Bonhoeffer, MD
 Mario Carminati, MD
 Anthony C. Chang, MD, MBA
 John P. Cheatham, MD
 Bharat Dalvi, MD, MBBS, DM
 Horacio Faella, MD
 Yun-Ching Fu, MD
 Felipe Heusser, MD
 Ziyad M. Hijazi, MD, MPH
 Ralf Holzer, MD
 Marshall Jacobs, MD
 R. Krishna Kumar, MD, DM, MBBS
 Gerald Ross Marx, MD
 Tarek S. Momenah, MBBS, DCH
 Toshio Nakanishi, MD, PhD
 Carlos A. C. Pedra, MD
 Daniel Penny, MD
 James C. Perry, MD
 P. Syamasundar Rao, MD
 Shakeel A. Qureshi, MD
 Andrew Redington, MD
 Carlos E. Ruiz, MD, PhD
 Girish S. Shirali, MD
 Horst Sievert, MD
 Hideshi Tomita, MD
 Gil Wernovsky, MD
 Zhuoming Xu, MD, PhD
 William C. L. Yip, MD
 Carlos Zabal, MD

FREE Subscription

Congenital Cardiology Today is available free to qualified professionals worldwide in pediatric and congenital cardiology. International editions available in electronic PDF file only; North American edition available in print. Send an email to Subs@CCT.bz. Be sure to include your name, title, organization, address, phone, fax and email.

Contacts and Other Information

For detailed information on author submission, sponsorships, editorial, production and sales contact, current and back issues, see website: www.CongenitalCardiologyToday.com



Two Perspectives. One Focus

International Symposium on the Hybrid Approach to Congenital Heart Disease
 with live presentations
 Columbus, OH



www.hybridsymposium.com
June 27-29, 2007

