

CONGENITAL CARDIOLOGY TODAY

Timely News and Information for BC/BE Congenital/Structural Cardiologists and Surgeons

May 2013; Volume 11; Issue 5
International Edition

IN THIS ISSUE

Intrapericardial Teratoma in Newborn Infant Revealed by Cardiac Tamponade

by Amadou Gabriel Ciss, MD; Momar Sokhna Diop, MD; Pap Adama Dieng, MD; Pap Salmane Ba, MD; Assane N'diaye, MD, PHD; Mouhamadou N'diaye, MD, PHD; Magaye Gaye, MD; Oumar Diarra, MD, PHD; Mohamed Leye, MD; Souleymane Diatta, MD; Arame Diallo, MD; Fall Lamine, MD; Etienne Birame Sène, MD; Oumar Kane, MD, PHD; Abdou Magib Gaye, MD
~Page 1

Image of the Month #5 – May 2013 Presented by The Archiving Working Group

Contributors: Jeffrey P. Jacobs, MD; Jorge M. Giroud, MD; Robert Anderson, MD; Vera D. Aiello, MD; Diane E. Spicer, BS
~Page 8

An Unusual Case of Prenatal Diagnosis of a Large Left Ventricular Diverticulum and Its Intermediate Term Outcome

by Resham Kaur, MD; Javier Paiz, RDCS; John Brownlee, MD; Umang Gupta, MBBS, DCH
~Page 10

Do Not Close the Door on Closing the Door!

by Stefan C. Bertog, MD; Laura Vaskelyte, MD; Ilona Hofmann, MD; Marius Hornung, MD; Simon Lam, MD; Sameer Gafoor, MD; Horst Sievert, MD
~Page 12

DEPARTMENTS

June Medical Meeting Focus: CSI 2013 and iCI 2013

~Page 13

Medical News, Products and Information

~Page 15

CONGENITAL CARDIOLOGY TODAY

Editorial and Subscription Offices

16 Cove Rd, Ste. 200

Westerly, RI 02891 USA

www.CongenitalCardiologyToday.com

© 2013 by Congenital Cardiology Today
ISSN: 1544-7787 (print); 1544-0499 (online). Published monthly. All rights reserved.

Intrapericardial Teratoma in Newborn Infant Revealed by Cardiac Tamponade

By Amadou Gabriel Ciss, MD; Momar Sokhna Diop, MD; Pap Adama Dieng, MD; Pap Salmane Ba, MD; Assane N'diaye, MD, PHD; Mouhamadou N'diaye, MD, PHD; Magaye Gaye, MD; Oumar Diarra, MD, PHD; Mohamed Leye, MD; Souleymane Diatta, MD; Arame Diallo, MD; Fall Lamine, MD; Etienne Birame Sène, MD; Oumar Kane, MD, PHD; Abdou Magib Gaye, MD

Summary

A 15-day-old boy presented with signs of heart failure and tamponade. Two-dimensional echocardiography revealed a complex intrapericardial mass (47 mm x 36 mm) with a large pericardial effusion compressing the heart. A CT scan defined the relationship with great vessels, pericardium and myocardium. Complete surgical resection was performed without complication. Histology of the tumor confirmed the presumptive imaging diagnosis of teratoma. Intrapericardial teratomas are rare primary cardiac tumors usually diagnosed in neonates and infants. They contain endodermic, mesodermic, and neuroectodermic germinal layers. Intrapericardial teratomas are usually benign tumors but may be life-threatening because of pericardial effusion and heart compression. Echocardiography was used to make a diagnosis by showing an intrapericardial heterogeneous mass compressing the heart. Bi-dimensional echocardiography was a performed exam in primary cardiac tumors di-

“A newborn with a teratoma has an excellent prognosis after excision. The diagnosis in utero with prenatal echography and resonance magnetic imaging in newborns prevents cardiac adiasy and improves the care.”

agnosis, but tomodesitometry (CT scan) and magnetic resonance imaging (MRI) have advantages in large tumors assessing the relationship between the tumor and adjacent tissues

Keys Words: tumor, teratoma, surgery, tamponade, cardiac

Introduction

Intrapericardial teratoma is a rare primary cardiac tumor usually diagnosed in newborns.¹ This tumor realized a compression of the cardiac cavities, the respiratory system and pericardial effusion. The authors report a case of intrapericardial tumor in a newborn revealed by cardiac tamponade.

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share?

Submit your manuscript to: **RichardK@CCT.bz**

NUCLEUS-X™

Balloon Aortic & Pulmonic
Valvuloplasty Catheter



Only U.S. Cleared
Balloon for
Balloon Aortic
Valvuloplasty





Need to Recruit a Pediatric Cardiologist?

Advertise in Congenital Cardiology Today, the only monthly newsletter dedicated to pediatric and congenital cardiologists.

Reach the most Board Certified or Board Eligible pediatric cardiologists worldwide.

Recruitment advertising includes full color in either the North American print edition, or the electronic PDF International edition.

Available in 1/3 and 1/2 page vertical Recruitment ad sizes. We can create the advertisement for you at no extra charge!

Contact:

Tony Carlson, Founder

Tel: +1.301.279.2005 or TCarlsonmd@gmail.com

the 7th post-operative day. Histopathology exam showed a benign mature teratoma with multiple cysts surrounded by conjunctive tissues, including endodermic, mesodermic, and neuroectodermic germinal layers.

Three months later, the child was asymptomatic with normal echocardiogram.



Figure 3: View of the tumor.



Figure 4: Teratoma after excision.

Discussion

Intrapericardial teratoma is a rare primary cardiac tumor.² This embryologic tumor is observed in newborns. Intrapericardial teratomas are usually benign tumors, but may be life-threatening because of pericardial effusion and heart compression.³ Today the majority of Intrapericardial teratomas are diagnosed in utero by prenatal echography.^{1,4} In this case, tamponade is due to late diagnosis, because echography is not commonly administered to low-income pregnant women.

Two dimensional echocardiography is the primary diagnostic imaging modality.⁵ In under development countries, explorations including magnetic resonance (MRI) or CT Scan are not accessible immediately for the majority of population. The only alternative for us in this emergency case of cardiac tamponade is pericardial drainage. MRIs and CT scans have advantages in assessing the relationship between large tumors and adjacent tissues.

The complete excision of the tumor was done 37 days after a diagnostic ultrasound. Surgery was performed with a beating heart. The bypass pump was on standby. In a few cases, these excisions are done with cardiopulmonary bypass and cross-clamping the aorta.¹ These tumors are usually removable and recurrences in mature teratomas are exceptional.



Medtronic

Technologies to Manage
Congenital Heart Disease

Every Step of the Way

**Melody® Transcatheter Pulmonary Valve
Ensemble® Transcatheter Valve Delivery System**

Indications: The Melody TPV is indicated for use in a dysfunctional Right Ventricular outflow Tract (RVOT) conduit (≥ 16 mm in diameter when originally implanted) that is either regurgitant (\geq moderate) or stenotic (mean RVOT gradient ≥ 35 mm Hg)

Contraindications: None known.

Warnings/Precautions/Side Effects:

- DO NOT implant in the aortic or mitral position.
- DO NOT use if patient's anatomy precludes introduction of the valve, if the venous anatomy cannot accommodate a 22-Fr size introducer, or if there is significant obstruction of the central veins.
- DO NOT use if there are clinical or biological signs of infection including active endocarditis.
- Assessment of the coronary artery anatomy for the risk of coronary artery compression should be performed in all patients prior to deployment of the TPV.
- To minimize the risk of conduit rupture, do not use a balloon with a diameter greater than 110% of the nominal diameter (original implant size) of the conduit for pre-dilation of the intended site of deployment, or for deployment of the TPV.
- The potential for stent fracture should be considered in all patients who undergo TPV placement. Radiographic assessment of the stent with chest radiography or fluoroscopy should be included in the routine postoperative evaluation of patients who receive a TPV.
- If a stent fracture is detected, continued monitoring of the stent should be performed in conjunction with clinically appropriate hemodynamic assessment. In patients with stent fracture and significant associated RVOT obstruction or regurgitation, reintervention should be considered in accordance with usual clinical practice.

Potential procedural complications that may result from implantation of the Melody device include: rupture of the RVOT conduit, compression of a coronary artery, perforation of a major blood vessel, embolization or migration of the device, perforation of a heart chamber, arrhythmias, allergic reaction to contrast media, cerebrovascular events (TIA, CVA), infection/sepsis, fever, hematoma, radiation-induced erythema, and pain at the catheterization site.

Potential device-related adverse events that may occur following device implantation include: stent fracture resulting in recurrent obstruction, endocarditis, embolization or migration of the device, valvular dysfunction (stenosis or regurgitation), paravalvular leak, valvular thrombosis, pulmonary thromboembolism, and hemolysis.

For additional information, please refer to the Instructions for Use provided with the product or call Medtronic at 1-800-328-2518 and/or consult Medtronic's website at www.medtronic.com.

Humanitarian Device. Authorized by Federal law (USA) for use in patients with a regurgitant or stenotic Right Ventricular Outflow Tract (RVOT) conduit (≥ 16 mm in diameter when originally implanted). The effectiveness of this system for this use has not been demonstrated.

Melody and Ensemble are trademarks of Medtronic, Inc.
UC201303735 EN © Medtronic, Inc. 2013;
All rights reserved.

The Melody® TPV offers children and adults a revolutionary option for managing valve conduit failure without open heart surgery.

Just one more way Medtronic is committed to providing innovative therapies for the lifetime management of patients with congenital heart disease.

Innovating for life.

“A newborn with a teratoma has an excellent prognosis after excision. The diagnosis in utero with prenatal echography and resonance magnetic imaging in newborns prevents cardiac adiasstoly and improves the care.”

Conclusion

A newborn with a teratoma has an excellent prognosis after excision. The diagnosis in utero with prenatal echography and resonance magnetic imaging in newborns prevents cardiac adiasstoly and improves the care.

References

1. De Geeter B; Kretz J G; Nisand I; Eisenmann B; Kieny M-Th; Kieny R. Intrapericardial teratoma in newborn infant: use of fetal echocardiography. *Ann Thorac Surg* 1983; 35: 664-6.
2. Adler RH, Taheri SA, Weintraub DF: mediastinal teratoma in infancy. *J Thorac Cardiovasc Surg* 1960; 30: 394.
3. Reynolds J L, Donahue J K, Pearce C W. Intrapericardial teratoma : a cause of acute pericardial effusion in infancy. *Pediatrics* 1969; 43: 71-78.
4. Laquay N, Ghazaouani S, Vaccaroni L, Vouhi P. Intrapericardial teratoma in newborn babies. *Eur Jour Cardiothorac Surg* 2003; 23: 642-4.
5. Summer TE, Crow JE, Klein A, Mc Kone RC, Weaver RL. Intrapericardial teratoma in infancy. *Pediatr Radiol* 1980; 10: 51-3.

CCT

Corresponding Author

*Amadou Gabriel Ciss, MD
Cardiovascular Surgeon; Specialist,
Congenital Heart Diseases; Maître
Assistant, Dakar University; Member,
African Society of Thoracic and
Cardiovascular Surgeons; Member,
French Society of Thoracic and
Cardiovascular Surgeons
University Hospital of Fann
BP 5035 Dakar Sénégal West Africa
Tel: 00 221 70 600 12 25
cissgaby@yahoo.fr*

*Momar Sokhna Diop, MD
Pap Adama Dieng, MD
Pap Salmane Ba, MD
Assane N'diaye, MD, PHD
Mouhamadou N'diaye, MD, PHD
Magaye Gaye, MD
Oumar Diarra, MD, PHD
Clinic of Thoracic and Cardiovascular
Surgery
University Hospital of Fann
BP 5035 Dakar Sénégal West Africa*

*Mohamed Leye, MD
Souleymane Diatta, MD
Arame Diallo, MD
Clinic of Cardiology
University Hospital of Fann
BP 5035 Dakar Sénégal West Africa*

*Fall Lamine, MD
Etienne Birame Sène, MD
Oumar Kane, MD, PHD
Intensive Care Unit
University Hospital of Fann
BP 5035 Dakar Sénégal West Africa*

*Abdou Magib Gaye, MD
Pathologist
Dakar University
BP 5035 Dakar Sénégal West Africa*

Letters to the Editor

Congenital Cardiology Today welcomes and encourages Letters to the Editor. If you have comments or topics you would like to address, please send an email to: LTE@CCT.bz, and let us know if you would like your comment

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share?

Submit your manuscript to:
RichardK@CCT.bz

- Title page should contain a brief title and full names of all authors, their professional degrees, and their institutional affiliations. The principal author should be identified as the first author. Contact information for the principal author including phone number, fax number, email address, and mailing address should be included.
- Optionally, a picture of the author(s) may be submitted.
- No abstract should be submitted.
- The main text of the article should be written in informal style using correct English. The final manuscript may be between 400-4,000 words, and contain pictures, graphs, charts and tables. Accepted manuscripts will be published within 1-3 months of receipt. Abbreviations which are commonplace in pediatric cardiology or in the lay literature may be used.
- Comprehensive references are not required. We recommend that you provide only the most important and relevant references using the standard format.
- Figures should be submitted separately as individual separate electronic files. Numbered figure captions should be included in the main Word file after the references. Captions should be brief.
- Only articles that have not been published previously will be considered for publication.
- Published articles become the property of the Congenital Cardiology Today and may not be published, copied or reproduced elsewhere without permission from Congenital Cardiology Today



ACHA - 6757 Greene Street, Suite 335 - Philadelphia, PA, 19119
P: (888) 921-ACHA - F: (215) 849-1261

A nonprofit organization which seeks to improve the quality of life and extend the lives of congenital heart defect survivors.

<http://achaheart.org>

Watch over 300 Live Case Videos, Presentations and Workshops Online from Leading Congenital and Structural Medical Meetings from Around the World - www.CHDVideo.com



- Transseptal Access Workshop from Cook Medical
- Workshop: Past Present and Future of Pediatric Interventions Cardiology - St. Jude & AGA Mmedical
- Symposium on Prevention of Stroke Clinical Trials at the Heart of the Matter - WL Gore Medical
- Imaging in Congenital & Structural Cardiovascular Interventional Therapies
- Morphology of The Atrial Septum
- Morphology of The Ventricular Septum
- Pre-Selection of Patients of Pulmonic Valve Implantation and Post-Procedural Follow-up
- Echo Paravalvular Leakage (PVL)
- ICE vs TEE ASD Closure in Children - PRO & CON ICE
- 3D Rotational Angiography - Why Every Cath Lab Should Have This Modality
- PICS Doorway to the Past - Gateway to the Future
- Follow-up From PICS Live Cases 2010 Presentation
- Intended Intervention - Transcatheter TV Implantation - *Live Case*
- Intended Intervention - LAA Closure Using Amplatzer Cardiac Plug Under GA & Real Time 3D
- Provided Intervention - LPA Stenting / Implantation of a Sapien Valve
- Intended Intervention - PV Implantation
- Intended Intervention - COA Stent Using Atrium Advanta V12 Covered Stent - *Live Case*
- Intended Intervention - ASD Closure - *Live Case*
- Intended Intervention - Transcatheter VSD Device Closure - *Live Case*
- Intended Intervention - COA Stenting Using Premounted Advanta V12 Covered Sten - *Live Case*
- Stunning Revelation - The Medical System is Changing - What Can You Do To Show Patients That Your Practice Does It Right? Patient Perspective
- Percutaneous Paravalvular Leak Closure Outcomes
- Intensive Management of Critically Ill Infants Undergoing Catheterization
- Off Label Device Usage - Careful!
- Evolving Hybrid Programs Communications is the Key
- Anti-Coagulation in Pediatric Interventions - Evidence Please! Are we Flying Blind?
- Shaping Your Structural Heart Interventional Career: How Do I Get the Necessary Training and Start a Program?
- Closure of Post-Infarct VSDs Limitations & Solutions
- Percutaneous Interventions During Pregnancy
- Percutaneous PVL Closure Techniques & Outcomes
- Percutaneous Therapy for MR
- Update on Percutaneous Aortic Valve Replacement: Time to Open the Flood Gates?
- 2011 PICS/AICS Achievement Award Presented To Horst Sievert, MD.
- The Cases I Learn The Most From
- If I was Starting Now I Would... Contemplating the Question.
- 10 Tips for the Trainee
- The Perfect CV
- Why I Left The Cath Lab
- Intended Intervention Occlude Right High Flow Superior Segment PAVM
- Intended Intervention Edwards Sapien XT Transcatheter Heart Valve - 3D Guidance - *Live Case*
- Closure of the Silent PDA - Indicated or Not
- Transcatheter Tricuspid Valve Replacement
- Recommendations For Device Closure of MVSDs
- Intended Intervention Diagnostic Angio/First Intervention Occlude Left Lower Lobe - *Live Case*
- Intended Intervention PFO Closure using Gore Helex Septal Occluder Under GA and TEE Guidance - *Live Case*
- Pulmonary Valvuloplasty - Pulmonary Oedema Presentation
- Intended Intervention LAA Closure Using the Amplatzer Cardiac Plug - *Live Case*
- Transcatheter Alternatives in Sick Neonates With Coartation of the Aorta; Balloon Angioplasty vs Surgery for Native Coartation of the Aorta in Newborns
- **and many more.... Plus, live cases from other major meetings**

Presented by **CONGENITAL CARDIOLOGY TODAY**




Image of the Month #5 – May 2013

Presented by The Archiving Working Group

Contributors: Jeffrey P. Jacobs, MD, Jorge M. Giroud, MD, Robert Anderson, MD, Vera D. Aiello, MD, Diane E. Spicer, BS

This is a special column that is published bi-monthly in *Congenital Cardiology Today* with contributors and images from the Archiving Working Group (AWG) of the *International Society for Nomenclature of Paediatric and Congenital Heart Disease*.


Please visit us at the AWG Web Portal at <http://ipccc-awg.net> and help in the efforts of the Archiving Working Group and the *International Society for Nomenclature of Paediatric and Congenital Heart Disease*.

The authors would like to acknowledge the Children's Heart Foundation (www.childrensheartfoundation.org) for financial support of the AWG Web Portal.

IPCCC:
07.10.01, 07.10.17, 12.14.02, 12.24.23

AEPC Derived Term:
Perimembranous ventricular septal defect (VSD) (07.10.01)

AWG Web Portal link for this series of images:
http://ipccc-awg.net/VSD_Page/VSD_07_10_17/VSD_07_10_17.html



Description: This view of the great vessels as they exit the ventricular mass shows the aorta arising slightly to the right of its usual position. The pulmonary trunk is dilated with a pulmonary artery band in place and a ligated arterial duct. The right atrium is dilated and there is right ventricular hypertrophy.
Contributor: Diane E. Spicer, BS

Ventricular septal defect (VSD) + malaligned outlet septum anteriorly (07.10.17)
Pulmonary trunk band (PA band) (12.14.02)
Ligation of patent arterial duct (PDA) (12.24.23)

EACTS-STIS Derived Term:


VSD, Type 2 (Perimembranous) (Paramembranous) (07.10.01)
VSD-modifier for infundibular septal morphology, VSD + malaligned outlet septum, Anterior deviation of infundibular septum (07.10.17).
PA Banding, Band on main pulmonary artery (12.14.02).
PDA closure, Surgical therapy, Ligation (12.24.23)

ICD10 Derived Term: Q21.0 Congenital malformations of the cardiac septa. Ventricular septal defect.

Commentary: Hearts with antero-cephalad malalignment of the muscular outlet, or conal septum are usually found in the setting of Tetralogy of Fallot. In the specimen shown here, however, the malalignment of the outlet septum did not cause sub-pulmonary obstruction. This arrangement, with an unobstructed outflow tract, is the lesion il-


lustrated by Victor Eisenmenger at the end of the nineteenth century, and known by many as the "Eisenmenger VSD". The very existence of this type of heart, with an unobstructed subpulmonary outlet, showed that something extra is needed over and above the anterocephalad deviation of the outlet septum so as to produce the phenotypic morphology of Tetralogy of Fallot. The additional feature in Tetralogy is an abnormal arrangement of the septoparietal trabeculations, which in most instances are additionally hypertrophied. It is the combination of the deviated outlet septum and the abnormal septoparietal trabeculations that produces the "squeeze" at the mouth of the subpulmonary infundibulum that is the essence of tetralogy of Fallot. We will discuss additional morphological aspects of the consequences of overriding of arterial valvar roots to the definition and description of holes between the ventricles is a future column from the Archiving Working Group.

Please visit us at the AWG Web Portal at <http://ipccc-awg.net/> and help in the efforts of the Archiving Working Group and the *International Society for Nomenclature of Paediatric and Congenital Heart Disease*.



Description: An image of the same heart demonstrating on the anterior apical view of the right ventricle the antero-superior malalignment of the outlet septum into the right ventricle with an unobstructed subpulmonary outlet. The aorta overrides the ventricular septum, although the aortic valve is not appreciated in this view. This defect is consistent with an Eisenmenger type of ventricular septal defect.

Contributor: Diane E. Spicer, BS



Description: In this anatomic view of the opened left ventricle, the free wall has been lifted away to demonstrate the aortic root. The aorta clearly overrides the interventricular septum with a good portion of the aortic valve supported in the right ventricle.

Contributor: Diane E. Spicer, BS



PICS-AICS
Pediatric and Adult Interventional Cardiac Symposium

SAVE THE DATE
JUNE 7-10, 2014
Marriott Chicago
DOWNTOWN
CHICAGO

WWW.PICSYMPOSIUM.COM



LIVE CASE DEMONSTRATIONS • ABSTRACT
SESSIONS • "MY NIGHTMARE CASE IN THE
CATH LAB" • HOT DEBATES • WORKSHOPS
• SMALLER BREAKOUT SESSIONS



RUSH UNIVERSITY
MEDICAL CENTER
Sponsored for CME credit by Rush University Medical Center

The authors would like to acknowledge the Children's Heart Foundation (www.childrens-heartfoundation.org/) for financial support of the AWG Web Portal.

CCT



Jeffrey P. Jacobs, MD
Archiving Working Group
Congenital Heart Institute of Florida,
St. Petersburg & Tampa, USA



Corresponding Author

Jorge M. Giroud, MD
Co-chairman, Archiving Working Group
Congenital Heart Institute of Florida &
Pediatrix Medical Group
St. Petersburg & Tampa, USA
jorgemgiroud@gmail.com



Robert Anderson, MD
Co-chairman, Archiving Working Group
Institute of Genetic Medicine, Newcastle
University, Newcastle upon Tyne,
United Kingdom



Vera D. Aiello, MD
Co-chairman, Archiving Working Group
Heart Institute (InCor). Sao Paulo
University School of Medicine, Brazil



Diane E. Spicer, BS
Senior Archivist, Archiving Working
Group
University of Florida, Department of
Pediatrics-Cardiology, Gainesville,
Florida and the Congenital Heart
Institute of Florida, St. Petersburg &
Tampa, FL USA



*And the members of the Archiving
Working Group of the International
Society for Nomenclature of Paediatric
and Congenital Heart Disease*

Letters to the Editor

Congenital Cardiology Today welcomes
and encourages Letters to the Editor. If
you have comments or topics you would
like to address, please send an email
to: LTE@CCT.bz, and let us know if you
would like your comment



Do You Use Medical Apps on Your Smartphone or Tablet?

Email us the names of some of your favorites and why.

Send them to: apps@CCT.bz



An Unusual Case of Prenatal Diagnosis of a Large Left Ventricular Diverticulum and Its Intermediate Term Outcome

By Resham Kaur, MD; Javier Paiz, RDCS; John Brownlee, MD; Umang Gupta, MBBS, DCH

Introduction

Congenital Ventricular Aneurysm or Diverticulum are very rare conditions. There are few reported cases of these lesions in the literature. Here we describe a rare case of Wide-mouthed Left Ventricular (LV) Free Wall Diverticulum that was identified by fetal echocardiography and subsequently followed in our clinic.

Case Report

A 24 week pregnant G4P2012 was referred for fetal echocardiography due to abnormality noted in the Left ventricular LV free wall and mild pericardial effusion.

The echocardiogram done in the clinic showed large LV free wall out-pouching with a wide mouth and some synchronous contractility with the rest of LV free wall. A very small pericardial effusion was seen. The LV systolic function appeared to be depressed with marked reduction in the contractility in and around the lesion. No arrhythmias were seen. No intracardiac thrombi were identified.

She was worked up extensively for conditions that could lead to dilated cardiomyopathy in fetus with no cause detected.

The patient was followed in the fetal clinic every 4 weeks by serial echocardiograms, the last being performed at 36 weeks of gestation (Figure 1). No changes were noted either in the size or the function of the LV and the diverticulum.

The fetus continued to show normal growth on prenatal examination and didn't develop hydrops and showed no arrhythmias. No mitral valve regurgitation was seen.

The baby was subsequently delivered through an uneventful spontaneous vaginal delivery weighing 3420 grams. At birth no other congenital defects were identified.

Figure 1. The four-chamber view at 36 week of gestational age showing the large diverticulum (Arrow) at the Left Ventricular Free Wall.



He was subsequently transferred to a transplant center for evaluation and underwent an extensive work up to rule out any genetic/metabolic abnormalities all of which were negative. He was started on carvedilol and enalapril and after a period of observation was discharged home. At the time of discharge patient was feeding very well, was very vigorous and was growing normally. No arrhythmias were identified at any time during the hospital stay.

He has subsequently been followed very closely in the cardiology clinic and at the time of his last visit was 7 months old and showed normal growth and development.

He has had no episodes of arrhythmias. His diverticulum continues to decrease in size relative to his LV chamber size (Figure 2) with continued improvement in contractility.

Discussion

Congenital Ventricular wall aneurysm or diverticulum is a very rare condition. There are few reported cases of congenital ventricular aneurysm in the literature.^{1,2,3,4,5,8} The finding of congenital diverticulum is even rarer.^{1,8,9,11,12} Many of these have been diagnosed after birth and only a handful have been diagnosed to date using fetal echocardiography.^{1,2,3,4,5,8,9,11}

The reasons for referral for fetal echocardiography in most cases have varied from abnormalities on the four-chamber views to hydrops/pericardial effusion to arrhythmias (including atrial arrhythmias) seen during routine prenatal ultrasound examination.^{1,2,3,4,5,8,9}

Our case was referred to us due to "abnormality" detected on the LV free wall and small pericardial effusion.

Most of these congenital diverticula are described as showing synchronous contractility with the rest of the chamber and usually have a narrow mouth, characteristics, that help differentiate them with aneurysms.¹⁰

In our case, the mouth of the lesion was wide, which had initially confounded us to believe it to be an aneurysm (Figure 1). Later imag-

Figure 2. Four-chamber view at 7 months of age showing the diverticulum (arrow) that appears to be much smaller relative to the Size of LV.







Figure 3 & Figure 4. Showing the diastolic and systolic frame of LV and diverticulum in parasternal short axis view. Note the synchronous contractility of the diverticulum with the rest of the LV.

ing proved it to have synchronous systolic contraction (Figure 3 & 4). The subsequent improvement in function after birth made the diagnosis of diverticulum more obvious.

Ever since they were first described a common association has been found between these diverticula and other congenital defects, most notably midline thoraco-abdominal defects.¹²

In our case, we didn't identify any other congenital defect.

A lot of focus has been placed on the outcome. However, literature remains inconclusive about the course and prognosis of these anomalies. Reported experience has varied from a relatively benign course to a more ominous presentation and outcome.^{1,2,3,4,5,6,7,8,9,10,11} The most common causes of death that have been reported seem to be arrhythmias or severe fetal hydrops.^{1,7}

Our patient remained stable throughout the pregnancy and showed normal intrauterine growth. His delivery remained uneventful and he has shown progressive improvement in his left ventricular function with decrease in the size of the diverticulum and improved contractility of the involved region.

References

1. Cavalle- Garrido T, Cloutier A, Harder J, Boutin C, Smallhorn JF: Evolution of fetal ventricular aneurysms and diverticula of the heart: an echocardiographic study. *Am J Perinatol.* 1997; (7): 393-400.
2. Matias A, Fredouille C, Nesmann C, Azancot A: Prenatal diagnosis of left ventricular aneurysm: a report of three cases and a review. *Cardiol Young.* 1999;9(2): 175-84.
3. Jacobson RL, Perez A, Meyer RA, Miodovnik M, Siddiqi TA: Prenatal diag-

“Ever since they were first described a common association has been found between these diverticula and other congenital defects...”

nosis of fetal left ventricular aneurysm: a case re-port and review. *Obstet Gynecol.* 1991; 78(3 Pt 2):525-8.

4. Sherman SJ, Leenhouts KH, Utter GO, Litaker M, Lawson P: Prenatal diagnosis of left ventricular aneurysm in the late second trimester: a case report. *Ultrasound* 1996; 7(6):456-7.
5. JHornberger LK,Dalvi B,Benacerraf BR: Prenatal sonographic detection of cardiac aneurysms and diverticula. *Ultrasound Med.* 1994; 13(12):967-70.
6. M K Mardini: Congenital diverticulum of the left ventricle. Report of two unusual cases. *Br Heart J.* 1984;51(3): 321-326.
7. Bernasconi A, Delezoide AL, Menez F, Vuillard E, Oury JF, Azancot A: Prenatal rupture of left ventricular diverticulum: a case report and review of the literature. *Prenat Diagn.* 2004; 24 (7): 504-7.
8. K. Balakumar: Prenatal diagnosis of left ventricular aneurysm. *Indian J Radiol Imaging.* 2009; 19(1): 84-86.
9. Perlitz Y, Mukary M, Lorber A, Ben-Ami M: Prenatal diagnosis of fetal cardiac right ventricular diverticulum disappearing at three months of age. A case report and literature review. *Fetal Diagn Ther.* 2009; 25(1):44-6.
10. Fernandez MS, Lopez A, Villa JJ, Lluna J, Miranda J: Cantrell's pentalogy: Report of four cases and their manage-

ment. *Pediatr Surg Int* 1997; 12: 428-31.

11. Prefumo F, Bhide A, Thilaganathan B, Carvalho JS: Fetal con-genital cardiac diverticulum with pericardial effusion: two cases with different presentations in the first trimester of pregnancy. *Ultrasound Obstet Gynecol.* 2005; 25(4):405-8.
12. Knight L, Neal WA, Williams HJ, Huseby TL, Edwards JE. Congenital left ventricular diverticulum: Part of a syndrome of cardiac anomalies and midline defects. *Minn Med* 1976; 59: 372-5.

CCT

Corresponding Author

Resham Kaur, MD
Driscoll Children's Heart Center
Driscoll Children's Hospital
3533 S. Alameda St.
Corpus Christi, TX 78411, USA
resham_thethi@yahoo.com
Tel: 36.877.9474; Fax: 361.855.9518
Corresponding email:
author-umangupta@yahoo.com

Javier Paiz, RDCS
Driscoll Children's Heart Center
Driscoll Children's Hospital
Corpus Christi, TX 78411, USA

John Brownlee, MD
Driscoll Children's Heart Center
Driscoll Children's Hospital
Corpus Christi, TX 78411, USA

Umang Gupta, MBBS, DCH
Driscoll Children's Heart Center
Driscoll Children's Hospital
Corpus Christi, TX 78411, USA

Do Not Close the Door on Closing the Door!

By Stefan C. Bertog, MD; Laura Vaskelyte, MD; Ilona Hofmann, MD; Marius Hornung, MD; Simon Lam, MD; Sameer Gafoor, MD; Horst Sievert, MD

The foramen ovale, a remnant of the fetal circulation remains patent in approximately 25% of the population based on autopsy and echocardiographic studies.^{1,2} Though population-based longitudinal studies do not support a higher risk of stroke in the presence of a patent foramen ovale (PFO),^{3,4} patients with cryptogenic stroke are more likely to have a PFO (at least 40%) than the general population.⁵ Moreover, passage of thrombus originating from the venous system via a PFO into the systemic circulation is physiologically plausible and has been demonstrated in a number of case reports unequivocally demonstrating paradoxical embolism via an interatrial communication.⁶⁻⁹ Intuitively, anticoagulation, antiplatelet therapy or closure of a PFO should abolish or reduce the risk of stroke from paradoxical embolism. Nevertheless, significant controversy remains as to how a patient should be treated who has experienced a stroke in the absence of an identifiable cause with the exception of a PFO.

For the affected individual, there are four options: no therapy at all, anticoagulation, antiplatelet therapy and surgical or percutaneous closure. Unfortunately, a study that provides us with a definitive answer does not exist.

Let us briefly examine the evidence for either of the available approaches. For the first option, no therapy, we do not know what the risk of a recurrent event is because the natural history (no therapy) of individuals with a PFO and cryptogenic stroke is unknown. Regarding the second and third options, we do not know if either anticoagulant or antiplatelet therapy reduces the rate of recurrent events, as neither has ever been compared with no therapy in a randomized controlled trial. Likewise, we do not have any conclusive evidence to support anticoagulant therapy over antiplatelet therapy or vice versa. Finally, percutaneous or surgical closure has never been compared to no therapy in a randomized controlled fashion. To conclude, in the absence of a randomized controlled trial comparing therapy to no therapy, we do not know if any therapy is better than no therapy. Does this imply that our response to the individual with a cryptogenic stroke and PFO should be that no therapy is required?

“To conclude, in the absence of a randomized controlled trial comparing therapy to no therapy, we do not know if any therapy is better than no therapy. Does this imply that our response to the individual with a cryptogenic stroke and PFO should be that no therapy is required?”

If we believe that a patient who has experienced a cryptogenic stroke in the absence of a PFO should be treated with antiplatelet therapy because the most likely mechanism of the event (with the exception of the rare pure atherosclerotic debris, fat, tumor or air embolism) is platelet aggregation and thrombus formation with or without embolization, then we should, likewise, favor antiplatelet therapy for the patient in question because these mechanisms may also be present in patients with cryptogenic stroke in the presence of a PFO. The data in support of this approach stems from trials in patients with strokes regardless of mechanism who benefit from aspirin administration. Hence at least an aspirin should be recommended. However, some would argue that, in patients with cryptogenic stroke and PFO, aspirin might not be sufficient because it does not provide adequate protection against thrombus formation in the venous circulation and paradoxical embolization via a PFO. On a theoretical basis, only PFO closure or anticoagulation may have merit in preventing venous thrombus and paradoxical embolism. No dedicated randomized trials comparing aspirin and anticoagulation with warfarin in patients with PFO and cryptogenic stroke exist. Available data stems from subgroup analysis of patients with PFO in stroke trials comparing aspirin and warfarin. Though there was a trend favoring warfarin over aspirin, this was not statistically significant.^{10,11} However, the trials were underpowered to detect a difference. Hence, in the absence of more convincing data, there is no conclusive evidence to favor warfarin over aspirin for the prevention of recurrent events in patients with PFO and cryptogenic stroke. How about PFO closure? There are

non-randomized studies comparing percutaneous PFO closure to medical therapy suggesting a lower stroke recurrence rate after closure^{12,13} and there is surgical data that suggests a very low rate of event recurrence after surgical PFO closure.^{14,15} More importantly, there are now three trials comparing percutaneous PFO closure to medical therapy, CLOSURE 1 (Evaluation of the STARFlex Septal Closure System in Patients with a Stroke and/or Transient Ischemic Attack Due to Presumed Paradoxical Embolism through a Patent Foramen Ovale),¹⁶ PC (Clinical Trial Comparing Percutaneous Closure of the Patent Foramen Ovale Using the Amplatzer PFO Occluder with Medical Treatment in Patients with Cryptogenic Embolism)¹⁷ and RESPECT (Randomized Evaluation of Recurrent Stroke Comparing PFO Closure to Established Current Standard of Care Treatment)¹ in patients with cryptogenic stroke and PFO.

In CLOSURE 1, patients were randomized to closure with the STARFlex device (NMT Medical) and in PC and RESPECT to closure with the Amplatzer PFO occluder (St. Jude Medical) versus medical therapy. In CLOSURE 1, 909 patients (mean age 45 years) were enrolled. After 2 years, though the rate of the primary event (stroke, transient ischemic attack and 30-day all cause and 30-day to 2 year neurologic mortality) and rate of recurrent strokes were numerically lower in patients who underwent PFO closure, the difference was not statistically significant.¹⁶

Three important aspects regarding CLOSURE 1 are worth mentioning. First, the closure rates were lower with the STARFlex device (no or minimal residual shunt: 86%) than with the Amplatzer PFO occluder in both PC (96%) and RESPECT (94%). Second, the follow-up was limited to 2 years. Third, the rates of thrombus formation and atrial fibrillation appear to be higher after STARFlex implantation than after implantation of the most commonly used Amplatzer devices.¹⁸ This may reduce or (in the case of device associated thrombus or atrial fibrillation) offset the potential benefits of PFO closure. In PC, 414 patients (mean age 44-45 years) were enrolled. After a mean of 4 years of follow-up, there was no difference in the primary endpoint (composite of death, nonfatal stroke, transient ischemic attack or peripheral embolism) in an intention-to-treat analysis.¹⁷ However, there was a trend toward a lower rate of strokes (0.5% versus 2.4%) in patients treated with PFO closure ($p=0.13$) by the definition used in the trial and a strong trend towards a reduction in

stroke risk based on the definition used in RESPECT ($p=0.07$). Nine-hundred and eighty patients (mean age 46 years) were enrolled in RESPECT.¹ Once again, there was a strong trend towards a reduction in stroke recurrence in patients treated with PFO closure compared to those treated medically (after a mean follow-up time of approximately 2 years) in an intention-to-treat analysis. In addition, there was a significant reduction in stroke recurrence in an as-treated analysis ($p=0.007$). Though this type of analysis may be confounded by non-random factors leading to deviation from the treatment protocol, it is worth mentioning that among the nine patients who experienced a recurrent stroke assigned to the device group, three had not undergone device closure, one of whom had a stroke after enrollment but before scheduled device implantation. The second patient, after treatment assignment, decided not to proceed with device implantation. The third patient underwent bypass surgery and surgical PFO closure after assignment to the device group but prior to planned implantation. Importantly, in subgroup analyses, there was a significant reduction in the primary endpoint in favor of PFO closure (compared with medical therapy only) in patients with an interatrial septal aneurysm or large right-to-left shunt.

Particularly the stronger treatment effect with a larger shunt size is in line with the observed higher event rate in patients with cryptogenic strokes and large shunts,^{19,20} both of which support the notion that PFO's promote strokes via paradoxical embolism and the rationale of its closure. In addition, there was a strong trend toward a reduction in the primary endpoint in patients whose index infarct had been located more superficially favoring an embolic etiology.

The event rates continued to separate further in favor of closure with time. For example, the stroke rates were 1.3%, 1.6% and 2.2% in the closure group and 1.7%, 3.0% and 6.4% in the medically treated group after one, two and five years, respectively. Though the absolute difference in event rates is small, the relative difference at 5 years is 65%. Assuming that the difference between the treatment arms is a true difference and not a result of chance, in young patients this may translate into a large number of strokes prevented. In addition, no device erosion or thrombus formation directly related to the device occurred in either the PC or RESPECT trial supporting the safety of closure with the Amplatzer PFO occluder.

If one focuses on strictly predefined intention-to-treat findings of these trials and allows only results with a p -value of <0.05 to be meaningful, discounts subgroup and as-treated analyses and ignores potential differences among PFO devices, withholding PFO closure from patients with a cryptogenic stroke may be the logical consequence. However, based on above described data, it appears more reasonable to conclude that the risk of harm due to device closure (i.e. device associated thrombus formation and device erosion) is very low. Further, there may be a benefit of closure compared with medical therapy particularly in certain patient subgroups. Therefore, though PFO closure in an unselected population is not appropriate, categorically closing the door to PFO closure is equally inappropriate. For example, mandating no therapy or aspirin therapy only in a young individual with a cryptogenic stroke, PFO with a large interatrial shunt or atrial septal aneurysm and superficial infarct seen on brain imaging seems to be premature. Rather, careful patient selection and a fair discussion with the patient outlining the therapeutic options, risks and limitations to current knowledge, but ultimately respecting the patient's decision seem to be most appropriate.

In this context, we would like to draw attention to our upcoming *ICI/CSI Conference* in Frankfurt Germany this June, where topics such as the above and many others related to congenital and structural

JUNE MEETING FOCUS

CSI 2013 and iCi 2013

Frankfurt am Main, Germany

June 26, 2013 (ICI 2013); June 27-29, 2013 (CSI 2013)

www.csi-congress.org or www.ici-congress.org

CSI Course Director: Horst Sievert, MD; CardioVascular Center Frankfurt (Germany)

CSI Co-Course Directors: Mario Carminati, MD; Instituto Policlinio San Donato (Italy); Ilona Hofman, MD; CardioVascular Center Frankfurt (Germany); Shakeel A Qureshi, MD; Evelina Children's Hospital (UK); Neil Wilson, MD; John Radcliffe Hospital (UK);

For the 16th time, **CSI 2013, the annual international Congress on Congenital and Structural Interventions of the Heart**, will take place from June 27th to 29th 2013 in Frankfurt am Main, Germany, with **ICI 2013, the Imaging in Cardiovascular Interventions** meeting taking place one day before on June 26th.

Starting in 1996 as a small one-day workshop on catheter closure of congenital shunts with 77 attendees, the **CSI Congress** has now become an established event for interventional adult and pediatric cardiologists, radiologists, cardiovascular surgeons, catheter laboratory nurses and technicians from all over the world. Over time, the variety of topics grew and, today, our attendees share their experience with procedural techniques, devices and latest technologies in all areas of interventional cardiology. From day one, CSI focused on the transmission of LIVE cases from cath labs around the world, including: CardioVascular Center Frankfurt, Instituto Policlinio San Donato, Sri Jayadeva Institute of Cardiology and Cedars-Sinai Medical Center. With more than 800 participants in 2012, the **CSI Congress** serves as a growing, but still familiar gathering.

In keeping with our tradition, transmission of LIVE cases from various cath labs will continue to be the main focus. Allowing experienced interventionalists, as well as beginners, to profit from the "see one, do one, teach one" concept, **CSI** offers a mixture of complex challenging as well as routine procedures. Every participant at the **Congress** center has the opportunity to directly interact with the interventionalist in the cath lab, assuring that questions directly related to the procedure are answered under real-time conditions. In addition, comprehensive lectures by leading interventionalists allow participants to keep up-to-date on specific topics of interest. Particularly, discussion of controversial topics is fostered. Due to the flexible concept of LIVE transmissions and scientific lectures, attention is directed to every step of the procedure from scientific background, patient selection, planning and execution to postprocedural care.

This year's **CSI** is again accompanied by **Imaging in Cardiovascular Interventions (iCi 2013)**, taking place one day before the CSI conference on June 26th. The main focus of **iCi** is the introduction and demonstration of new imaging methods and modalities. Main focus is the discussion of the utility of existing and novel imaging techniques in diagnosing congenital and structural heart disease and its application to optimize patient selection. Procedural strategy and choice and sizing of interventional equipment will be discussed in detail. In addition, new imaging methods and modalities (for example 3D-ICE or Echo-Fluoro-Overlay) will be introduced and demonstrated. During **iCi**, LIVE demonstrations will take place and simulator training will be available.

heart disease will be discussed at length by experts in the respective subjects. For more information visit: www.ici-congress.org and www.csi-congress.org.

References

- Putala J, Metso AJ, Metso TM, et al. Analysis of 1008 consecutive patients aged 15 to 49 with first-ever ischemic stroke: the Helsinki young stroke registry. *Stroke; a journal of cerebral circulation* 2009;40:1195-203.
- Hagen PT, Scholz DG, Edwards WD. Incidence and size of patent foramen ovale during the first 10 decades of life: an autopsy study of 965 normal hearts. *Mayo Clin Proc* 1984;59:17-20.
- Meissner I, Khandheria BK, Heit JA, et al. Patent foramen ovale: innocent or guilty? Evidence from a prospective population-based study. *J Am Coll Cardiol* 2006;47:440-5.
- Di Tullio MR, Sacco RL, Sciacca RR, Jin Z, Homma S. Patent foramen ovale and the risk of ischemic stroke in a multiethnic population. *J Am Coll Cardiol* 2007;49:797-802.
- Lechat P, Mas JL, Lascault G, et al. Prevalence of patent foramen ovale in patients with stroke. *N Engl J Med* 1988;318:1148-52.
- Falk V, Walther T, Krankenberg H, Mohr FW. Trapped thrombus in a patent foramen ovale. *Thorac Cardiovasc Surg* 1997;45:90-2.
- Koullias GJ, Elefteriades JA, Wu I, Jovin I, Jadbabaie F, McNamara R. Images in cardiovascular medicine. Massive paradoxical embolism: caught in the act. *Circulation* 2004;109:3056-7.
- Fukumoto A, Yaku H, Doi K, et al. Images in cardiovascular medicine. Continuous thrombus in the right and left atria penetrating the patent foramen ovalis. *Circulation* 2005;112:e143-4.
- Mascarenhas V, Kalyanasundaram A, Nassef LA, Lico S, Qureshi A. Simultaneous massive pulmonary embolism and impending paradoxical embolism through a patent foramen ovale. *J Am Coll Cardiol* 2009;53:1338.
- Homma S, Sacco RL, Di Tullio MR, Sciacca RR, Mohr JP. Effect of medical treatment in stroke patients with patent foramen ovale: patent foramen ovale in Cryptogenic Stroke Study. *Circulation* 2002;105:2625-31.
- Mohr JP, Thompson JL, Lazar RM, et al. A comparison of warfarin and aspirin for the prevention of recurrent ischemic stroke. *N Engl J Med* 2001;345:1444-51.
- Schuchlenz HW, Weihs W, Berghold A, Lechner A, Schmidt R. Secondary prevention after cryptogenic cerebrovascular events in patients with patent foramen ovale. *Int J Cardiol* 2005;101:77-82.
- Wahl A, Juni P, Mono ML, et al. Long-term propensity score-matched comparison of percutaneous closure of patent foramen ovale with medical treatment after paradoxical embolism. *Circulation* 2012;125:803-12.
- Devuyst G, Bogousslavsky J, Ruchat P, et al. Prognosis after stroke followed by surgical closure of patent foramen ovale: a prospective follow-up study with brain MRI and simultaneous transesophageal and transcranial Doppler ultrasound. *Neurology* 1996;47:1162-6.
- Ruchat P, Bogousslavsky J, Hurni M, Fischer AP, Jeanrenaud X, von Segesser LK. Systematic surgical closure of patent foramen ovale in selected patients with cerebrovascular events due to paradoxical embolism. Early results of a preliminary study. *Eur J Cardiothorac Surg* 1997;11:824-7.
- Furlan AJ, Reisman M, Massaro J, et al. Closure or medical therapy for cryptogenic stroke with patent foramen ovale. *N Engl J Med* 2012;366:991-9.
- Meier B, Kalesan B, Mattle HP, et al. Percutaneous closure of patent foramen ovale in cryptogenic embolism. *N Engl J Med* 2013;368:1083-91.
- Taaffe M, Fischer E, Baranowski A, et al. Comparison of three patent foramen ovale closure devices in a randomized trial (Amplatzer versus CardioSEAL-STARflex versus Helex occluder). *Am J Cardiol* 2008;101:1353-8.
- Lippmann H, Rafferty T. Patent foramen ovale and paradoxical embolization: a historical perspective. *Yale J Biol Med* 1993;66:11-7.
- Homma S, Di Tullio MR, Sacco RL, Mihalatos D, Li Mandri G, Mohr JP. Characteristics of patent foramen ovale associated with cryptogenic stroke. A biplane transesophageal echocardiographic study. *Stroke; a journal of cerebral circulation* 1994;25:582-6.

CCT

Stefan C. Bertog, MD
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany

Laura Vaskelyte, MD
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany

Ilona Hofmann, MD
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany

Marius Hornung, MD
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany

Simon Lam, MD
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany

Sameer Gafoor, MD
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany

Corresponding Author

Prof. Dr. Med. Horst Sievert
CardioVascular Center Frankfurt
Seckbacher Landstrasse 65
60389 Frankfurt
Germany
Phone: +49-69-4603-1344
Fax: +49-69-4603-1343

info@cvcfrankfurt.de



Archiving Working Group
International Society for Nomenclature of
Paediatric and Congenital Heart Disease
ipccc-awg.net

Medical News, Products and Information

Philips Announces FDA Clearance for its EchoNavigator Live Image-Guidance Tool

Royal Philips Electronics announced that it has received 510(k) clearance from the US Food and Drug Administration (FDA) to market its innovative EchoNavigator live image-guidance tool (EchoNavigator). This world-first technology helps interventional cardiologists and cardiac surgeons perform minimally-invasive structural heart disease repairs by providing an intelligently integrated view of live X-ray and 3D ultrasound images. Following the CE marking of EchoNavigator in Europe, Philips will now be able to introduce the system globally.¹ The first systems have already been installed in Europe and the US.

Philips' EchoNavigator has been developed in response to a clear upward trend in the use of both X-ray imaging and 3D cardiac ultrasound imaging (also known as echocardiography, or echo for short) during structural heart disease procedures - an area of interventional cardiology that is growing at around 40% per year. During such procedures, ultrasound imaging provides critical insights into the heart's soft tissue anatomy, while X-ray imaging has particular strengths in visualizing the catheters and the heart implants.

Working in collaboration with partner hospitals in Europe and the US, Philips designed EchoNavigator to address the unique challenges associated with working with live X-ray and 3D ultrasound images simultaneously. Accurately recognizing the heart structures from these medical images takes years of training and experience, and the use and interpretation of both imaging techniques during the procedures can be challenging, especially when manipulating and steering the catheters that carry the implants. Moreover, the communication during the critical moment of the procedure between the interventional cardiologist or surgeon steering the catheters, and the echocardiographer operating the 3D ultrasound equipment, has been identified as particularly demanding.

"Together with Philips, we set out to bring two separate medical imaging techniques together in a way that provides clear visual guidance," said Professor John Carroll, MD,

Interventional Cardiologist, University of Colorado Hospital, Denver, US. "A world-first, EchoNavigator is enabling us to use X-ray images combined with real-time 3D ultrasound images to navigate catheters and deploy implants in the right position in the heart, making such treatments more straightforward."

Philips' EchoNavigator will enable clinicians to perform procedures more efficiently by providing intelligently integrated X-ray and 3D ultrasound images into one intuitive and interactive view, as well as providing easy-to-use system navigation and better communication between the multidisciplinary team carrying out the procedure. As a result, EchoNavigator helps save valuable time and enhances patient care.

"We have learned that ideally two live imaging technologies are needed to guide catheter-based repairs to the heart and a multidisciplinary team is needed to perform it," said Professor Roberto Corti, MD, Interventional Cardiologist, University Hospital Zurich, Switzerland. "This adds to the complexity of such procedures. The development of a more sophisticated imaging technology such as EchoNavigator will definitely provide us with a better understanding of the complex structures of the heart and its repair."

"As the global market leader in interventional cardiology, we have worked with our partners to lead the way with pioneering solutions such as our real-time 3D ultrasound technology and more recently our HeartNavigator navigation tool," said Gene Saragnese, CEO for Imaging Systems at Philips Healthcare. "EchoNavigator is further evidence of our commitment to transforming healthcare through the introduction of innovations that enables best-in-class minimally-invasive procedures."

Philips offers a comprehensive interventional cardiology portfolio that includes hybrid operating room solutions and imaging solutions, plus advanced interventional tools that work smoothly in sync with them. In 2011, Philips introduced HeartNavigator, a procedure planning and image guidance tool optimized for minimally invasive aortic heart valve replacements. Philips' new EchoNavigator unites the company's

strengths in interventional X-ray and ultrasound, as well as its strengths in clinical information solutions. Philips introduced Live 3D Trans Esophageal Echo (Live 3D TEE) technology² as an industry-first in 2007 on its iE33 ultrasound system and last year introduced CX50 xMATRIX - the world's first compact portable ultrasound system to incorporate Live 3D TEE technology.³

"In the emerging field of complex structural heart disease interventions, the information obtained by merging imaging technologies, as now possible with HeartNavigator and EchoNavigator, will be of tremendous value to the interventionalist, and in turn to the patient," said Dr. Carlos Ruiz, MD, Director of the Structural and Congenital Heart Disease program, Department of Interventional Cardiology at Lenox Hill Hospital, New York (US).

¹ Not available for sale in Brazil, Saudi Arabia, Serbia, and Australia pending clearance by the relevant regulatory authorities.

² With this technology an ultrasound scan is taken by placing a special 3D ultrasound transducer into the patient's esophagus so that it is positioned close to the heart valves.

³ Both iE33 xMATRIX and CX50 xMATRIX support EchoNavigator.

Lower-Profile SAPIEN XT Transcatheter Heart Valve Associated With Improved Procedural Outcomes

Preliminary Results From Inoperable Cohort of The PARTNER II Trial Presented at ACC 2013

SAN FRANCISCO, CA—(Marketwire - March 10, 2013) - Edwards Lifesciences Corporation (NYSE:EW), the global leader in the science of heart valves and hemodynamic monitoring, announced that preliminary results from The PARTNER II Trial demonstrated similar one-year outcomes in mortality and major clinical events between the Edwards SAPIEN XT transcatheter aortic valve and the Edwards SAPIEN valve, yet fewer vascular events with the lower-profile SAPIEN XT valve. These data from The PARTNER II Trial studying transcatheter aortic valve replacement (TAVR) in inoperable patients with severe, symptomatic aortic stenosis were presented today as a late-



CHINA CALIFORNIA HEART WATCH

Providing free clinical care and doctor training to medically underserved farming communities in Yunnan, a province of wild beauty but also one of the poorest regions in China.

We have a beautiful volunteer-created hard cover photo book taking you on a journey through the landscapes and peoples of Yunnan, showing the world in which China Cal works and the lives we've affected. Available to all donors who make a tax deductible donation of at least \$150. Visit www.chinacal.org to order.



The PARTNER II Trial, Inoperable Cohort (Cohort B) (i)

Outcome	SAPIEN			SAPIEN		
	XT	SAPIEN	p value	XT	SAPIEN	p value
All-cause mortality - %	3.5	5.1	0.36	22.5	23.7	0.706
Stroke: all - %	4.3	4.1	0.88	5.9	5.7	0.935
Vascular events: major - %	9.6	15.5	0.04			
Vascular events: minor - %	5.0	7.4	0.23			

Outcome	SAPIEN			SAPIEN		
	XT	SAPIEN	p value	XT	SAPIEN	p value
All-cause mortality - %	3.5	5.1	0.36	22.5	23.7	0.706
Stroke: all - %	4.3	4.1	0.88	5.9	5.7	0.935
Vascular events: major - %	9.6	15.5	0.04			
Vascular events: minor - %	5.0	7.4	0.23			

(i) All percents for mortality and stroke data in this press release are Kaplan-Meier estimates.

breaking clinical trial at the American College of Cardiology's (ACC) 62nd Annual Scientific Session in San Francisco.

"We are very pleased to see improved outcomes in both the SAPIEN and SAPIEN XT patients, as compared to results from earlier trials," said Larry L. Wood, Edwards' Corporate Vice President, Transcatheter Heart Valves. "The SAPIEN XT valve was designed to reduce complications of the TAVR procedure, and we believe this has been demonstrated by today's results. Edwards is proud to lead the advancement of cutting-edge innovations for these high-risk patients, supported by rigorous clinical evidence."

The PARTNER II Trial enrolled 560 patients deemed inoperable for traditional open-heart surgery at 28 hospitals in the United States between April 2011 and February 2012. Patients were randomized to receive one of the two Edwards transcatheter aortic heart valves: 276 received the SAPIEN valve, and 284 received the SAPIEN XT valve.

The FDA approved the SAPIEN valve in November 2011 for the treatment of inoperable patients, and expanded the indication to high-risk surgical patients in October 2012. The SAPIEN XT valve is an investigational device not yet available commercially in the United States.

Edwards anticipates submitting data from the inoperable cohort (Cohort B) of The PARTNER II Trial to the United States Food and Drug Administration (FDA) in the second quarter. The company expects to complete enrollment in the intermediate risk cohort (Cohort A) of The PARTNER II Trial mid-year.

Background Information on The PARTNER Trial, Inoperable Cohort (Cohort B)

The PARTNER Trial studied 358 patients with severe, symptomatic aortic stenosis deemed inoperable for traditional open-heart surgery, and enrolled between April 2007 and March 2009. Patients were evenly randomized to receive either the Edwards SAPIEN valve or standard therapy. Please note that the results from The PARTNER Trial and The PARTNER II Trial are not directly comparable.

ECG Screening for Competitive Athletes Would not Prevent Sudden Death

The risk of cardiovascular sudden death was very small and only about 30% of the incidence were due to diseases that could be reliably detected by pre-participation screening, even with 12-lead ECGs, according to research in a US high school athlete population presented March 10th at the *American College of Cardiology (ACC) Scientific Sessions*.

Sudden death in young competitive athletes due to cardiovascular disease is an important community issue, which could impact the design of population-based screening initiatives. The frequency with which these tragic events occur impacts considerations for selecting the most appropriate screening strategy. Currently, athletes are assessed through a healthcare professional performing a physical exam and reviewing the individual's clinical history.

"Screening initiatives for high school-aged athletes have the potential to impact 10-15 million young adults in the US," says the study's lead author, Barry J. Maron, MD, Director of the Hypertrophic Cardiomyopathy Center at the Minneapolis Heart Institute Foundation in Minneapolis. "This is a controversial issue because some are suggesting that all young competitive athletes should be screened with a 12-lead ECG screening, which would be a massive and costly undertaking. Also, we do not have any evidence to show whether this is clinically necessary."

To assess this need, Maron and his colleagues interrogated the forensic case records of the U.S. National Registry of Sudden Death in Athletes over a 26-year period (1986-2011) to identify those events judged to be cardiovascular in origin occurring in organized competitive interscholastic sports participants in Minnesota. There were more than 4.44 million sports participations, including 1,930,504 individual participants among 24 sports.

There were 13 incidences of sudden deaths in high school student-athletes related to physical exertion during competition (7) or at practice (6). The ages were 12 to 18 and each was a white male. Most common sports involved were basketball, wrestling or cross-country running. Sudden deaths occurred in 1 out of 150,000 participants.

Autopsy examination documented cardiac causes in 7 of the 13 deaths. In only 4 athletes (31%) could the responsible cardiovas-

MEDICAL CONFERENCES 2013	ICI 2013 Imaging in Cardiovascular Interventions Frankfurt, Germany 	CSI 2013 Congenital, Structural & Valvular Interventions Frankfurt, Germany 	TRENDS ASIA-PACIFIC 2013 Catheter-based Neurohumoral Remodeling Taiwan, The Republic of China 	LAA 2013 Left Atrial Appendage Closure Frankfurt, Germany 	www.ici-congress.org www.csi-congress.org www.csi-trends.org www.csi-laa.org
	JUNE 26	JUNE 27-29	SEPTEMBER 21	NOVEMBER 22-23	
	www.ici-congress.org	www.csi-congress.org	www.csi-trends.org	www.csi-laa.org	

cular diseases be reliably detected by history, physical exam or 12-lead ECG, which is equivalent to 1 in 1 million participants.

"This very low event rate does not warrant changing the current national screening strategy, especially because only one-third of the deaths would have been detectable through additional screening," says Maron. "These findings demonstrate that these tragic events are rare. In addition to these data, no evidence in the medical literature has shown that ECGs reduce mortality in a broad-based screening effort."

Close-to-the-Heart Catheters Safer for Hospitalized Children

Location, location, location. A new Johns Hopkins Children's Center study shows the real-estate mantra also holds true when it comes to choosing correct catheter placement in children.

The research findings, described online March 18th in *JAMA Pediatrics*, show that catheters in children inserted in a vessel in the arm or leg and not threaded into a large vein near the heart are nearly four times as likely to dislodge, cause vein inflammation or dangerous blood clots as are catheters advanced into major vessels near the heart.

A peripherally inserted central venous catheter, or PICC line, is a tube placed into a small blood vessel, usually in the arm, and threaded toward a major blood vessel near the lungs and heart to serve as a temporary portal for medications, nutrients or fluids. However, clinicians sometimes forego threading close to the heart and leave the PICC line in a peripheral vein in the arm or leg instead — a choice dictated by the ease and speed of placement or a child's overall condition or anatomy.

The study findings, however, suggest that leaving the device in a non-central vein should only be done as last resort, the researchers say.

"Clinicians should carefully weigh the ease and speed of non-central vein placement against the higher complication risk that our study found goes with it," says senior investigator and pediatric infectious disease specialist Aaron Milstone, MD, MHS.

Non-central, smaller veins, especially those in the arm, are narrower, thinner and more prone to injury than major vessels near the heart, the researchers say. Thus, a catheter can easily damage the protective coating on the walls of such veins and encourage the formation of blood clots that, in the worst-case scenario, can dislodge and travel to the lungs or heart, causing a pulmonary embolism or heart damage.

Conducted among more than 1,800 pediatric patients hospitalized at Johns Hopkins over six years, the study found that such non-centrally positioned catheters accounted for a mere 16% of the central lines, but for 44% of all complications that led to catheter removal.

Children in the study cumulatively underwent more than 2,500 catheter insertions, of which more than 500 — one in five — had to be removed due to complications. Three-quarters of problems stemmed from mechanical malfunction such as device breakage or dislodgement, clot formation or blood vessel inflammation. The rest were due to infection, which traditionally has been the greatest worry with central lines. Vein location, however, played no role in infection risk, the research showed.

Despite the higher risks seen with non-centrally position catheters, overall complications rates dropped significantly over the six-year study period, a trend that should get a further boost by emerging technologies, the investigators say.

"We are already adopting new technologies that render PICC placement near the heart easier, safer and faster, and which will drive complications rates further down," says Leslie Gosey, RN, MS, leader of the pediatric catheter-insertion team at Johns Hopkins.

The study was funded by the National Institute of Allergy and Infectious Diseases (K23 AI081752) and the National Institute of Nursing Research (R03 NR012558).

Co-investigators included: Ketan Jumani, BDS, MPH; Sonali Advani, MBBS, MPH; and Nicholas Reich, PhD, all from Johns Hopkins.

Depression in Kids Linked to Cardiac Risks in Teens

Teens who were depressed as children are far more likely than their peers to be obese,

smoke cigarettes and lead sedentary lives, even if they no longer suffer from depression.

The research, by scientists at Washington University School of Medicine in St. Louis and the University of Pittsburgh, suggests that depression, even in children, can increase the risk of heart problems later in life.

The researchers reported their findings March 15th at the annual meeting of the *American Psychosomatic Society* in Miami, FL.

"Part of the reason this is so worrisome is that a number of recent studies have shown that when adolescents have these cardiac risk factors, they're much more likely to develop heart disease as adults and even to have a shorter lifespan," says first author Robert M. Carney, PhD, a professor of psychiatry at Washington University. "Active smokers as adolescents are twice as likely to die by the age of 55 than nonsmokers, and we see similar risks with obesity, so finding this link between childhood depression and these risk factors suggests that we need to very closely monitor young people who have been depressed."

Researchers have known for years that adults with depression are likely to have heart attacks and other cardiac problems, but it hasn't been clear when risk factors for heart disease such as smoking, obesity and sedentary lifestyle join forces with depression to increase the risk for heart problems.

"We know that depression in adults is associated with heart disease and a higher risk of dying from a heart attack or having serious complications," Carney says. "What we didn't know is at what stage of life we would begin to see evidence of this association between depression and these cardiac risk factors."

The researchers studied children who had participated in a 2004 study of the genetics of depression. At the time, their average age was 9. The investigators surveyed 201 children with a history of clinical depression, along with 195 of their siblings who never had been depressed. They also gathered information from 161 unrelated age- and gender-matched children with no history of depression.

In 2011, when the study participants had reached the age of 16, the researchers surveyed them again, looking at rates of smok-



Archiving Working Group
International Society for Nomenclature of
Paediatric and Congenital Heart Disease
ipccc-awg.net

ing, obesity and physical activity in all three groups of adolescents.

"Of the kids who were depressed at age 9, 22% were obese at age 16," Carney says. "Only 17% of their siblings were obese, and the obesity rate was 11% in the unrelated children who never had been depressed."

Carney and his colleagues found similar patterns when they looked at smoking and physical activity.

"A third of those who were depressed as children had become daily smokers, compared to 13% of their nondepressed siblings and only 2.5% of the control group," he says.

In terms of physical activity, the teens who had been depressed were the most sedentary. Their siblings were a bit more active, and members of the control group were the most active.

When the researchers took a closer look and used statistical methods to eliminate other factors that potentially could have influenced smoking or obesity rates in the depressed children, Carney's team found that the effects of depression grew even more pronounced.

"The siblings of depressed children were five times more likely to smoke than members of the study's control group, so depression wasn't the only risk factor for smoking," he explains. "But the depressed children in the study were another 2½ times more likely to smoke than their non-depressed siblings."

And the heart disease risk factors were more common in formerly depressed children whether or not they still were clinically depressed at the time of the second survey. In fact, Carney says, for most of the adolescents, depression was in remission by the time the second survey was conducted in 2011, with only 15% of them reporting depression.

The results suggest that any history of depression in childhood appears to influence the presence of cardiac risk factors during adolescence, according to Carney.

"Depression seems to come first," he says. "It's playing an important, if not a causal, role. There may be some related genetic influences that give rise to both depression and to

heart disease, or at least to these types of cardiac risk behaviors, but more study will be required before we can draw any firm conclusions about that."

Carney RM, Rottenberg J, Freedland KE, Kovacs M. Childhood major depressive disorder and cardiovascular risk factors in adolescents. Presented at the *Annual Meeting of the American Psychosomatic Society*, March 15th, 2013.

Washington University School of Medicine's 2,100 employed and volunteer faculty physicians also are the medical staff of Barnes-Jewish and St. Louis Children's hospitals. The School of Medicine is one of the leading medical research, teaching and patient care institutions in the nation, currently ranked sixth in the nation by *U.S. News & World Report*. Through its affiliations with Barnes-Jewish and St. Louis Children's hospitals, the School of Medicine is linked to BJC HealthCare.

Novel Approach To Treating Children With Irregular Heart Beat

Newswise – A new retrospective study from The Children's Hospital at Montefiore (CHAM) found similar safety and efficacy at lower cost using a novel three-catheter approach for ablation in children with Wolff-Parkinson-White (WPW) Syndrome, a condition where an extra, abnormal electrical pathway in the heart causes rapid heart rate or tachycardia. The current standard of care for ablation of left-sided accessory pathways in children with WPW is a five-catheter approach and patients treated with this approach served as the age and gender control match in this study. These data were presented March 10th at *The American College of Cardiology's 62nd Annual Scientific Session* in San Francisco.

"With increased concern over healthcare expenditures, it is paramount that we maintain quality, efficacy and safety while also maintaining or reducing costs," said Robert H. Pass MD, Director of the Pediatric Cardiac Catheterization Laboratory, CHAM, and Associate Professor of Pediatrics at Albert Einstein College of Medicine of Yeshiva University. "Our study showed that using a three-catheter approach in this patient population can reduce the cost by more than 20% with comparable outcomes."

Wolff-Parkinson-White affects one to three people in every 1,000 worldwide. The long-term treatment is catheter ablation, where radiofrequency energy is applied to the extra electrical pathway, disabling the heart's ability to beat irregularly. Typically five catheters are used to monitor both sides of the heart, but in patients with WPW who have a left-sided pathway, an electrocardiogram test prior to the procedure enables surgeons to accurately predict the location of the pathway. That information allows for reduction in the number of catheters used on the right side of the heart.

The four-year study included 56 children with WPW who were treated with catheter ablation. One group (n=28) received the new approach, where three catheters were inserted through the patient's groin and neck and threaded through to the heart where the extra electrical pathway was identified and disabled, and had a 100% success rate. The control group (n=28) received the standard, five-catheter approach and had a 96% success rate. The average catheter cost was lower in the three-catheter group (\$1,940 vs. \$2,620).

"In Wolff-Parkinson-White ablation cases where we can predict that a pathway is located on the left side of the heart, we are confident in this method and believe it should be applied more often," said Dr. Pass. "Most importantly, with the growing cost of healthcare, we believe any such effort to provide outstanding and consistent outcomes while garnering significant savings is paramount."

The Pediatric Cardiac Catheterization Laboratory is part of the Pediatric Heart Center at CHAM. It was the first hybrid catheterization lab in the New York metropolitan region, allowing for maximal efficiency and coordination of care for children requiring both surgical and interventional procedures. The Pediatric Heart Center is recognized as a world leader in providing advanced cardiovascular care for young patients with congenital heart diseases, treating patients of all ages, from newborns to adults, from minor arrhythmias to heart transplantation.

For more information please visit www.montefiore.org and www.montekids.org.



CHDResources.org
Free downloadable
patient education materials

A service of
California Heart Connection
a nonprofit support network
caheartconnection.org
info@caheartconnection.org
877-824-3463

Beware: Newly Recognized Heart Cardiomyopathy Is Not Always Benign - Largely Present in Women, 'Broken Heart Syndrome' is Often Triggered by Stress

Even though a newly recognized cardiomyopathy, which mainly impacts women, is typically treatable, Tako-tsubo cardiomyopathy can also be deadly when compounded by other co-morbidities, such as heart failure, according to a study was presented March 9 at the *American College of Cardiology (ACC) Scientific Sessions*.

This condition, formally known as Tako-tsubo Cardiomyopathy (TTC) and informally known as stress cardiomyopathy or Broken Heart Syndrome, has abrupt onset of symptoms and is characterized by a distinctive left ventricular (LV) contraction profile. Ninety percent of the time, this condition affects women, who are usually middle-aged and older, and the condition usually is triggered by a stressful event.

"Although TTC is typically reversible and considered to have favorable clinical outcomes, we have identified an important subset of patients, particularly those with severe heart failure and hypotension, who can have a substantial mortality risk," says the study's lead author Scott W. Sharkey, MD, a research cardiologist at the Minneapolis Heart Institute Foundation and a physician at the Minneapolis Heart Institute® at Abbott Northwestern Hospital in Minneapolis. "It's also important that physicians are aware this is not a rare a condition, as it is present in nearly 10% of women who present to the hospital with suspected heart attacks."

MHIF researchers reviewed 250 TTC patients who presented to the Minneapolis Heart Institute at Abbott Northwestern Hospital between 2001 and 2012. Then, they segregated those TTC patients presenting with particularly severe heart failure and very low pressure, or hypotension (systolic blood pressure < 100 mm Hg), who required supportive treatment.

They found that severe hypotensive heart failure occurred in 45 patients. In this subset, 9 female patients died in-hospital despite aggressive treatment intervention, representing the only TTC-related hospital deaths in the 250 patient cohort.

Therefore, Sharkey and his colleagues concluded that TTC is not necessarily

a benign condition. Severe hypotensive heart failure of severity necessitating vasopressor and/or intra-aortic balloon pump occurs in nearly 20% of patients. Also, all TTC-related hospital deaths occurred in the hypotensive heart failure subgroup with an overall mortality of 3.5%.

Importantly, triggering physical stressors related to severe co-morbid non-cardiac conditions (8) or advanced age (1) were present in all 9 non-survivors, Sharkey notes.

"Unfortunately, there are not any guidelines or criteria to instruct diagnosis and treatment of these patients at this time," says Sharkey. "Therefore, this study could be a starting point for this process, as it provides a more complete profile of the clinical spectrum of TTC and provides useful guidance for the effective management of these acutely ill patients."

To raise additional awareness and improve care of these patients, he adds that guidelines would be helpful at this time, in order to standardize diagnosis and treatment across varied healthcare settings.

Congenital Cardiology Today

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share?

Submit your manuscript to:
RichardK@CCT.bz

Letters to the Editor

Congenital Cardiology Today welcomes and encourages Letters to the Editor. If you have comments or topics you would like to address, please send an email to: **LTE@CCT.bz**, and let us know if you would like your comment

CONGENITAL CARDIOLOGY TODAY

© 2013 by Congenital Cardiology Today
(ISSN 1554-7787-print; ISSN 1554-0499-online).
Published monthly. All rights reserved.

Publication Headquarters:

8100 Leaward Way, Nehalem, OR 97131
USA

Mailing Address:

PO Box 444, Manzanita, OR 97130 USA
Tel: +1.301.279.2005; Fax: +1.240.465.0692

Editorial and Subscription Offices:

16 Cove Rd, Ste. 200, Westerly, RI 02891
USA

www.CongenitalCardiologyToday.com

Publishing Management:

- Tony Carlson, Founder, President & Sr. Editor - TCarlsonmd@gmail.com
- Richard Koulbanis, Group Publisher & Editor-in-Chief - RichardK@CCT.bz
- John W. Moore, MD, MPH, Medical Editor - JMoore@RCHSD.org
- Virginia Dematatis, Assistant Editor
- Caryl Cornell, Assistant Editor
- Loraine Watts, Assistant Editor
- Chris Carlson, Web Manager
- William Flanagan, Strategic Analyst
- Rob Hudgins, Graphic Designer - RobH@CCT.bz

Editorial Board: Teiji Akagi, MD; Zohair Al Halees, MD; Mazeni Alwi, MD; Felix Berger, MD; Fadi Bitar, MD; Jacek Bialkowski, MD; Philipp Bonhoeffer, MD; Mario Carminati, MD; Anthony C. Chang, MD, MBA; John P. Cheatham, MD; Bharat Dalvi, MD, MBBS, DM; Horacio Faella, MD; Yun-Ching Fu, MD; Felipe Heusser, MD; Ziyad M. Hijazi, MD, MPH; Ralf Holzer, MD; Marshall Jacobs, MD; R. Krishna Kumar, MD, DM, MBBS; John Lamberti, MD; Gerald Ross Marx, MD; Tarek S. Momenah, MBBS, DCH; Toshio Nakanishi, MD, PhD; Carlos A. C. Pedra, MD; Daniel Penny, MD, PhD; James C. Perry, MD; P. Syamasundar Rao, MD; Shakeel A. Qureshi, MD; Andrew Redington, MD; Carlos E. Ruiz, MD, PhD; Girish S. Shirali, MD; Horst Sievert, MD; Hideshi Tomita, MD; Gil Wernovsky, MD; Zhuoming Xu, MD, PhD; William C. L. Yip, MD; Carlos Zabal, MD

Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors, and are not necessarily the views of Congenital Cardiology Today.



The Barth Syndrome Foundation

P.O. Box 974, Perry, FL 32348

Tel: 850.223.1128

info@barthsyndrome.org

www.barthsyndrome.org

Symptoms: Cardiomyopathy, Neutropenia, Muscle Weakness, Exercise Intolerance, Growth Retardation



TINY HEARTS

inspired

**HYBRID LABS WITH ACCESS
FOR BIG TEAMS.**

Fixing a heart from birth through adulthood takes big teams working together. So we examined the needs of leading clinicians when designing our hybrid solutions. The result: our Infinix™-i with 5-axis positioners and low profile detectors, stays out of the way, but right where needed, providing the best possible access to patients. To lead, you must first listen. medical.toshiba.com



youtube.com/toshibamedical



[@ToshibaMedical](https://twitter.com/ToshibaMedical)