

CONGENITAL CARDIOLOGY TODAY

Timely News and Information for BC/BE Congenital/Structural Cardiologists and Surgeons

December 2017; Volume 15; Issue 12
North American Edition

IN THIS ISSUE

Introducing Allogeneic Human Mesenchymal Stem Cell-based Therapy to Improve Clinical Outcomes in Patients with Hypoplastic Left Heart Syndrome

By Sunjay Kaushal, MD; Helina Mehta, MD; Kristopher Deatruck, MD; Michael Slack, MD; Joshua Hare, MD
~Page 1

Tetralogy of Fallot with a Double Aortic Arch: A Case Report

By Gregory Aird, BA; Joel Hayden, BA; Randy Richardson, MD
~Page 8

Medical News, Products & Information

~Page 14

MEDICAL MEETINGS

Device Therapies for Heart Failure
Dec. 15-16, 2017; Berlin, Germany
www.csi-congress.org/dhf.php?go=0

NeoPREP - An Intensive Review & Update of Neonatal/Perinatal Medicine, Sponsored by the American Academy of Pediatrics & the AAP Section on Neonatal-Perinatal Medicine
Jan. 20-26, 2018; Atlanta, GA USA
shop.aap.org/live-activities

CSI Asia Pacific
Jan. 31 - Feb. 3, 2018; Ho Chi Minh City, Vietnam
www.csi-congress.org/csi-asia-pacific.php

15th International Conference on Pediatric and Pediatric Cardiology
Feb. 19-20, 2018; Paris France
pediatriccardiology.conferenceseries.com/europe/

ACC 67th Annual Scientific Session & Expo
Mar. 10-12, 2018; Orlando, FL USA
<https://accscientificsession.acc.org/Information-Pages/future-meetings>

CONGENITAL CARDIOLOGY TODAY

Editorial and Subscription Offices

16 Cove Rd, Ste. 200
Westerly, RI 02891 USA
www.CongenitalCardiologyToday.com

Follow on Twitter @CCardiology

Recruitment as: pages 5, 15, 16, 17, 18

Introducing Allogeneic Human Mesenchymal Stem Cell-based Therapy to Improve Clinical Outcomes in Patients with Hypoplastic Left Heart Syndrome

By Sunjay Kaushal, MD; Helina Mehta, MD; Kristopher Deatruck, MD; Michael Slack, MD; Joshua Hare, MD

Introduction

Hypoplastic Left Heart Syndrome (HLHS) is characterized by significant underdevelopment of the left side of the heart, which is defined by hypoplasia of the left ventricle associated with severe mitral and aortic stenosis or atresia, and hypoplasia of the ascending aorta and aortic arch. Since there is essentially no left ventricle, the right ventricle pumps blood to both pulmonary and systemic circulations. After delivery, survival depends on a Patent Ductus Arteriosus (PDA) and nonrestrictive Atrial Septal Defect (ASD), as well as the absence of aortic restriction (coarctation) of PDA blood flow to the coronary arteries via the ascending aorta.

Until the introduction of the first stage of surgical palliation by Norwood in 1981, HLHS was universally fatal within the first weeks of life. Currently, treatment strategies include cardiac transplantation or staged surgical palliation, consisting of a series of operations whereby the right ventricular outflow is reconstructed to provide systemic circulation. The first stage of palliation is now known as the Norwood operation and is performed within the first week of life, with the tripartite goal of providing: unobstructed systemic

blood flow, unrestricted mixing of atrial blood, and restrictive pulmonary blood flow. The second stage is broadly termed a superior cavopulmonary anastomosis, and can be constructed as either a "Bidirectional" Glenn, or a Hemi-Fontan between 4 to 6 months of age, with the goal of establishing a direct systemic venous – pulmonary arterial connection, and unloading the volume overload imposed by the systemic-pulmonary arterial connection imposed at the stage I operation. The third stage is universally known as the Fontan operation, and creates a total cavopulmonary connection. This operation results in the entirety of the systemic venous blood to be directed to the pulmonary arterial circulation, providing separation of the systemic arterial and venous circulations, while directing the full systemic venous flow to the pulmonary circulation. This may be performed as early as 18 months, but is more commonly done between 2 and 4 years of age. The overall objective of the Fontan circulation is to rely on the right ventricle to support the systemic circulation and for the systemic venous return to be pulled through the pulmonary circulation in the absence of a ventricular pumping chamber, relying on the vis-a-tergo and negative intrathoracic pressure to facilitate pulmonary circulation.

Approximately 949 infants with HLHS are born in the United States annually, and 318 die during the neonatal period for a neonatal mortality of 33%.¹ In spite of dramatic

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share? Submit your manuscript to:
RichardK@CCT.bz

NOW ENROLLING:

Harmony™ Transcatheter Pulmonary Valve (TPV) Clinical Study

WHAT IS THE MEDTRONIC HARMONY TPV CLINICAL STUDY?

The study is a multi-center prospective, non-randomized, interventional pre-market study in the United States. The purpose of the study is to evaluate the safety and effectiveness of the Harmony TPV system in patients who have congenital heart disease and are clinically indicated for pulmonary valve replacement. The trial will involve up to 40 subjects implanted at 10 study centers.

WHO CAN PARTICIPATE?

Patients who have pulmonary regurgitation:

- Severe pulmonary regurgitation by echocardiography, or
- Pulmonary regurgitant fraction $\geq 30\%$ by cardiac magnetic resonance imaging

Patients who have a clinical indication for surgical placement of a RV-PA conduit or prosthetic pulmonary valve:

- Subject is symptomatic secondary to pulmonary insufficiency (e.g., exercise intolerance, fluid overload), or
- Right ventricular end diastolic volume index ≥ 150 mL/m², or
- Subject has RVEDV:LVEDV Ratio ≥ 2.0

HOW CAN I LEARN MORE ABOUT THE HARMONY TPV CLINICAL STUDY?

For additional information about the program, please contact Medtronic at:
RS.HarmonyTPVClinicalStudy@medtronic.com

CAUTION: Investigational Device. Limited by Federal Law (USA) to Investigational Use.



medtronic.com

710 Medtronic Parkway
Minneapolis, MN 55432-5604
USA
Tel: (763) 514-4000
Fax: (763) 514-4879
Toll-free: (800) 328-2518

LifeLine
CardioVascular Technical Support
Tel: (877) 526-7890
Tel: (763) 526-7890
Fax: (763) 526-7888
rs.cstechsupport@medtronic.com




Figure 1. cMRI of an ELPIS patient at baseline. Allogeneic mesenchymal stem cells will be injected to the right ventricle of patients with Hypoplastic Left Heart Syndrome at the time of the bidirectional cavopulmonary connection operation administered intramyocardially at 6-10 injection sites (blue X's = sites of injection).

improvements in pre-, intra- and postoperative care, the mortality rate of these infants is as high as 25% to 35% during the first year of life and the overall 5-year survival rate remains limited with reported survivals over 50%-60%. Even with cardiac transplantation remaining as the only current viable alternative for patients with failing RV function, survival rate continues to remain poor after HLHS surgical staged palliation and transplantation.^{2,3,4} After the Fontan completion operation, the circulatory state is that of chronic systemic venous hypertension. This can lead to multiple clinically difficult and life-limiting complications, including hepatic congestion and cirrhosis, plastic bronchitis, and protein-losing enteropathy. Furthermore, even with optimal care in infancy and childhood, all patients with a Fontan will ultimately experience right heart failure, which may be related to increasing pressure in the pulmonary vascular bed or failure of the systemic right ventricle due to the much higher afterload. Given these surgical limitations, innovative treatment options to regenerate and remodel the RV myocardium to make it more capable of sustaining systemic circulation remain a significant unmet medical need.^{5,6,7}

We are currently one of the first pediatric centers pioneering the clinical application of stem cell therapy for infants with HLHS in a trial entitled "Allogeneic Human Mesenchymal Stem Cell Injection in Patients with Hypoplastic Left Heart Syndrome: An Open Label Pilot Study (ELPIS)." In this investigation, we are evaluating the intramyocardial delivery of adjunctive cell-based therapy by directly injecting allogeneic human mesenchymal stem cells (allo-MSCs) during the Stage II operation. To our knowledge, this is the first randomized clinical trial to use allogeneic stem cells in children with HLHS in the world.

Significance of the Study

We aim to overlay a novel cell therapeutic strategy on the staged surgical procedures that HLHS patients typically undergo. During the Stage II superior cavopulmonary operation, allogeneic mesenchymal stem cells will be injected intramyocardially into the right ventricle.




Figure 2. ELPIS Enrollment Plan.

Primary Objective

Evaluate the safety and feasibility of intramyocardial injection of allogeneic mesenchymal cells during the Stage II operation.

Secondary Objective

Observe effects on clinical outcomes including right ventricular myocardial function, severity of tricuspid regurgitation, incidence of serious adverse events, re-hospitalizations, changes in health status, the need for transplantation, or mortality.

Methodology

Study Design and Sample Size

We plan to enroll a total of thirty patients with HLHS in a staged enrollment process. In this open-labeled study, a maximum of 20

“We aim to overlay a novel cell therapeutic strategy on the staged surgical procedures that HLHS patients typically undergo. During the Stage II superior cavopulmonary operation, allogeneic mesenchymal stem cells will be injected intramyocardially into the right ventricle.”

Test	Baseline	Day 2	Week 1	Week 2	Week 4	Week 16	Week 24	Week 36	Week 48
Demographic Medical History	SOC								
PE Vital Signs and AE's	SOC	SOC	SOC	SOC	SOC	SOC	SOC	SOC	SOC
Chemistry:									
Urinalysis	SOC	SOC					SOC		SOC
Comprehensive Metabolic Panel (CMP)	SOC	SOC							
Troponin I		SOC							
Brain Natriuretic Peptide (BNP)	SOC		SOC				R		R
Hematology:									
CBC w/diff, includes platelet	SOC	SOC							
PTT (if pt on anti-coags)*	SOC*	SOC*							
PT/INR (if pt on anti-coags)	SOC*	SOC*							
Miscellaneous Labs:									
Immune Monitoring: CD markers	R	R					R		R
Immune Monitoring: cPRA	SOC						R		R
HLA Typing	R								
Exosomes	R	R	R						
Endothelial Progenitor Cells	R		R				R		
Diagnostic Imaging:									
Cardiac MRI	R						R		R
CT Chest/Abdomen/Pelvic (First 10 pts only)									R
Electrocardiogram (EKG)	SOC	SOC	SOC	SOC	SOC	SOC	SOC	SOC	SOC
Transthoracic Echocardiogram (TTE)	SOC	SOC	SOC	SOC	SOC	SOC	SOC	SOC	SOC
Cardiac Catheterization	SOC								

Figure 3. ELPIS Schedule for standard of care (SOC) and research events (R).

patients will eventually receive intramyocardial injection of the allogeneic mesenchymal stem cells and 10 control patients with no cell injection. The enrollment of the patients will occur in two staged groups: Group A and Group B. In Group A, 10 consecutive HLHS patients will be initially enrolled in the allogeneic MSCs treatment arm to determine feasibility and safety. After six months of the last enrolled patient in Group A, all Group A patients will be assessed in order to determine whether the methodology is feasible and safe, including the harvesting, processing, and administering of the allogeneic MSCs. Thereafter, Group B will start enrolling a total of 20 HLHS patients, who will be randomized to the treatment and control arms in a 1:1 ratio, respectively, in order to have 10 allogeneic MSCs treated patients and 10 control patients. At the completion of this Phase I clinical study, the total enrolled cohort will be 20 patients treated with allogeneic MSCs and 10 patients in the control arm.

Study Setting and Period

Recruitment of subjects for the study are ongoing at the University of Maryland Children's Heart Program and Johns Hopkins Hospital. We will be recruiting subjects at additional sites in the near future, including Children's Healthcare of Atlanta.

Inclusion Criteria

In order to participate in this study, a patient must meet all of the inclusion criteria:

1. Subjects with HLHS (all types) requiring Stage II surgery.

Exclusion Criteria

In order to participate in this study, a patient must not:

1. Have HLHS and restrictive or intact atrial septum.
2. Be undergoing the Norwood procedure that does not have HLHS.
3. Have significant coronary artery sinusoids.
4. Require mechanical circulatory support prior to Stage II operation.
5. Have an underlying evidence of arrhythmia requiring anti-arrhythmia therapy.
6. Have a parent or guardian unwilling or unable to comply with necessary follow-up.
7. Be serum positive for HIV, hepatitis BsAg or viremic hepatitis C.
8. Be unsuitable for inclusion in the study in the opinion of the investigators.
9. Need for concomitant surgery for aortic coarctation or tricuspid valve repair.

Sample Size Considerations

Analyses of bioactivity will be exploratory in nature in order to aid in selection of dosage, endpoints, time points, and sample size determination for subsequent larger trials, but is sufficiently powered to determine safety.

Therapeutic Stem Cell Intervention

During the Stage II operation when the patient is on cardiopulmonary bypass, previously harvested, isolated, and

expanded allogeneic mesenchymal stem cells will be delivered into the right ventricle, directly into the myocardium using a 27-gauge needle syringe. The stem cell transplant occurs after the completion of the repair but before separating from cardiopulmonary bypass. The process and manufacturing of the allogeneic mesenchymal stem cells will be delivered as specified in the POSEIDON-DCM Phase I/II Trial (IND #: 14419; NCT01392625), led by Dr. Joshua M. Hare (ISCI / University of Miami Miller School of Medicine, FL). The allogeneic hMSC cells will have undergone control testing on the final enriched cell product prior to administration and the cell doses employed will be 2.5 x 10⁵ cells per kg of recipient body weight (5 million / 20kg). The entire dose of the cells, 600 microliters, will be divided and delivered in 6-10 intramyocardial injections. The intramyocardial injection of cells will take approximately 5 to 10 minutes, with a minimal addition to the total bypass time.^{8,9}

Plan for Analysis

All subjects will be cared for according to our standard postoperative protocols. All subjects will receive post-treatment assessments completed at set intervals between two days and 48 weeks after cell injection. Vital signs, complete physical examination, 12 lead electrocardiogram, and complete TTE as well as selected blood work will be included at two days, 24 weeks, and 48 weeks. Also, 24-hour Holter monitor (Month 6 and Month 12), and AE monitoring will be included. The only additional

assessment that is not part of the routine standard of care for follow-up after the Stage II or III operated HLHS patients are the following: cardiac MRI scan performed at pre-operation, 24 and 48 weeks post-cell injection; immune monitoring for graft rejection at two days, 24 weeks, and 48 weeks. The following markers will be used for analysis to assess for activated T-cells based upon a CD3+CD25+ or CD3+CD69+ phenotype; biomarkers will be performed at 2 days, 1 week, 4 weeks, 24 weeks, and 48 weeks.

Primary Endpoints

- We will measure safety and feasibility of intramyocardial delivery of allogeneic mesenchymal stem cells in subjects with HLHS after one year of injections. Note: We will monitor major adverse cardiac events including: death, sustained/symptomatic ventricular tachycardia requiring intervention with inotropic support, aggravation of heart failure, new myocardial infarction, unplanned cardiovascular option for cardiac tamponade and infection in the first month after injection, and serially afterwards.¹⁰

Secondary Endpoints

- To measure improvements of right ventricular function: change from baseline in right ventricular function by measuring right ventricular ejection fraction or fractional area change (RVEF/FAC), right ventricular end-diastolic volume, right ventricular end-systolic volume, right ventricular end-systolic diameter, and tricuspid regurgitation as measured by serial echocardiograms and MRI scans.¹⁰
- To measure incidence of mortality or need for transplantation after the BDCPA operation up to one year follow-up.¹⁰
- To measure changes in somatic growth velocity over time (weight, height, head circumference) from the BDCPA operation out to 12 months post-op.¹⁰
- To measure assessments of co-morbidity: cardiovascular mortality, all-cause mortality, cardiovascular morbidity, re-hospitalizations, and need for heart transplantation.¹⁰

Discussion

The objective of the ELPIS Trial is to advance the clinical application of MSCs in the treatment of HLHS patients and to potentially uncover the mechanism by which MSCs can improve ventricular function in this unique patient subset. In Phase I/II adult clinical trials, we have previously shown the safety and preliminary efficacy of allogeneic MSC treatment in adult ischemic patients by the proposed remodeling mechanisms of direct cell-cell interactions with myocardial cells and indirect paracrine effects in order to reduce myocardial fibrosis, stimulate angiogenesis, and stimulate endogenous c-kit+ cardiac progenitor cells (CPCs) in the left ventricle (LV).^{11,12} We have recently shown MSC injections preserve RV ventricular dysfunction by activating endogenous pathways of ventricular remodeling in a swine model of RV pressure overload, replicating features of HLHS.^{13,14} We hypothesize that MSCs will facilitate remodeling of the RV in HLHS patients by both direct and indirect mechanisms and improve ventricular function. We will overlay our novel therapeutic MSC strategy on the stage II bidirectional cavopulmonary anastomosis operation in HLHS patients and rigorously validate potential mechanisms for RV improvement.¹³



PEDIATRIC INTERVENTIONAL CARDIOLOGIST FACULTY POSITION

The Division of Pediatric Cardiology at the University of Utah School of Medicine and based at Primary Children's Hospital is recruiting a BE/BC Pediatric Cardiologist with subspecialty training in Interventional Cardiology, providing focused care to patients with congenital heart disease needing cardiac catheterization for diagnosis and management. The interventional cardiologist will join a division of 29 faculty and a team of 2 pediatric interventional cardiologists, a dedicated nurse practitioner, and experienced cath lab personnel, while working in a growing, nationally ranked Heart Center. The division has a flourishing catheterization program (640 cases last year) and coordinates care with cardiothoracic surgery, cardiac intensive care, and cardiac anesthesia. In addition, the division's thriving programs in fellowship training and research will provide the candidate with tremendous opportunities for teaching and participation in basic, translational, and/or clinical research (our program is one of the core participating centers in the Pediatric Heart Network funded by the NIH).

The successful candidate will receive a faculty appointment in the Department of Pediatrics on the Clinical or Tenure track at the academic level commensurate with experience and qualifications. The University of Utah offers an excellent benefits package, including 20.2% retirement contributions that vest immediately and excellent health care choices. The Department also offers an education loan repayment program. The local area provides an exceptional quality of life with immense cultural and recreational opportunities.

To read more about this opportunity and apply, please visit:

- **Pediatric Interventional Cardiologist:**
<http://utah.peopleadmin.com/postings/69669>

For additional information, please contact Lloyd Tani, M.D., Division Chief, at Lloyd.Tani@hsc.utah.edu.

The University of Utah is an EOE/AA employer and educator

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share? Submit your manuscript to: RichardK@CCT.bz

MSCs have been thoroughly investigated in animal models, and shown the ability to regenerate the heart directly through formation of new tissue and indirectly through paracrine effects.^{20,21} MSCs are an attractive candidate for stem cell therapy for multiple reasons. First, MSCs have well-defined characteristics due to their ability to differentiate into selected terminally differentiated lineages in vitro and their expression of specific, well-established markers (CD90, CD105, CD73).^{18,19} Second, MSCs are reproducibly isolated from bone marrow and expand robustly in vitro.^{18,19} Third, since MSC engraftment occurs at low rates with <0.1% of injected cells, the current paradigm suggests that MSCs interact with the injured native cells in order to limit tissue destruction or enhance repair by a variety of mechanisms. Cross-talk mechanisms, by which MSCs exert effects on various native signaling pathways include:

- a. secretion of bioactive proteins that act in a paracrine or autocrine fashion;
- b. upregulation of genes that attenuate destructive inflammation and facilitate repair and;
- c. secretion of extracellular microvesicles (EVs) that contain proteins and microRNAs.²²

In the injured heart, MSCs remodel the injured or distressed myocardium by modulating endogenous remodeling, attenuating fibrosis, promoting neovessel formation, stimulating cardiomyocyte proliferation, and activating endogenous c-kit+ CPCs.²² Fourth, MSCs have been studied in both cardiac and other conditions with more than 120 clinical trials, listed on clinicaltrials.gov, and have repeatedly been shown to be safe in clinical studies. Finally, MSCs have recently been evaluated in clinical trials in multiple forms of adult cardiac disease, including acute myocardial infarction and chronic heart failure, and treatment has decreased myocardial fibrosis and improved clinical status.^{8,9,11,12}

Allogeneic MSC injections are as effective as autologous MSCs in promoting functional remodeling of the treated ventricle and did not increase immunophenotyping in adult patients with ischemic cardiomyopathy patients.^{11,12} Follow-up on the original POSEIDON trial, (POSEIDON-DCM trial NCT01392625) demonstrated that allogeneic MSC are safe and efficacious. In an early Phase I/IIb study, patients with chronic nonischemic dilated cardiomyopathy (NIDCM) were randomized to receive intramyocardial injections of either autologous or allogeneic MSCs.¹¹ Twelve-month SAE incidence was 28.2% following allogeneic MSC injection versus 63.5% with autologous MSC injection ($p=0.1$). The ejection fraction increased following allogeneic MSC injection by 8.0% ($p=0.004$) and 5.4% with autologous MSC injection ($p=0.116$; allo vs auto $p=0.4887$).¹¹ This improvement in LV function correlated with increased functional capacity as measured by the 6-minute walk test.¹¹ Therefore, allogeneic MSCs are safe and potentially effective in improving left ventricular ejection fraction. Duration of this affect still remains to be determined.

The development of new therapies for right ventricular dysfunction in patients with HLHS remains challenging. The ability to conduct laboratory investigation is limited because no available animal models replicate all of the physiologic derangements present, particularly with respect to a morphologic right ventricle providing the systemic cardiac output. Nevertheless, treatment of other models of ventricular failure with stem cell therapy has been promising, and this promise has been replicated in clinical trials for adult patients.

The hypothesis of our ELPIS Trial protocol is that MSCs transplanted into the myocardium of the right ventricle of patients with HLHS will stimulate favorable remodeling and thus improve myocardial function in the long term. We further hypothesize patients treated with MSC injection will have improved long-term survival and functional capacity compared to non-treated control patients, who currently have a average mortality rate of 54% at five years. The existing pre-

clinical and clinical data with MSC demonstrates that these cells have a strong regenerative ability. If the anticipated 10% to 15% improvement of right ventricular ejection fraction is seen in these HLHS patients, as was seen in the adult trial and preclinical animal data, we believe that this will lead to significantly improved survival and quality of life. Preserving, but more importantly, improving right ventricular function is critical for long-term survival of the HLHS patient.

Although trials of stem cell therapy for adults with cardiac disease have shown great promise, we believe children may be the best responders to stem cell therapy. In the ELPIS trial, we will be testing MSC injection in the right ventricular myocardium of pediatric patients with HLHS. As previous research has shown that neonatal and pediatric cardiomyocytes and cardiac progenitor cells are more responsive to the biological cues directed by transplanted stem cells. We also believe that these patients may be better candidates for functional improvements than adults.¹⁵ While this hypothesis needs further testing with clinical trials, indirect evidence is available from previous human studies examining the growing young human myocardium. Case reports using bone marrow-derived cell preparations and autologous cord blood have shown striking improvements in ventricular function in young HLHS patients. In a recent Japanese Phase I study of CDC therapy for HLHS, the best treatment responders were the smallest, youngest infants with the lowest ejection fractions.^{16,17} While encouraging, this study was limited by the lack of a proper control group and the heterogeneity of the patients enrolled. These observations are consistent with our data showing 3-fold higher c-kit+ CPC counts in neonates than in infants.¹⁸ Interestingly, levels of c-kit + CPCs in patients with heart failure are similar to those in normal neonatal hearts.²⁰ These findings may have important clinical implications for HLHS patients when the MSCs remodel the RV myocardium by stimulating human c-kit+ CPCs to proliferate and differentiate, as we have shown in swine myocardial infarction model.¹⁹ Although there is no large animal model of HLHS, in a neonatal porcine model of RV failure, we observed that MSC treatment facilitated remodeling of the RV myocardium at a histological and functional levels.¹³ Based on these preliminary insights, the young myocardium may be the best responder to MSC therapies to improve cardiac performance since the developing myocardium is more biologically responsive.

Conclusion

This landmark clinical trial will potentially result in a paradigm shift in HLHS treatment to meet an unmet medical need in this patient population. Our hypothesis that allogeneic MSCs can improve function in the right ventricle by both direct and indirect mechanisms builds on the advances in both surgery and cellular biology over the past thirty years. The ELPIS Trial is intended to address the remaining obstacles by using an allogeneic stem cell-based therapy to improve long-term ventricular function in patients with HLHS. We propose that a cell-based therapy for HLHS patients may prevent RV dysfunction, reducing both mortality and the need for transplantation.

Acknowledgement:

The cells are produced by Longeveron, LLC in Miami, and have been awarded TEDCO funding from the Maryland Stem Cell Research Fund and Children's Heart Foundation.

References

1. MMWR. Centers for Disease Control and Prevention, Hospital Stays, Hospital Charges, and In-Hospital Deaths Among Infants with Selected Birth Defects- United States, 2003, MMWR 2007; 56(5-29).
2. Bove EL. Current status of staged reconstruction for hypoplastic left heart syndrome. *Pediatr Cardiol* 1998;19:308-315.

3. Williams DL, Gelijns AC, Moskowitz AJ et al. Hypoplastic left heart syndrome: valuing the survival. *J Thorac Cardiovasc Surg* 2000;119:720-731.
4. Morrow WR, Naftel D, Chinnock R et al. Outcome of listing for heart transplantation in infants younger than six months: predictors of death and interval to transplantation. The Pediatric Heart Transplantation Study Group. *J Heart Lung Transplant* 1997;16:1255-1266.
5. Bernstein, D., Naftel, D., Chin, C. et al. Pediatric Heart Transplant Study. Outcome of listing for cardiac transplantation for failed Fontan: a multi-institutional study. *Circulation*. 2006; 114 (4): 273-280.
6. Gajarski RJ, Towbin JA, Garson A, Jr. Fontan palliation versus heart transplantation: a comparison of charges. *Am Heart J* 1996;131:1169-1174.
7. Tweddell JS, Hoffman GM, Mussatto KA et al. Improved survival of patients undergoing palliation of hypoplastic left heart syndrome: lessons learned from 115 consecutive patients. *Circulation* 2002;106:182-189.
8. Karantalis V, DiFede DL, Gerstenblith G et al. Autologous mesenchymal stem cells produce concordant improvements in regional function, tissue perfusion, and fibrotic burden when administered to patients undergoing coronary artery bypass grafting: The Prospective Randomized Study of Mesenchymal Stem Cell Therapy in Patients Undergoing Cardiac Surgery (PROMETHEUS) trial. *Circ Res*. 2014 Apr 11;114(8):1302-10.
9. Hare JM, Fishman JE, Gerstenblith G et al. Comparison of allogeneic vs autologous bone marrow-derived mesenchymal stem cells delivered by transendocardial injection in patients with ischemic cardiomyopathy: the POSEIDON randomized trial. *JAMA* 2012;308:2369-2379.
10. Ishigami S, Ohtsuki S, Eitoku T et al. Intracoronary Cardiac Progenitor Cells in Single Ventricle Physiology: The PERSEUS (Cardiac Progenitor Cell Infusion to Treat Univentricular Heart Disease) Randomized Phase 2 Trial *Circ Res* 2017 Mar 31;120(7): 1162-1173.
11. Hare JM, DiFede DL, Castellanos AM et al. Randomized Comparison of Allogeneic Vs. Autologous Mesenchymal Stem Cells for Non-Ischemic Dilated Cardiomyopathy: POSEIDON-DCM Trial. *Journal of the American College of Cardiology*. 2017;69(5):526-537.
12. Hare JM, Fishman JE, Gerstenblith G et al. Comparison of allogeneic vs autologous bone marrow derived mesenchymal stem cells delivered by transendocardial injection in patients with ischemic cardiomyopathy: the POSEIDON randomized trial. *JAMA* 2012;308:2369-2379.
13. Wehman B, Sharma S, Pietris N et al. Mesenchymal stem cells preserve neonatal right ventricular function in a porcine model of pressure overload. *Am J Physiol Heart Circ Physiol*. 2016 Jun 1;310(11):H1816-26.
14. Makkar R, Schatz R, Traverse J et al. Abstract 20536: allogeneic heart stem cells to achieve myocardial regeneration (allstar): the one year phase I results. *Circulation*. 2014;130:A20536.
15. Tarui, S., Sano, S. & Oh, H. Stem cell therapies in patients with single ventricle physiology. *Methodist DeBakey cardiovascular journal* 2014;10:77-81.
16. Ishigami, S. et al. Intracoronary autologous cardiac progenitor cell transfer in patients with hypoplastic left heart syndrome: the TICAP prospective phase 1 controlled trial. *Circulation research* 2015;116:653-664.
17. Tarui, S. et al. Transcoronary infusion of cardiac progenitor cells in hypoplastic left heart syndrome: Three-year follow-up of the Transcoronary Infusion of Cardiac Progenitor Cells in Patients With Single-Ventricle Physiology (TICAP) trial. *The Journal of thoracic and cardiovascular surgery* 2015;150:1198-1207, 1208 e1191-1192.
18. Mishra, R. et al. Characterization and functionality of cardiac progenitor cells in congenital heart patients. *Circulation* 2011;123:364-373.
19. Hatzistergos, K. E. et al. Bone marrow mesenchymal stem cells stimulate cardiac stem cell proliferation and differentiation. *Circulation research* 2010;107:913-922.
20. Wehman, B. et al. Pediatric End-Stage Failing Hearts Demonstrate Increased Cardiac Stem Cells. *The Annals of thoracic surgery* 2015;100:615-622.
21. Nelson TJ, Cantero Peral S. Stem Cell Therapy and Congenital Heart Disease. *Journal of Cardiovascular Development and Disease*. 2016; 3(3):24.
22. Sharma S, Mishra R, Bigham G et al. A Deep Proteome Analysis Identifies the Complete Secretome as the Functional Unit of Human Cardiac Progenitor Cells. *Circulation* 2017;120:816-834.

CCT

Corresponding Author



*Sunjay Kaushal MD, PhD
Division of Cardiac Surgery
University of Maryland Medical Center
110 S. Paca Street, 7th floor
Baltimore MD 21201
410-328-5842 (O); 410-328-2750 (F)
skaushal@som.umaryland.edu*

*Helina Mehta, MD
Division of Pediatric Cardiology,
University of Maryland School of
Medicine
110 S. Paca Street, 7th Fl.
Baltimore, MD USA*

*Kristopher Deatrck, MD
Division of Pediatric Cardiology,
University of Maryland School of
Medicine
110 S. Paca Street, 7th Fl.
Baltimore, MD USA*

*Michael Slack, MD
Division of Pediatric Cardiology,
University of Maryland School of
Medicine
110 S. Paca Street, 7th Fl.
Baltimore, MD USA*

*Joshua Hare, MD
Interdisciplinary Stem Cell Institute,
University of Miami Miller School of
Medicine
Miami, FL USA*



Archiving Working Group
**International Society for Nomenclature of
Paediatric and Congenital Heart Disease**
ipccc-awg.net

Tetralogy of Fallot with a Double Aortic Arch: A Case Report

By Gregory Aird, BA; Joel Hayden, BA; Randy Richardson, MD

Abstract

Tetralogy of Fallot (TOF) is a relatively common Congenital Heart Defect (CHD) known to correspond with 0.28 of every 1000 live births. Very rarely, TOF may be associated with other cardiac abnormalities, including Double Aortic Arch, a pathology that leads to the formation of a vascular ring that typically surrounds the trachea and esophagus that can potentially present in early childhood with symptoms of dysphagia and difficulty breathing. CT angiogram has been heavily relied upon as it also allows for 3D reconstruction and optimal visualization of the vascular ring pathology. This particular case presents an infant with TOF further complicated by double aortic arch that was reconstructed with 3D-CT angiogram to better visualize and understand the consequences of these congenital abnormalities on the cardiopulmonary system, as well as contribute to a fund of knowledge about the diagnosis and management of similar cases.

Introduction

Tetralogy of Fallot is a Cyanotic Congenital Heart Disease characterized by right ventricular hypertrophy, Ventricular Septal Defect (VSD), dextropositioning of the aorta to override the ventricular septum, and right ventricular outflow obstruction that ultimately leads to a right-to-left shunt and low oxygen saturation.¹ Although the tetralogy is characterized by a similar set of cardiac abnormalities, the condition can present with numerous variations to the pulmonary artery and aortic arch anatomies.^{1,2} Among the variations of the associated anatomy is double aortic arch. Double aortic arch is a vascular ring pathology in which the ascending aorta branches into two limbs that surround the trachea and esophagus before joining to form a single descending aorta.³ This

condition is the most common symptomatic aortic arch variant, often presenting with dysphagia or recurrent respiratory infections within the first six months after birth.³

Here, we discuss a rare case of Tetralogy of Fallot with double aortic arch diagnosed with the assistance of cardiac CT angiogram to create exceptionally detailed and educational three-dimensional images.

Case Report

A newborn female born to a 35-year-old G3P2 at 38 and 4/7 weeks via elective Cesarean section due to antenatally diagnosis of Tetralogy of

“Tetralogy of Fallot (TOF) is a relatively common Congenital Heart Defect (CHD) known to correspond with 0.28 of every 1000 live births. Very rarely, TOF may be associated with other cardiac abnormalities, including Double Aortic Arch, a pathology that leads to the formation of a vascular ring that typically surrounds the trachea and esophagus that can potentially present in early childhood with symptoms of dysphagia and difficulty breathing.”




Figure 1. 3D volumetric reconstruction of a double aortic arch with measurements.




Figure 2. 3D reconstruction posterior view of cardiac anatomy.

Fallot with multiple aortopulmonary collateral circulations is taken to the Neonatal Intensive Care Unit (NICU). The pregnancy was complicated by maternal supraventricular tachycardia treated with Labetolol, otherwise all immunizations were up-to-date and titers were normal. The patient's birth weight was 2900 grams, Apgars were 9 and 9 at 1 minute and 5 minutes, and vitals were found to be within normal limits with a temperature of 36.9 degrees Celsius, respiratory rate of 47 breaths per minute, blood pressure of 65/40 mmHg, pre-ductal saturations of 91%, post-ductal saturations of 100%, and SpO₂ at 96%.

On physical exam the patient was found to be stable with no signs of acute distress or symptoms of dyspnea and dysphagia.




Figure 3. 3D reconstruction anterior view of cardiac anatomy.




Figure 4. Posterior view of cardiac anatomy with removal of structures for optimal view of Double Aortic Arch.

Follow-up CT cardiogram with 3D reconstruction confirmed antenatal diagnosis of TOF with pulmonary atresia and revealed the presence of a double aortic arch with a dominant left arch and smaller, high-riding right arch that reconvene to form a single descending aorta (Figures 1-7). "In addition, the trachea is encircled, but not constricted by the double aortic arch (Figure 7). The typical findings of TOF including a large Ventricular Septal Defect (VSD) and ascending aorta supplied mainly by an enlarged right ventricle were noted. The trachea and esophagus did not appear to be compressed by the two aortic arches, and the right-sided vessel of the vascular ring was observed to fill a diminutive right main pulmonary artery in the absence of a main and left main pulmonary artery. Large, bilateral Major Aortopulmonary Collateral Arteries (MAPCA) were also seen to originate from the descending aorta (Figures 5 and 6).

Assessment of cardiac chamber size and function was performed by echocardiogram. While the left atria and left ventricle were normal in size, the echocardiogram revealed mild right atrial enlargement, as well as right ventricular hypertrophy with sustained systolic function. Blood flow measurements demonstrated a Patent Foramen Ovale (PFO) with right-to-left shunting, as well as a large Ventricular Septal Defect (VSD) with bidirectional shunting. Consistent with the findings of the CT cardiogram with 3D recon, the echocardiogram showed signs of a double aortic arch with the right-sided arch being smaller and more superior than the left-sided arch. Two MAPCAs were noted to arise off of the descending aorta.

Discussion

Tetralogy of Fallot is a Congenital Heart Defect that is present at birth as a right-to-left shunting disorder. Classically, there are four defects that characterize the anomaly: a Ventricular Septal Defect, pulmonary stenosis, right ventricular hypertrophy, and an overriding aorta. Symptoms noticeable at birth include cyanosis, heart murmur, and fatigue while breastfeeding. Although TOF is often seen as an isolated condition, it can rarely be associated with other abnormalities





Figure 5. Anterior view of cardiac anatomy with removal of structures for optimal view of Double Aortic Arch.

MAKING A DIFFERENCE



CP STENT™ UNMOUNTED OR PRE-MOUNTED ON A BIB® CATHETER
 FOR TREATMENT OF COARCTATION OF THE AORTA

NIT-OCCLUD® PDA COIL SYSTEM
 FOR TREATMENT OF PATENT DUCTUS ARTERIOSUS



INDICATIONS FOR USE:

The CP Stent™ is indicated for use in the treatment of native and/or recurrent coarctation of the aorta involving a compliant aortic isthmus or first segment of the descending aorta where there is adequate size and patency of at least one femoral artery and the balloon angioplasty is contraindicated or predicted to be ineffective. **WARNINGS / PRECAUTIONS:** Coarctation of the aorta involving the aortic isthmus or first segment of the descending aorta should be confirmed by diagnostic imaging. The CP stent has not been evaluated in patients weighing less than 20kg. As with any type of implant, infection secondary to contamination of the stent may lead to aortitis, or abscess. Over-stretching of the artery may result in rupture or aneurysm formation. Crimping the stent on a balloon catheter smaller than 12mm may cause damage to the stent. This device is intended for single use only. Do not sterilize and/or reuse it, as this can potentially result in compromised device performance and increased risk of cross-contamination. **CONTRAINDICATIONS:** Patients too small to allow safe delivery of the stent without compromise to the systemic artery used for delivery. Unfavorable aortic anatomy that does not dilate with high pressure balloon angioplasty. Curved vasculature. Occlusion or obstruction of systemic artery precluding delivery of the stent. Clinical or biological signs of infection. Active endocarditis. Known allergy to aspirin, other antiplatelet agents, or heparin. Pregnancy.

INDICATIONS FOR USE:

The Covered CP Stent™ is indicated for use in the treatment of native and/or recurrent coarctation of the aorta involving the aortic isthmus or first segment of the descending aorta where there is adequate size and patency of at least one femoral artery associated with one or more of the following: Acute or chronic wall injury; Nearly atretic descending aorta of 3 mm or less in diameter; A non-compliant stenotic aortic segment found on pre-stent balloon dilation; A genetic or congenital syndrome associated with aortic wall weakening or ascending aortic aneurysm. **WARNINGS / PRECAUTIONS:** Coarctation of the aorta involving the aortic isthmus or first segment of the descending aorta should be confirmed by diagnostic imaging. The CP stent has not been evaluated in patients weighing less than 20kg. As with any type of implant, infection secondary to contamination of the stent may lead to aortitis, or abscess. Over-stretching of the artery may result in rupture or aneurysm formation. Crimping the stent on a balloon catheter smaller than 12mm may cause damage to the stent. Excessive handling and manipulation of the covering while crimping the stent may cause the covering to tear off of the stent. This device is intended for single use only. Do not sterilize and/or reuse it, as this can potentially result in compromised device performance and increased risk of cross-contamination. **CONTRAINDICATIONS:** Patients too small to allow safe delivery of the stent without compromise to the systemic artery used for delivery. Unfavorable aortic anatomy that does not dilate with high pressure balloon angioplasty. Curved vasculature. Occlusion or obstruction of systemic artery precluding delivery of the stent. Clinical or biological signs of infection. Active endocarditis. Known allergy to aspirin, other antiplatelet agents, or heparin. Pregnancy.

INDICATIONS FOR USE:

The Nit-Occlud® PDA coil is a permanently implanted prosthesis indicated for percutaneous, transcatheter closure of small to moderate size patent ductus arteriosus with a minimum angiographic diameter less than 4mm. Nit-Occlud Brief Statement: Do not implant the Nit-Occlud PDA into patients who have endocarditis, endarteritis, active infection, pulmonary hypertension (calculated PVR greater than 5 Wood Units), thrombus in a blood vessel through which access to the PDA must be obtained, thrombus in the vicinity of the implantation site at the time of the implantation or patients with a body weight < 11 lbs (5 kg). An angiogram must be performed prior to implantation for measuring length and diameter of the PDA. Only the pfm medical implantation delivery catheter should be used to implant the device. Administration of 50 units of heparin per kg body weight should be injected after femoral sheaths are placed. Antibiotics should be given before (1 dose) and after implantation (2 doses) in order to prevent infection during the implant procedure. Do not implant the Nit-Occlud PDA in an MR environment. Do not pull the Nit-Occlud coil through heart valves or ventricular chambers. Contrast media should not be injected through the implantation catheter. The catheter must not be connected to high pressure injectors. Patients may have an allergic response to this device due to small amounts of nickel that has been shown to be released from the device in very small amounts. If the patient experiences allergic symptoms, such as difficulty in breathing or swelling of the face or throat, he/she should be instructed to seek medical assistance immediately. Antibiotic prophylaxis should be performed to prevent infective endocarditis during first 6 months after coil implantation. Potential Adverse Events: Air embolism, Allergic reaction to drug/contrast, Apnea, Arrhythmia requiring medical treatment or pacing, Arteriovenous fistula, Bacterial endocarditis, Blood loss requiring transfusion, Chest pain, Damage to the tricuspid or pulmonary valves, Death, Embolization of the occluder, requiring percutaneous or surgical intervention, Endarteritis, False aneurysm of the femoral artery, Fever, Headache/migraine, Heart failure, Hemolysis after implantation of the occluder, Hypertension, Hypotension or shock, Infection, Myocardial infarction, Occluder fracture or damage, Perforation of the heart or blood vessels, Stenosis of the left pulmonary artery or descending thoracic aorta, Stroke/TIA, Thromboembolism (cerebral or pulmonary), Valvular Regurgitation, Vessel damage at the site of groin puncture (loss of pulse hematoma, etc.).

Refer to the Instructions for Use for complete indications, relevant warnings, precautions, complications, and contraindications.

CP Stent is a trademark of NuMED, Inc. BIB is a registered trademark of NuMED, Inc. Nit-Occlud is a registered trademark of pfm medical, Inc.

Rx only CV9058 - 9/16 ©2016 B. Braun Interventional Systems Inc.



Distributed by:

B. Braun Interventional Systems Inc. | 824 Twelfth Avenue | Bethlehem, PA 18018 USA

Tel 877 836 2228 | Fax 610 849 1334 | www.bisusa.org

to the pulmonary and aortic vasculature. Among these variations include the rare condition known as Double Aortic Arch.

Double Aortic Arch (DAA) is an uncommon anomaly in which the ascending aorta branches into two vessels that form a vascular ring surrounding the trachea and esophagus before joining into a single descending aorta. Three subtypes of the vascular ring pathology are seen, with the most common being right-arch dominant and the least common being codominant. Although little is known about the exact etiology of the condition, it is often associated with the deletion of chromosome 22q11 which includes other Congenital Heart Defects, such as Tetralogy of Fallot.

The condition typically presents within the first month of the neonatal period and within the first 6 months of life. Although not always present, symptoms of double aortic arch arise due to compression of the esophagus and trachea by the vascular ring formed by the two vessels. As a result, the individual may have difficulty swallowing, inspiratory or expiratory stridor, and wheezing. Similarly, the presence of the defect predisposes the patient to recurrent respiratory infections, as well as a nagging, persistent cough.

While most incidences of DAA can be found prenatally via fetal ultrasound, confirmation of the diagnosis can be made using other imaging modalities such as echocardiography, magnetic resonance imaging, or computed tomography. The use of CT angiograms with contrast is the preferred diagnostic modality




Figure 6. Sagittal view of cardiac anatomy with removal of structures for optimal view of double aortic arch.

Pediatric Cardiac Intensivist

The Congenital Heart Collaborative (TCHC), an affiliation between University Hospitals Rainbow Babies & Children's Hospital (Cleveland OH) and Nationwide Children's Hospital (Columbus OH) heart programs, seeks candidates at the Assistant or Associate Professor rank for a faculty position in the Division of Pediatric Critical Care at UH Rainbow Babies & Children's Hospital in the role of Pediatric Cardiac Intensivist. Candidates should have dual training in Pediatric Critical Care and Cardiology, or Pediatric Critical Care with an additional year of Pediatric Cardiac Critical Care training. The successful candidate will be expected to perform at the highest level, with dedicated clinical service and night call in the Pediatric CTICU. The candidate will also have opportunities to participate in quality improvement initiatives, clinical research, and education of medical students, residents, and fellows.


The successful candidate will be well-supported at a world-class children's hospital with over 60 years of experience in the care of pediatric and adult congenital heart disease patients; an outstanding educational and research enterprise at Case Western Reserve University School of Medicine and an internationally recognized program partner with the Nationwide Children's Hospital Heart Center. TCHC is a dedicated service line with a common executive administration and functions as one program on two campuses with the commitment to expand access to high-quality comprehensive cardiac care regardless of patient age to the communities we serve while equally embracing a scholarly and educational mission. TCHC provide excellent cardiothoracic surgical, cardiac interventional, electrophysiologic, and non-invasive services. The candidate would be immediately accountable to the Pediatric Critical Care Division Chief and to TCHC senior leadership.

Please send letter of interest and curriculum vitae to:

Alexandre Rotta, MD, FCCM
Division Chief of Pediatric Critical Care
UH Rainbow Babies & Children's Hospital,
Alex.Rotta@uhhospitals.org

*Equal Opportunity Employer – minorities/females/
veterans/individuals with disabilities/sexual orientation/
gender identity.*

The Congenital Heart Collaborative | University Hospitals
Rainbow Babies & Children's
Nationwide Children's Hospital



7th Annual Scientific Sessions of the Cardiac Neurodevelopmental Outcome Collaborative

JUNE 6-8, 2018

In collaboration with  **Ward Family Heart Center**
CHILDREN'S MERCY KANSAS CITY






Figure 7. 3D reconstruction aerial view of double aortic arch.

owing to its visualization of the defect and its relationship with the structures nearby, such as the trachea and esophagus.

Treatment of these conditions is determined based on the severity and presence of symptoms. If left untreated, cases of TOF have a 35% mortality rate, and can present with complications such as growth retardation later on in life. As a result, complete surgical correction is often warranted to improve patient hemodynamics and outcome. Surgical correction of the DAA, which consists of ligation and dissection of the non-dominant vessel, is necessary only when symptoms of tracheal and esophageal compression are present.² Asymptomatic cases of DAA do not require surgery, and are carefully monitored until complications appear. Patients with TOF and DAA receive surgical correction of both defects regardless of the symptomatology of the double aortic arch.

Conclusion

While TOF is a relatively common Congenital Heart Defect, all suspected cases must be further explored with imaging modalities to monitor for the presence of other variations to

the pulmonary and aortic vasculature, such as Double Aortic Arch. While imaging techniques such as echocardiography can detect the presence of such vascular anomalies, CT cardiograms with 3D reconstruction can accurately diagnose the vascular ring pathology and show the relationship of the condition with the structures around it. Careful three-dimensional reconstruction of DAA in patients with TOF can also aid in the surgical correction of both defects, improving both patient care and outcome.

References

1. Apitz, C., Webb, G.D., Redington, A.N. Tetralogy of Fallot. *Lancet*. 2009; 374: 1462-1471.
2. Pankaj, B., Munesh, T., Bhan, A. Dysphagia in an adult Tetralogy of Fallot with double aortic arch. *Images in Pediatric Cardiology*. 2013; 15(3): 6-13.
3. Noguchi, K., Hori, D., Nomura, Y. et al. Double aortic arch in an adult. *Interact Cardiovascular Thoracic Surgery*. 2012; 14(6): 900-902.

CCT

Corresponding Author

Gregory Alan Aird is a third year medical student at Creighton University School of Medicine. He received his Bachelor of the Arts degree in Cognitive Science at Rice University where he graduated Magna Cum Laude. He is also a member of the Phi Beta Kappa Honors Society and Creighton's Radiology Interest Group.

Corresponding Author



Gregory Alan Aird, BA
3rd Year Medical Student
Creighton University School of Medicine
Phoenix Regional Campus
Phoenix, AZ USA

Mailing Address:
4140 N, Central Ave., Apt 1084
Phoenix, AZ 85012 USA
Phone: 949.243.4689
gaa66960@creighton.edu



Joel James Hayden, BA
4th Year Medical Student
Creighton University School of Medicine
Phoenix Regional Campus
Phoenix, AZ USA

Randy Richardson, MD
Creighton University School of
Medicine Phoenix Regional Campus
and Department of Radiology
St. Joseph's Hospital and Medical Center
Phoenix, AZ USA

Medical News, Products & Information

Compiled and Reviewed by Tony Carlson, Senior Editor

SCAI Launches Educational Initiative as Nation Shifts to Value-Based Healthcare — SCAI's Transradial Program Leads the Future of PCI Procedures

The Society for Cardiovascular Angiography and Interventions (SCAI) announced late October the launch of TRAnSition for VALUE, a multi-faceted educational initiative supported by Medtronic, to inform cardiologists, hospital administrators, and cath lab staff about the benefits of expanding the adoption of transradial (wrist access) percutaneous coronary intervention (PCI) in the U.S.

Approximately 70% of PCI procedures in the U.S. are performed through transfemoral (groin area) access. While this approach is often successful, it is associated with challenges, including an increased risk of bleeding complications at the insertion site and a slower recovery process for the patient.

As the country's healthcare system moves toward a value-based approach, TRAnSition for VALUE highlights the clinical and economic advantages of transradial intervention, which has increased from 16 to nearly 30% in the past decade. While not all patients can benefit from this approach, eligible patients may experience improved safety with reductions in major bleeding and vascular complications, increased comfort, a shorter hospital stay, and lower risk of mortality.

Despite the potential benefits, usage rates among U.S. interventionalists remain lower than those in similar healthcare systems, with cardiologists in major European and Asian countries reporting transradial usage at over 70%. Research indicates that a lack of education and training has impacted U.S. transradial adoption rates.

"Value for your healthcare dollar is an increasingly important driving force in the current healthcare environment. It is demanded by patients, payers and regulators," said Adhir Shroff, MD, MPH, FSCAI, chair of TRAnSition for VALUE, and director, Cardiovascular Catheterization Laboratories, University of Illinois. "The transradial approach represents an opportunity to provide value-based healthcare through improving outcomes for patients and streamlining care delivery within medical centers."

In addition, the transradial approach increases hospital efficiency by allowing same-day discharge, reducing recovery time and hospital staffing needs, and decreasing the cost of care. It also may expand treatment options for those who may not be candidates for transfemoral access PCI, such as those who are obese, elderly or have orthopedic disabilities that make it difficult to lie flat.

"Transradial PCI is truly a patient-centric approach with undisputed clinical benefits and patient preference," said Sunil V. Rao, MD, FSCAI, associate professor of Medicine, Duke University Medical Center, and co-chair of the initiative. "The issue has been a lack of training opportunities for physicians interested in adopting this

approach. SCAI's comprehensive TRAnSition for VALUE program offers the resources, training, and opportunity to learn from the nation's experts."

The TRAnSition for VALUE education curriculum will be guided by the SCAI Quality Improvement Committee and supports SCAI's consensus statement on transradial best practices. The primary education destination is a new website, TRAnSitionForVALUE.org, where visitors can register to receive custom education and training opportunities. The site will also host a modular, eight-part, quality improvement toolkit with accompanying webinars hosted by SCAI Quality Improvement experts on the following topics:

- How Value-Based Healthcare Affects Your Clinical Practice
- The Clinical Benefits of Transradial Access
- The Economic Benefits of Transradial Access
- Femoral to Radial 101 – Where to Begin for MDs
- Clinical Best Practices, Tips & Tricks for TRA Technique
- Rethinking Care Delivery in a TRA Cath Lab
- The Cath Lab Staff's Role in TRA Adoption
- Your TRA Support Network: How to Initiate Change

The TRAnSition for VALUE program will also be featured at the annual SCAI Scientific Sessions, as well as other regional events in partnership with hospitals and healthcare systems throughout the U.S.

The Society for Cardiovascular Angiography and Interventions is a 4,300-member professional organization representing invasive and interventional cardiologists in more than 75 countries. SCAI's mission is to promote excellence in invasive/interventional cardiovascular medicine through physician and allied health professional education and representation, and advancement of quality standards to enhance patient care. For more information about SCAI, visit www.SCAI.org.

Re-Interventions Are Common in Long-Term Survivors of Childhood Heart Operations - CHOP-Led Study: Surgical or Catheter Procedures Needed in Majority of Long-Term Survivors of Fontan Procedure

Newswise — Among patients who undergo childhood heart surgery for the severe birth defect Single-Ventricle Disease, two-thirds of survivors require a surgical or catheter-based procedure within 20 years. Pediatric cardiology researchers note that doctors should counsel families about the likelihood of re-interventions.

"Unfortunately, for many patients, the Fontan is not the final intervention," said study leader Andrew Glatz, MD, MSCE, referring to the Fontan operation, the third in a series of reconstructive operations performed on children with a severely underdeveloped ventricle, one of the heart's two pumping chambers. Glatz is a pediatric interventional cardiologist in the Cardiac Center at Children's Hospital of Philadelphia (CHOP).

Glatz and colleagues published their study on September 1, 2017, in *Circulation: Cardiovascular Interventions*. Other key members of

www.picsymposium.com

the study team include Tacy Downing, MD and Kiona Allen, MD (both were pediatric cardiology fellows at CHOP during the work); and David Goldberg, MD and William Gaynor, MD (current faculty members in the Cardiac Center at CHOP).

The study team performed a retrospective review of 773 patients who underwent the Fontan operation at CHOP between 1992 and 2009.

Although the Fontan procedure offers high survival rates for a condition that previously was universally fatal during infancy, it cannot provide normal blood circulation, and carries long-term risks of complications that continue to be analyzed. Clinicians and researchers were aware of the need for re-interventions in long-term Fontan survivors, but there was little detailed knowledge of re-intervention rates until now.

In the current study, the researchers found that 65% of Fontan survivors underwent a re-intervention by 20 years after their operation, with a median time to first re-intervention slightly less than 10 years. The re-interventions were either operations or catheterizations, with catheterizations being more common—often to close unwanted openings or to widen narrowed blood vessels. Among operations, the most common were to place or revise a pacemaker.

“The important message from this work is that, for many patients, the Fontan operation is not the ‘final’ procedure, as it is sometimes referred to. Instead, many patients require further interventions after the Fontan to continue to try to optimize the circulation, as best as possible. It’s important for families and doctors to understand this, so expectations are clear. This also highlights the need for close and careful ongoing follow-up after the Fontan operation by pediatric cardiologists familiar with potential complications that could befall a Fontan patient,” said Glatz.

Tacy Downing, MD is now an interventional and adult congenital cardiologist at Nicklaus Children’s Hospital and Kiona Allen MD, a pediatric cardiac intensivist at Ann & Robert H. Lurie Children’s Hospital.

Tacy E. Downing, et al “Surgical and Catheter-based Re-Interventions Are Common in Long-Term Survivors of the Fontan Operation” *Circulation: Cardiovascular Interventions*, September 1, 2017. doi: 10.1161/CIRCINTERVENTIONS.116.004924.

Children’s Hospital of Philadelphia: Children’s Hospital of Philadelphia was founded in 1855 as the nation’s first pediatric hospital. Through its long-standing commitment to providing exceptional patient care, training new generations of pediatric healthcare professionals, and pioneering major research initiatives. Its pediatric research program is among the largest in the country. In addition, its unique family-centered care and public service programs have brought the 546-bed hospital recognition as a leading advocate for children and adolescents. For more information, visit www.chop.edu.

Families of Survivors of Extracorporeal Membrane Oxygenation (ECMO) for Heart Conditions Report Favorable Quality of Life

Newswise — One of the few large studies to report long-term outcomes in cardiac patients treated in childhood with extracorporeal membrane oxygenation (ECMO) has found overall favorable outcomes among survivors, as reported by families. ECMO provides short-term breathing



Chief for its Division of Cardiology and Co-Executive Director of The Heart Institute

Children’s Hospital Los Angeles (CHLA), in partnership with University of Southern California (USC) and the Keck School of Medicine of USC (KSOM), seeks to recruit a Chief for its Division of Cardiology and Co-Executive Director of The Heart Institute. We seek a visionary leader to further develop a successful clinical, research and academic program that serves a large and diverse patient population in our world-class children’s hospital. The Division Chief provides leadership for the Division, overseeing clinical care delivery and quality, budget and financial performance, strategic planning, recruitment and program development, and ensuring the integration of training and research programs with clinical services throughout the hospital. The Chief will also lead the Division’s support of the departmental academic goals in research, education and faculty recruitment and development and join the Department of Pediatrics Leadership team.

The successful candidate will have strong academic, clinical, and leadership/management experience in either a research-oriented children’s hospital or an academic pediatric department affiliated with a university medical school, with a minimum of 10 years of progressive medical and research experience and an increasing scope of responsibility including faculty and trainee mentoring and program development. The candidate will have a track record of scholarly accomplishment, ideally with extramural funding.

The candidate must have an MD, DO or MD/PhD degree, must have completed at least three years of post-doctoral or clinical fellowship training, and must be board-certified by the American Board of Pediatrics and board-certified in Pediatric Cardiology. The candidate should have clinical as well as research experience in a pediatric setting and qualify for appointment as Associate Professor or Full Professor at KSOM.

To apply, please direct inquiries, nominations, and applications, including resume and a letter of interest, in confidence to:

Children’s Hospital Los Angeles
Attn: Valerie Campana, Director of Faculty Affairs and Development
4650 Sunset Blvd., M/S #71
Los Angeles, California 90027
Vcampana@chla.usc.edu

Electronic submission is encouraged.

CHLA and USC are EO/AA Employers



The congenital heart professionals network exists to facilitate communications between congenital heart professionals locally, regionally, and globally.

JOIN TODAY

www.chipnetwork.org



Funded by Cincinnati Children’s Heart Institute

and heart support for critically ill children while doctors treat the underlying illness.

A research team from Children's Hospital of Philadelphia (CHOP) published the ECMO study in the August 2017 issue of *Pediatric Critical Care Medicine*.

The team analyzed a cohort of 396 patients with cardiac disease treated with ECMO at CHOP from 1995 to 2012. Overall mortality was 66a5 at a median follow-up of 6 years after ECMO therapy, which remains consistent with outcomes seen in previous decades.

In phone surveys or written surveys among the families of survivors, a majority reported positive outcomes regarding health and physical limitations. Over 90% of families reported good or excellent health, and approximately 86% reported no or mild physical limitations.

However, the authors noted a discrepancy between family-reported favorable outcomes and a relatively high rate of medical and behavioral issues revealed by more detailed questioning. Almost 25% of patients had below-average school performance and required special education, and almost 50% had parental-reported learning disabilities. These results may help families define realistic expectations regarding long-term outcomes for children supported with ECMO due to an underlying cardiac condition.

Matthew D. Elias, MD, a pediatric cardiologist at CHOP and first author of the study, noted that ECMO use in children with Congenital Heart Disease (CHD) has increased markedly over the past several decades, as increased experience in pediatric cardiology and cardiac surgery has allowed ECMO use to expand to more complex patients. Senior author Matthew J. O'Connor, MD, also a CHOP pediatric cardiologist, added that "several factors have potentially improved long-term outcomes, such as increasing experience with ECMO and CHD in general. But the inclusion of a more medically complex population in the recent era may mitigate these improvements in outcomes, accounting for the fact that overall mortality rates haven't changed much."

Although this single-center study represents one of the largest cohorts of ECMO patients undergoing detailed assessments of outcomes and quality of life, Elias said that further research in larger, multicenter studies should further investigate family experiences and

PHYSICIAN OPPORTUNITY

Peoria, Illinois

Seeking Assistant Professor, Pediatric Cardiology

Job Specifics

- The new pediatric cardiologists will join a well-established team of ten pediatric cardiologists with 30 plus years of success in the region.
- The new pediatric cardiologists will join the group who staffs Children's Hospital of Illinois (CHOI) in Peoria and supports outreach clinics located in surrounding community cities.
- Professional efforts will be bolstered by the support of two pediatric cardiovascular surgeons.
- Preferred candidates should have training in pediatric interventional cardiac catheterization
- Clinical activities will also include in-patient rotations, clinic coverage including satellite clinics
- Scholarly and research interests are highly desirable, along with experience in teaching medical students and residents.
- The candidate must be board-certified or board-eligible in pediatric cardiology and will report to the division head of pediatric cardiology.
- EEO Employer M/F/Vet/Disabled



The Community

- Peoria, Illinois has a greater metro area population of over 350,000, offers a remarkably low cost of living with the big city comforts and attractions.
- Quality public and private pre-school to University educational options, the University of Illinois College of Medicine, Bradley University and a junior college nearby, educational options are abundant.
- Peoria offers a vibrant downtown with a beautiful skyline, river front, cultural events and entertainment.

Children's Hospital of Illinois is the primary pediatric teaching facility for the University Of Illinois College Of Medicine, a 136-bed facility that offers over 50 pediatric programs and services, including one of the largest pediatric surgery programs in Illinois. It is supported by a growing referral base of 140 CHOI Pediatric Sub-Specialists, 72 faculty members at the University of Illinois College of Medicine-Peoria, over 68 OSF Medical Group practice locations in Illinois and over 800 physicians on staff at OSF Saint Francis Medical Center. Children's Hospital of Illinois is the only full service tertiary hospital for children and adults with congenital heart disease in downstate Illinois.

Please contact or send CV to:

Stacey Morin,
OSF HealthCare Physician Recruitment
Ph: (309) 683-8354
Email: stacey.e.morin@osfhealthcare.org
Web: www.osfhealthcare.org

 OSF HEALTHCARE

 OSF HealthCare
Children's Hospital
of Illinois

 UNIVERSITY OF ILLINOIS
COLLEGE OF MEDICINE AT PEORIA

**CONGENITAL
CARDIOLOGY
TODAY**

Looking for Career Opportunities in Congenital/Structural Heart Disease (CHD)?

Go to: www.CongenitalCardiologyToday.com/Recruit/Recruit.html

long-term patient outcomes. He added, "In the meantime, our findings should allow for improved family counseling in discussing long-term quality-of-life for children with heart disease."

The ECMO Center at Children's Hospital of Philadelphia recently received the "Platinum Level ELSO Award for Excellence in Life Support" from the Extracorporeal Life Support Organization (ELSO), an international consortium of centers offering ECMO (extracorporeal membrane oxygenation) for support of failing organ systems in infants, children and adults. CHOP's ECMO Center has been recognized as an ELSO Center of Excellence since 2008. The Platinum Level is the highest awarded honor, and is rarely achieved by ELSO member institutions, especially pediatric centers. The ECMO Center at CHOP is the only ECMO center in the Philadelphia region designated by ELSO as a Platinum Center of Excellence, and is one of the most active in the country. It has supported more than 1,300 patients since it was established in 1990. The program's multidisciplinary team is comprised of pediatric surgeons, neonatologists, intensivists, anesthesiologists, perfusionists, specially trained nurses and respiratory therapists.

Matthew D. Elias, et al. "Long-Term Outcomes of Pediatric Cardiac Patients Supported by Extracorporeal Membrane Oxygenation," *Pediatric Critical Care Medicine*, August 2017.

<http://doi.org/10.1097/PCC.0000000000001227>.

For more information visit www.chop.edu.

CHOP Pediatric Specialty Consultation Services Will Provide Education, Clinical Care in the UAE

Newswise — On Oct. 5th, The Ministry of Health and Prevention of the United Arab Emirates (MOHAP) and Children's Hospital of Philadelphia (CHOP) entered into a memorandum of understanding regarding a pediatric specialty consultation program to provide clinical and educational services to MOHAP hospitals.

"The Ministry of Health and Prevention of the United Arab Emirates is dedicated to providing quality care for our patients and

education for our clinicians," said Yousif Mohammad Al Serkal, Assistant Undersecretary for Hospitals Sector of MOHAP. "CHOP is a global leader in pediatric medicine, with top-rated clinical programs, groundbreaking research, educational and training programs, and state-of-the-art facilities."

"CHOP has an investment in the healthcare of children in the Middle East region and is excited to offer The Ministry of Health of the United Arab Emirates our pediatric experts on a short-term visiting basis to provide education and clinical care in the MOHAP hospitals," said Matthew Bayley, Senior Vice President and Chief Strategy Officer at Children's Hospital of Philadelphia. "As a result of this important initiative, both clinicians and children in the Middle East region will benefit from medical education and consultations by CHOP healthcare providers who are experts in the field of pediatric medicine."

The Visiting Consultants Program intends to bring the expertise of CHOP physicians to patients at MOHAP hospitals within the United Arab Emirates. The visiting physicians will meet with the clinical team, perform clinical consultations as needed, perform inpatient consultations and give lectures to the MOHAP team.

The visits are advertised through mainstream media and through the MOHAP Hospitals to ensure proper outreach.

For more information visit www.chop.edu

The Ministry of Health and Prevention is a ministry of the Government of United Arab Emirates, and is responsible for the implementation of health care policy in all areas of technical, material, and coordination with the Ministries of State, and cooperation with the private sector in health locally and internationally.

Quality Improvement Through the ACPC Quality Network

By Devyani Chowdhury, MBBS, FACC

ACC's Adult Congenital and Pediatric Cardiology (ACPC) Quality Network is a national network of Congenital Heart



Boston Children's Hospital
Until every child is well

Faculty Position Available Pediatric Cardiology Inpatient Service

The Inpatient Division of the Department of Cardiology at Boston Children's Hospital, a teaching hospital of Harvard Medical School, is recruiting new faculty at the assistant professor level. The Pediatric Cardiology Inpatient Service, among the largest in North America, comprises patients with congenital and acquired pediatric heart disease. Care teams include residents, fellows, and advance practice nurses supervised by faculty cardiology attending physicians.

Candidates must be board certified in pediatrics and pediatric cardiology, have at least five years of relevant experience beyond fellowship, and have excellent clinical, teaching, and academic skills. Preference will be given to individuals with strong interest and experience in medical student and resident education and high potential for leadership in this area.

Interested individuals should send inquiries and curriculum vitae to:

Roger E. Breitbart, MD
Chair, Inpatient Cardiology Search Committee

**Department of Cardiology
Boston Children's Hospital
300 Longwood Avenue
Boston, Massachusetts 02115**

Email:

roger.breitbart@cardio.chboston.org

Boston Children's Hospital is an Equal Opportunity Employer. All qualified applicants will receive consideration for employment without regard to race, color, religion, sex, sexual orientation, gender identity, national origin, disability status, protected veteran status, or any other characteristic protected by law.



**HARVARD MEDICAL SCHOOL
TEACHING HOSPITAL**

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share? Submit your manuscript to: RichardK@CCT.bz

Disease (CHD) Centers dedicated to collecting and sharing data, collaborating and developing best practices for CHD patients. The network offers data aggregation, benchmarking to improve patient care, and collaborative quality improvement. There are currently 36 sites participating in the ACPC Quality Network, representing academic and private practice with a total of five quarters of data collection.

The Quality Network was developed to operationalize a subset of ACPC quality metrics allowing participating centers to compare their performance anonymously across a variety of quality metrics. Participating sites select from available metrics and submit data on a quarterly schedule.

To augment the data collection and reporting activities, the College's ACPC Member Section sponsors a series of activities to facilitate discussion between participating Quality Network centers. It is expected that these discussions will lead to increased sharing of current practices and ultimately improve practices in specific areas across the CHD field. Activities promoting collaborative quality improvement include in-person meetings at ACC's *Annual Scientific Session*, webinars or in-person learning sessions, as well as a dedicated listserv.

Two of the Quality Network metrics – counseling for patients with body mass index >85% and assessment of 22q11 status in Tetralogy of Fallot patients – have been approved by the American Board of Pediatrics (ABP) as a Quality Improvement Activity. Physicians at participating Quality Network Centers meeting meaningful participation requirements outlined by the ACC and the ABP can earn Part IV Maintenance of Certification by collecting and submitting data for these two metrics.

Enrollment in the ACPC Quality Network is available to pediatric cardiology and adult congenital heart disease practices who have at least one fellow of the ACC within the practice.

For more information about the ACPC Quality Network, including current metrics and details about earning MOC Part IV

credit, visit ACC.org/QNet or email ACPCQNet@acc.org.

Cardiac MR Added to Digisonics Congenital Cardiology Solution with Medis Plug-in

Digisonics and Medis have partnered to provide a comprehensive single system solution that adds Cardiac MR to Digisonics Congenital Echo and Cath system.

The industry leading QMass and QFlow applications from Medis will help solidify Digisonics as the only congenital cardiology specializing in structured reporting and image management for Echo, Cath and Cardiac MR.

Digisonics provides top-rated clinical image management and structured reporting systems for cardiovascular (CVIS), radiology, and obstetrics & gynecology. Digisonics structured reporting solutions combine high performance image review workstations, a powerful PACS image archive, an integrated clinical database, comprehensive analysis capabilities and highly configurable reporting for multiple modalities. Key applications are complemented with interfaces to information systems and 3rd party vendors, providing facilities with a seamless, efficient clinical workflow. To learn more, please visit www.digisonics.com.

Cardiology Societies to Expand Maintenance of Certification Offerings

The American College of Cardiology (ACC), Heart Failure Society of America (HFSA), Heart Rhythm Society (HRS) and Society for Cardiovascular Angiography and Interventions (SCAI) have announced a partnership to develop new modules to help subspecialty cardiologists potentially meet collaborative maintenance pathway requirements equivalent to satisfying the ABIM's current 10-year MOC examination.

"It is a shared goal of ACC, HFSA, HRS and SCAI to help our collective members ensure their patients are receiving the highest quality, evidence-based care," said ACC President Mary Norine Walsh, MD, FACC.



Congenital Cardiac Intensivist

The Heart Center (THC) at Nationwide Children's Hospital, the primary pediatric teaching facility for The Ohio State University in Columbus Ohio, is recruiting an attending physician, at any academic level, for the Cardiothoracic Intensive Care Unit (CTICU). This individual would join a group of seven multi-background academic cardiac intensivists and eight dedicated nurse practitioners devoted to the CTICU providing 24/7 in house coverage.

Our current independently-managed free-standing CTICU is a 20 bed unit with approximately 800 total admissions per year (medical and surgical) and an average daily census of 12. Candidates must have completed fellowship training in pediatric cardiac anesthesia, critical care and/or cardiology that includes advanced cardiac intensive care training. Preference will be given to those with experience and are boarded in pediatric cardiology.

THC embraces a culture of patient safety and quality, transparency, translational/outcome research, education, cost-containment and public health awareness. These create ample participation and leadership opportunities for the candidate's professional growth. THC is comprehensive with services including an active hybrid palliation center, a comprehensive single ventricle program, thoracic organ transplantation program, blood conservation strategies, and cardiac mechanical support team. The CTICU is supported by world-class and innovative interventional cardiology, cardiac imaging, cardiothoracic surgery and adult congenital heart experts. Current annual clinical metrics for THC includes: over 500 cardiothoracic surgeries, over 700 cardiac catheterizations and EP procedures, and over 13,000 cardiology outpatient visits. We have a pediatric and pediatric/adult combined cardiology fellowship programs. We participate in numerous multicenter clinical trials and quality initiatives including the JCCHD QI Collaborative. We are directly linked to our Center for Cardiovascular and Pulmonary Research which has an NIH T-32 training grant.

Interested candidates are encouraged to submit their curriculum vitae to:

Janet Simsic, MD
Director of the Cardiothoracic Intensive Care Unit,
Nationwide Children's Hospital, T2296
700 Children's Drive
Columbus, OH 43205
or janet.simsic@nationwidechildrens.org

KINGSTON
FREELANCE CONSULTING
MEDTECH CONSULTING FROM CONCEPT TO COMMERCIALIZATION

We can help you navigate and manage the challenges and risks of achieving market success from concept to commercialization.

AMBER KINGSTON
2305 WOODMONT CIRCLE, STE. K
MACUNGIE, PA 18062
TEL: 610.463.4964
AKKINGSTON@ICLOUD.COM
[LINKEDIN.COM/IN/AMBERKINGSTON](https://www.linkedin.com/in/amberkingston)

“In offering additional pathways for cardiologists who wish to maintain their professional certification, we can more effectively and efficiently help busy clinicians keep up with current knowledge in their specific areas of practice.”

Through this partnership, which will move forward upon reaching agreement with the ABIM, the ACC will enhance its existing ACC Self-Assessment Program (ACCSAP) product line with CathSAP, EPSAP and Heart Failure SAP products to help fulfill the MOC needs of interventionalists, electrophysiologists and heart failure specialists. An analogous product for ABIM diplomates in Adult Congenital Heart Disease will be developed by 2021.

“This program and partnership will deliver great value for all of our members. The sophistication of these self-assessment modules will allow a more streamlined pathway for maintaining credentials,” said HFSA President Christopher O’Connor, MD, FHFA.

“The Heart Rhythm Society is excited about exploring a partnership with ACC and the other societies in this initiative,” said HRS President George Van Hare, MD, FHRS, FACC. “We have long advocated for a process of recertification that recognizes the advantages of continuous learning, as opposed to a single high-stakes exam. HRS looks forward to having an opportunity to collaborate on developing a new MOC process.”

“SCAI leadership is eager to partner with ACC, to share in the work of building out best-in-class re-certification materials. By leveraging the breadth and depth of talent within both organizations, and maintaining close communication with ABIM, we can facilitate a quicker, easier, and more relevant MOC experience for our members,” said SCAI President Kirk N. Garratt, MD, MSc, FSCAI.

Development is in the early stages, but the new products may launch as early as 2019. During development the current ACCSAP 9 remains in place to help clinicians maintain professional competence.

The American College of Cardiology is the professional home for the entire cardiovascular care team. The mission of the College is to transform cardiovascular care

and to improve heart health. The ACC leads in the formation of health policy, standards and guidelines. The College operates national registries to measure and improve care, offers cardiovascular accreditation to hospitals and institutions, provides professional medical education, disseminates cardiovascular research and bestows credentials upon cardiovascular specialists who meet stringent qualifications.

The Heart Failure Society of America, Inc. represents the first organized effort by heart failure experts from the Americas to provide a forum for all those interested in heart function, heart failure, and congestive heart failure (CHF) research and patient care. For more information, visit www.hfsa.org.

The Heart Rhythm Society is the international leader in science, education and advocacy for cardiac arrhythmia professionals and patients, and the primary information resource on heart rhythm disorders. Its mission is to improve the care of patients by promoting research, education and optimal health care policies and standards. For more information, visit www.HRSonline.org.

The Society for Cardiovascular Angiography and Interventions is a 4,300-member professional organization representing invasive and interventional cardiologists in approximately 70 countries. SCAI’s mission is to promote excellence in invasive/interventional cardiovascular medicine through physician and allied health professional education and representation, and advancement of quality standards to enhance patient care. For more information about SCAI, visit www.SCAI.org. Follow @SCAI on Twitter for the latest heart health news.

CONGENITAL CARDIOLOGY TODAY

CALL FOR CASES AND OTHER ORIGINAL ARTICLES

Do you have interesting research results, observations, human interest stories, reports of meetings, etc. to share? Submit your manuscript to: RichardK@CCT.bz

CONGENITAL CARDIOLOGY TODAY

© 2017 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499-online). *Published monthly. All rights reserved.*

www.CongenitalCardiologyToday.com

Publication Company Address:

11502 Elk Horn Dr. Ste. 201
Clarksburg, MD 20871 USA
Tel: +1.301.279.2005

Publishing Management:

- Tony Carlson, Founder, President & Sr. Editor - TCarlsonmd@gmail.com
- Richard Koulbanis, Group Publisher & Editor-in-Chief - RichardK@CCT.bz
- John W. Moore, MD, MPH, Group Medical Editor - JMoore@RCHSD.org

Editorial Board:

Teiji Akagi, MD; Zohair Al Halees, MD; Mazeni Alwi, MD; Felix Berger, MD; Fadi Bitar, MD; Jacek Bialkowski, MD; Mario Carminati, MD; Anthony C. Chang, MD, MBA; John P. Cheatham, MD; Bharat Dalvi, MD, MBBS, DM; Horacio Faella, MD; Yun-Ching Fu, MD; Felipe Heusser, MD; Ziyad M. Hijazi, MD, MPH; Ralf Holzer, MD; Marshall Jacobs, MD; R. Krishna Kumar, MD, DM, MBBS; John Lamberti, MD; Gerald Ross Marx, MD; Tarek S. Momenah, MBBS, DCH; Toshio Nakanishi, MD, PhD; Carlos A. C. Pedra, MD; Daniel Penny, MD, PhD; James C. Perry, MD; P. Syamasundar Rao, MD; Shakeel A. Qureshi, MD; Andrew Redington, MD; Carlos E. Ruiz, MD, PhD; Girish S. Shirali, MD; Horst Sievert, MD; Hideshi Tomita, MD; Gil Wernovsky, MD; Zhuoming Xu, MD, PhD; William C. L. Yip, MD; Carlos Zabal, MD

Free Subscription to Qualified

Professionals: Send your name, title(s), hospital or practice name, work address and url, phone, fax and email to: sub@cct.bz.

Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors, and are not necessarily the views of Congenital Cardiology Today.



2nd International Training Workshops

aEEG and NIRS

www.munich-neocon.com

March 23rd – 25th, 2018

Congenital Cardiac Care Providers in North America at Hospitals That Offer Open Heart Surgery for Children

DIRECTORY 2016

ver. 12/1/16

Published by

CONGENITAL CARDIOLOGY TODAY

www.CongenitalCardiologyToday.com

ATTENTION:

**Division Chiefs of Pediatric Cardiology
and Fellowship Directors**

The Directory is Being Updated

Over the next few weeks, we will be sending emails to you with your hospital's information as listed in the 2015 directory.

View your current listing at:

<http://goo.gl/yi3Tq6>

If you or your hospital are not listed,
please send an email to: DIRECTORY@CCT.BZ