# CONGENITAL CARDIOLOGY TODAY

Timely News and Information for BC/BE Congenital/Structural Cardiologists and Surgeons

Volume 5 / Issue 12 December 2007 North American Edition

### IN THIS ISSUE

Role of Interventional Cardiology in Neonates – Part I - Non-Surgical Atrial Septostomy by P. Syamasundar Rao, MD Page 1

Interview with Richard Van Praagh- Part 2: Family Life by Bradley W. Robinson, MD Page 13

### **DEPARTMENTS**

Medical News, Products and Information Page 21

#### CONGENITAL CARDIOLOGY TODAY

Editorial and Subscription Offices 16 Cove Rd, Ste. 200 Westerly, RI 02891 USA

#### www.CongenitalCardiologyToday.com

© 2007 by Congenital Cardiology Today ISSN: 1544-7787 (print); 1544-0499 (online). Published monthly. All rights reserved. Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors, and are not necessarily the views of Congenital Cardiology Today.

> Recruitment Ads on Pages: 2, 5, 6, 7, 8, 10, 11, 12, 14, 15, 16, 18, 20, 22 and 23

### Role of Interventional Cardiology in Neonates – Part I - Non-Surgical Atrial Septostomy

By P. Syamasundar Rao, MD

"Role of Interventional Cardiology in Neonates – Part I" is the first in a series of three articles by P. Syamasundar Rao, MD, Professor of Pediatrics and Medicine; Director, Division of Pediatric Cardiology; University of Texas-Houston Medical School. The second and third articles will appear in upcoming issues.

### INTRODUCTION

Whereas the seeds of interventional pediatric cardiology were planted in the 1950s by Rubio-Alvarez and Limon-Lason by their work on transcatheter pulmonary and tricuspid valvotomy [1,2], by Dotter and Jedkins in the early 1960s by their gradational vascular dilatation techniques [3], by Rashkind in the mid-1960s by introduction of balloon atrial septostomy[4], by Porstmann in late 1960s by percutaneous closure of patent ductus arteriosus[5,6], and by King and Mills mid-1970s by non-operative closure of in atrial septal defects [7,8], it is not until Kan and her associates [9] adapted Gruntzig's technique [10] of balloon angioplasty with double-lumen balloon catheter for pediatric use in early the 1980s did transcatheter therapy in children become a reality [11-15]. These techniques were extended slowly but surely to treat neonates with congenital heart defects [16-20]. In this review, various catheter interventional techniques (Table I)

that are currently being used in neonates will be discussed; in Part I, Non-surgical atrial septostomy will be reviewed, and the remaining items listed in Table I will be dealt with in subsequent issues of this publication.

### Table I. Catheter InterventionalTechniques Used in the Neonate

- Non-surgical atrial septostomy
- Balloon angioplasty/valvuloplasty
- Radiofrequency perforation of atretic pulmonary valve
- Transcatheter occlusion of shunts
- Stents

In 1966, Rashkind and Miller [4] described a technique, now called Rashkind balloon atrial septostomy, which was extensively used to improve atrial mixing in neonates with transposition of the great arteries. It was subsequently applied to many other disease entities in which enlarging the atrial defect is beneficial [21]. In mid to late 1970s, Park and his associates extended the utility of balloon septostomy by introducing blade atrial septostomy to enlarge defects with

### Table II. Septostomy Procedures

- Rashkind balloon atrial septostomy
- Blade atrial septostomy
- · Balloon angioplasty
- Atrial septal perforation
- Stent implantation

Do you or your colleagues have interesting research results, observations, human interest stories, reports of meetings, etc. that you would like to share with the congenital cardiology community?

> If so, submit a brief summary of your proposed article to Congenital Cardiology Today at: Article@CCT.bz The final manuscript may be between 400-3,500 words, contain pictures, graphs, charts and tables.



### Vanderbilt Children's Hospital Division of Pediatric Cardiology is currently recruiting two BE/ BC cardiologists with expertise in echocardiography, including fetal and transesophageal studies.

The addition of these two candidates would bring the total complement of echocardiographers to five. Participation in the current cardiac MRI program is an option, but not required. In addition to clinical duties, the candidate would provide training at the fellowship and residency levels and would be expected to participate in the active research program of this division. This is a fully digital laboratory which functions within the framework of a comprehensive cardiac center, located in the state-of the art, 222 bed Vanderbilt Children's Hospital. The rapidly growing cardiology division includes 14 cohesive members who cover all aspects of cardiac care. There are two excellent cardiac surgeons who perform more than 500 operations per year, of all types, with impressive results.

The Children's Hospital is located on the beautiful Nashville campus of Vanderbilt University, nestled in the rolling hills of middle Tennessee. Nashville is a very safe, mid sized metropolitan area with top notch public and private schools, inviting parks and lakes, professional hockey and football, and the best music anywhere. The members of the cardiology division, in addition to valuing clinical and academic expertise, also place a high priority on family time and development of interests outside of the medical realm.



If you'd like to know more, please contact Ann Kavanaugh-McHugh, M.D., echocardiography lab director, for further information. Her phone number is 615-322-7447, and email is ann.kavanaugh-mchugh@vanderbilt.edu thick atrial septae [22]. A built-in retractable blade (knife) cuts the lower margin of the patent foramen ovale (PFO) which is followed by balloon atrial septostomy. More recently, balloon angioplasty [21,23,24], stents [25,26], Ross transseptal puncture [27], radiofrequency ablation [28-30] and cutting balloons [30] were applied to create and/or enlarge the atrial defects.

Cardiac defects in which atrial septostomy is likely to be useful will be reviewed followed by a description of septostomy procedures.

#### SEPTOSTOMY PROCEDURES

Various septostomy procedures that are currently used will be discussed hereunder.

#### TRANSPOSITION OF THE GREAT ARTERIES

In transposition of the great arteries (TGA), the aorta arises from the right ventricle and the pulmonary artery from the left ventricle. Consequently, the systemic venous return is pumped back into the body and the pulmonary venous return is ejected back into the lungs. Instead of having a normal in-series circulation, the TGA patients have parallel circulation. Without either an intra-cardiac or extra-cardiac shunt, the infants with TGA will not survive. The fetal circulatory pathways {PFO and patent ductus arteriosus (PDA)} will initially provide some mixing. In most neonates with TGA, the PFO and PDA tend to undergo spontaneous closure, resulting in a severely cyanotic infant. Rashkind balloon atrial septostomy has been extensively used in the palliation of the neonate with complete TGA. The improved mixing at the atrial level allows the neonate with transposition to grow up to an age (usually 3 to 6 months) at which time a venous switch (Mustard or Senning) procedure could safely be performed. With the introduction of arterial switch (Jatene) procedure which is usually performed at approximately one week of age, balloon atrial septostomy is not necessary in all babies. If naturally present PFO and/or Prostaglandin E1 (PGE1) infusion do not result in reasonably good oxygen saturations (60 to 70% without metabolic acidosis), balloon atrial septostomy should be performed, preparatory to arterial switch procedure.

### TRICUSPID ATRESIA

Tricuspid atresia may be defined as congenital absence or agenesis of the morphologic tricuspid valve [31,32]. In tricuspid atresia an obligatory right-to-left shunt occurs at the atrial level. Usually, this shunting is through a PFO. Because of the obligatory nature of the shunt, this fetal pathway persists in the postnatal period; this is in part related to low left atrial pressure. But, the entire systemic venous return must pass through the patent foramen ovale. Therefore, interatrial obstruction is anticipated, but very few patients with tricuspid atresia have clinically significant obstruction [33]. The right-to-left shunt occurs in late atrial diastole with augmentation during atrial systole ('a' wave [34,35]. A mean interatrial pressure difference greater than 5 mm Hg with very prominent 'a' waves (15 to 20 mm Hg) in the right atrium is generally considered to represent obstructed interatrial septum [33,36]. Balloon atrial septostomy [37], if unsuccessful, blade atrial septostomy [22,38], and rarely surgical atrial septostomy, may be necessary to relieve the obstruction. Significant interatrial obstruction requiring atrial septostomy in the neonate is rare and unusual although this can be a significant problem later in infancy [36,39].

#### PULMONARY ATRESIA WITH INTACT VENTRICULAR SEPTUM

Pulmonary atresia with intact ventricular septum is a complex cyanotic congenital heart defect characterized by complete obstruction of the pulmonary valve, two distinct ventricles, a patent tricuspid valve and no ventricular septal defect. The right ventricle is usually, but not invariably, small and hypoplastic. Because of atretic pulmonary valve, there is no forward flow from the right ventricle and the blood regurgitates back into the right atrium. Therefore, an obligatory right-to-left shunt occurs across the atrial septum. The objectives of any treatment plan are to achieve a four-chamber, bi-ventricular, completely separated circulation [40-43]. This aim may be achieved only in the absence of: a) right ventricular dependent coronary circulation b) severe right ventricular hypoplasia, and c) infundibular atresia. In the subgroup of neonates with unipartite or extremely small right ventricle with infundibular atresia and/or right ventricular dependent coronary circulation, a modified Blalock-Taussig shunt to provide pulmonary blood flow and atrial septostomy to decompress the right atrium should be performed. The usefulness of atrial septostomy in improving outcomes of systemic-topulmonary artery shunts in pulmonary atresia patients is demonstrated in early studies [44]. Balloon atrial septostomy, if unsuccessful blade atrial septostomy, and rarely surgical atrial septostomy, as described in tricuspid atresia section, may be necessary to promote egress of right atrial blood. However, it should be noted that no atrial septostomy should be undertaken in patients in whom transcatheter or surgical opening of the atretic pulmonary valve is contemplated: the objective in this scenario is to encourage anterograde pulmonary flow across the opened pulmonary valve.

#### HYPOPLASTIC LEFT HEART SYNDROME

In hypoplastic left heart syndrome (HLHS), there is hypoplasia of the left heart structures. Similar to other congenital heart defects, HLHS also shows a spectrum of severity. In the most severe form, aortic and mitral valves are atretic with a diminutive ascending aorta and markedly hypoplastic left ventricle [45,46]. The left atrium is usually smaller than normal. Because of obstruction at the mitral valve, pulmonary venous blood must cross the atrial septum via a PFO and mix with desaturated systemic venous blood in the right atrium. In some patients the PFO may





Working Together to Develop a Better Tomorrow

### December 2007

### CONGENITAL CARDIOLOGY TODAY



Figure 1. Selected cinfluroscopic frames of the Rashkind's balloon septostomy procedure. Note the position of the inflated balloon in the left atrium (A) and in right atrium and inferior vena cava in successive frames, as it is rapidly and forcefully withdrawn across the atrial septum (B,C, & D). After it reaches the inferior vena cava (D), it is rapidly advanced into the right atrium (E & F) in order not to inadvertently occlude the inferior vena cava in case of failure to deflate the balloon (which is quite rare).

be restrictive and occasionally the atrial septum may be intact. In the neonate, obstruction at the level of the PFO may be treated with conventional Rashkind balloon atrial septostomy [4]. However, since the left atrium is small, Rashkind septostomy may not be feasible. In addition, the septum may be too thick to be torn by balloon septostomy; therefore, Park blade septostomy [22] has been considered. But again, because of left atrial hypoplasia, blade septostomy may not be feasible. Static dilatation of the atrial septum [21,23,24] with a balloon angioplasty catheter may be used which may not only relieve the obstruction, but also keep some restriction such that there is no rapid fall in the pulmonary vascular resistance. Static balloon dilatation is preferred by the author. In some patients the atrial septum may be intact or have a tight patent foramen ovale which may not even allow passage of a catheter. In such situations, puncture of the atrial septum by Ross technique [27] or radiofrequency perforation of the atrial septum [28-30] followed by static balloon atrial septal dilatation [21,23,24] or stent implantation [25,26] may become necessary.

#### MITRAL ATRESIA

The considerations for mitral atresia are similar to those described for HLHS and the utility of balloon septostomy in this group of patients has been well documented [47,48].

### TOTAL ANOMALOUS PULMONARY VENOUS CONNECTION

In this entity, all the pulmonary veins drain into systemic veins, most commonly they drain into a common pulmonary vein which is then connected to left innominate vein, superior vena cava, coronary sinus, portal vein or other rare sites. Occasionally individual veins drain directly into the right atrium. Irrespective of the type, all pulmonary venous blood eventually gets back into right



Figure 2. Selected video frames showing a very small inter-atrial opening (A) which became larger (B) following balloon atrial septostomy.

atrium, mixes with systemic venous return and gets redistributed to the systemic and pulmonary circulations. The systemic flow is through the PFO. Consequently, restrictive PFO will cause decreased systemic perfusion and may indirectly result in pulmonary venous obstruction. In the neonate, the most common form is obstructive infra-diaphragmatic type causing severe pulmonary venous obstruction because the pulmonary venous return has to go through hepatic circulation. Occasionally, however, cardiac and supracardiac types may have restrictive PFO and in such patients balloon atrial septostomy is beneficial [49].

#### **Rashkind Balloon Atrial Septostomy**

In TGA patients who are stable, the usual hemodynamic data including cine-angiography, as needed, are performed. If the infant is unstable or has extremely low oxygen saturations, one may proceed directly with balloon septostomy. In such situations, aortic saturation and pressure pullback across the atria and echocardiographic size of atrial defect are recorded. The balloon septostomy procedure involves inserting a balloon septostomy catheter, usually via a sheath percutaneously placed in the femoral vein, into the left atrium via the PFO. The balloon is inflated with diluted contrast material to a sub-maximal amount (usually 3 ml) and rapidly pulled back across the atrial septum (Figure 1) after ensuring that the catheter tip is located in the left atrium either by lateral fluoroscopy or by echocardiography. Once the catheter is pulled back to the inferior vena cava, the

4

December 2007



Figure 3. Selected video frames showing a very small color flow jet across the inter-atrial opening (A) which became larger (B) following balloon atrial septostomy.

catheter should be rapidly advanced into the right atrium; all this is done as a single motion. The balloon should be deflated as the catheter is repositioned into the right atrium. This jerking motion of the contrast filled balloon catheter produces a tear in the lower margin of the PFO (septum primum) which is very thin and frail in the newborn. We usually perform one additional septostomy following what may be considered good septostomy.

Increase in systemic arterial oxygen saturation, disappearance of pressure gradient across the atrial septum and echographic increase in the size of the atrial defect (Figure 2) with nonrestrictive Doppler flow across the atrial septum (Figure 3) are demonstrated in successful procedures. In patients in whom atrial septostomy is performed to relieve interatrial obstruction, trans-atrial pressure gradient is reduced or abolished. Some cardiologists balloon-size the atrial defect both prior to and following balloon septostomy and this is another method of assessment of result of the septostomy.



DEPARTMENT O F PEDIATRICS (http://www-pediatrics.ucsd.edu) AND CHILDREN'S SPECIALISTS OF SAN DIEGO (http://childrensspecialists.com) are jointly recruiting a Director of Noninvasive Imaging for the unified Division of Pediatric Cardiology at Rady Children's Hospital, San Diego. This 220-bed facility serves as a major regional tertiary care hospital for children and is the major teaching facility for the Department of Pediatrics of the UCSD School of Medicine. This is a unique opportunity to develop an exceptional CT and MRI Imaging Program, in partnership with Rady Children's Hospital and the University of California San Diego, in the seventh largest and most desirable city in the USA. The successful candidate for this position should possess the qualifications for appointment, preferably, at the rank of Assistant or Associate Professor. Academic series will be determined based on the successful candidate's background and qualifications. Candidates must be Board Certified in Pediatric Cardiology and licensed or licensable to practice medicine in the State of California. This appointment will require demonstrated administrative capabilities, excellent skills in clinical care and teaching, and research accomplishment. The Division provides a full range of Pediatric Cardiology services. It currently has seven Pediatric Cardiologists, two Cardiothoracic Surgeons, and an ACGME approved fellowship program. The faculty performs approximately 400 cardiac catheterizations and 450 cardiac surgical procedures yearly. Extensive opportunities to perform clinical, epidemiologic or basic science research exist at UCSD and Rady Children's Hospital.

Applications received by March 1st or until the position is filled, will receive full consideration. Salary commensurate with University of California policy.

Please send Curriculum Vitae to: John Moore, M.D., email <u>imoore@rchsd.org</u> or by mail to 3020 Children's Way, MC 5004; San Diego, CA 92123.

UCSD is an AA/EOE with a strong institutional commitment of excellence through diversity.

PFO Detection with Power M-Mode Doppler • Higher Sensitivity than TEE\* • Non-invasive method enables a measured valsalva strain • Easy to perform, office-based testing °J of Neuroimaging, Vol 14, No 4, Oct 2004.



5

### December 2007

6

### CONGENITAL CARDIOLOGY TODAY

In the initial description of balloon septostomy by Rashkind [4], the catheter was introduced into the femoral vein by cut-down. To avoid femoral venous cut-down, insertion of the catheter and performance of balloon septostomy via the umbilical vein [50] has been advocated. When percutaneous technology became available, the balloon catheter was introduced via an appropriate sized percutaneously inserted femoral venous sheaths [51,52].

"In the vast majority of the patients, the septostomy procedures are successful in creating an appropriate sized opening. In the rare cases, surgical septostomy may be required."

Our first choice is to perform balloon septostomy via the umbilical venous route. Therefore, we encourage our neonatology colleagues to place an umbilical venous line early on, with its tip well into the right atrium, before the ductus venosus constricts. At the time of septostomy, this line is exchanged over a wire with an appropriate-sized sheath.

The feasibility of performing balloon septostomy bedside, under echo guidance, has been demonstrated [53,54]. But, most cardiologists perform the procedure in the catheterization laboratory which is preferred.

Initially, Rashkind balloon septostomy catheters (USCI, Boston, MA) were used. Because the catheters were straight, sometimes making it difficult to advance the catheter into the left atrium, and because of the limited volume of fluid that these balloons would take, most cardiologists have switched to Edwards septostomy catheters (American Edwards Baxter, McGow Park, IL). These catheters have a gentle curve at the tip, facilitating easy access into the left atrium and larger volume of fluid could be injected into these balloons. More recently, atrioseptostomy catheters (B/Braun, Bethlehem, PA) have become available. There are no studies comparing the relative effectiveness of the available catheters and therefore, the selection of the type of catheter used is at the discretion of the operator.

#### **Blade Atrial Septostomy**

In older patients and in some conditions such as hypoplastic left heart syndrome, the lower margin of the PFO is thick and can't be ruptured by conventional balloon septostomy. The septostomy may simply stretch and not tear the lower margin of the PFO. Park and



PEDIATRIX.

# PEDIATRIC CARDIOLOGISTS

### Central and South Florida

Central and South Florida offer a world-class lifestyle, great schools and enviable year round activities. Because of exceptional growth, we have openings for three BC/BE Pediatric Cardiologists that are seeking to make a difference. Our practices are mature and established and are all located in growing, family-friendly communities. As part of a national company, both research and leadership opportunities are available.

Pediatrix offers physicians competitive salaries and excellent benefits, including professional liability insurance, CME allowance, comprehensive health/life benefits, stock purchase plan and 401(k).

For more information, please contact Jenifer Krutchik, Physician Relations Specialist, jenifer\_krutchik@pediatrix.com 800.243.3839.ext. 5165

Pediatrix Medical Group 1301 Concord Terrace, Sunrise, FL 33323 www.pediatrix.com



### HYPERTROPHIC CARDIOMYOPATHY ASSOCIATION 11TH ANNUAL HCMA MEMBERS MEETING May 30-June 2, 2008 Morristown, NJ

www.4hcm.org

www.CongenitalCardiologyToday.com



The Division of Pediatric Cardiology at the University of Virginia Children's Hospital is seeking to expand our current program of 12 cardiologists and 2 surgeons. A full range of services is provided including pediatric cardiac transplantation, a dedicated adult congenital heart disease program and full service interventional cardiology and Electrophysiology programs. There is an active pediatric cardiology fellowship program funded by an NIH sponsored training grant.

The division seeks to expand with the following faculty opportunity:

**NONINVASIVE PEDIATRIC CARDIOLOGIST:** The Division seeks an academic minded pediatric cardiologist to direct the non-invasive service and attend on the inpatient cardiology service. Skills in transthoracic, transesophageal and fetal echocardiography are required. Other duties for this position will depend on applicant interest but could include exercise physiology or preventive cardiology. Academic duties will include teaching fellows, residents and medical students and participation in clinical research. Applicants are required to be board eligible or board certified in Pediatric Cardiology. Faculty appointment will be at the Assistant or Associate Professor level.

Interested persons should send a cover letter expressing their interest and qualifications along with a curriculum vita to:

G. Paul Matherne MD Professor of Pediatrics Division Head Pediatric Cardiology University of Virginia Health System PO Box 800386 Charlottesville, VA 22908-0386 phone: (434)982-0260 fax: (434)924-5656 gpm2y@virginia.edu

Position will remain open until filled.



Figure 4. Selected cinfluroscopic frames of the blade septostomy procedure. The top three figures (A,B & C) show the position of the catheter with the blade open in the posterior-anterior projection while bottom three (D,E & F) show the lateral view. Note the position of the blade pointing to left and anteriorly. This procedure is performed by slow withdrawal of the blade catheter in contradistinction to very rapid pullback in the Rashkind's balloon septostomy procedure.

his associates [22] developed catheters with build-in blade (knife) (Cook Inc., Bloomington, IN) to address such thick atrial septae. Three blade sizes are commercially available. The selected catheter (smallest size for the newborn) is positioned across the PFO, the position of the tip of the catheter is confirmed and the blade opened. While pointing the blade anteriorly and to the left (Figure 4), the catheter is slowly withdrawn (not a jerky motion as in balloon septostomy), thus cutting the lower margin of the PFO. This is repeated one or two more times, varying the angle slightly. This is followed by balloon septostomy. Evaluation of the results is similar to that described in the balloon septostomy section. Success rate ranged between 70 and 90% [55,56].

#### **Balloon Angioplasty**

Mitchell et al [57] and Sideris et al [58] performed static dilatation of the atrial septum successfully in animal models. The first clinical application was reported by Shrivastava and her colleagues. [23]. The procedure involves advancing either an end-hole or a multipurpose catheter from the right atrium to the left atrium across the PFO and from there into a left pulmonary vein (we prefer left lower). An exchange length, extra-stiff Amplatz guide wire is positioned in the pulmonary vein via the catheter and the



Eleventh Annual Update of Pediatric Cardiovascular Disease

New and Evolving Concepts and Practices www.chop.edu/cardiology2008

February 6-10, 2008

Hyatt Regency Scottsdale Resort and Spa Gainey Ranch \* Scottsdale, AZ



# CARDIOLOGIST Denver, Colorado

Seeking a 4th BC/BE pediatric cardiologist to join a well-established practice. This busy group practices in two office locations, admits patients to two major hospitals and does consults at several local community hospitals. The ideal candidate will be comfortable doing transthoracic echocardiography independently. This group works closely with our affiliated group of Neonatologists, Pediatric Hospitalists/Intensivists and Maternal-Fetal Medicine Specialists. Denver, ranked as one of the top 10 places to live and work, offers a diverse business economy, top-ranked educational institutions, vibrant cultural opportunities and championship sports teams as well as an abundance of recreational activities.

Pediatrix offers physicians competitive salaries and excellent benefits, including professional liability insurance, CME allowance, comprehensive health/life benefits, stock purchase plan and 401(k).

For more information, please contact Lori Abolatia lori\_abolatia@pediatrix.com 800.243.3839.ext. 5209

Pediatrix Medical Group 1301 Concord Terrace, Sunrise, FL 33323 www.pediatrix.com



Figure 5. Selected cinfluroscopic frames of balloon angioplasty procedure (to enlarge the patent foramen ovale) demonstrating an inflated balloon in the lateral view showing waisting of the balloon (A) which was completely abolished following further inflation of (B) the balloon (B).

catheter removed, leaving the guide wire in place. Selected balloon angioplasty catheter is advanced over the wire, positioned across the atrial septum and the balloon inflated to 3 to 5 atmospheres of pressure (Figure 5), taking care not to inadvertently dilate the pulmonary vein. The balloon inflation is repeated once or twice. The recommended duration of inflation is 5 seconds with a 5-minute interval in between each dilatation. Waisting of the balloon during the initial phases of balloon inflation and the disappearance of the waist indicates that the PFO is stretched beyond its initial size. Lack of waist in subsequent balloon inflations indicates that there is enlargement of the PFO. There are no data to indicate the most appropriate size of the balloon that should be used; 8 to 20 mm balloons have been used [21]. Based on theoretical considerations, the final diameter of the balloon should be three to four times the echographic size of the PFO is a good choice [21]. The balloon size should not exceed the size of the atrial septum. We usually end up with a balloon diameter of 14 to 15 mm. Evaluation of the results, again, is similar to that described in the balloon septostomy section. Increase in the size of the defect (similar to Figure 2) and in color Doppler flow width (Figure 6) demonstrates success of the procedure. There are only anecdotal reports of success, but no systematic studies to evaluate the results are published.

### **Atrial Septal Perforation**

In a small percentage of patients, the atrial septum is intact (no PFO), particularly in HLHS. In such situations the septum can't be crossed with conventional catheters. Traversing the septum either by Ross/Brockenbrough's transseptal technique or radiofrequency perforation may be required.

Transseptal catheterization was initially described by Ross, Brockenbrough and their colleagues [59,60]. This was subsequently adapted to pediatric patients [61,62]. More recently, the





For information, registration, and abstract submission go to: www.picsymposium.com

8



Figure 6. Selected video frames showing a very small color flow jet across the inter-atrial opening (A) which became larger (B) following balloon angioplasty of the atrial septum.

technique was extended to neonates [27]. In the Ross technique, initially an end-hole or multipurpose catheter is advanced from the femoral vein into the superior vena cava through which a guide wire is introduced and the catheter removed. The transseptal sheath assembly (Cook, Bloomington, IN) is advanced over the wire and the wire removed. The tip of the transseptal catheter should be pointing posteriorly and to the left (medial) and slowly withdrawn under fluoroscopy in the lateral view. As the tip of the catheter enters the right atrium, it flips suddenly; at this point the position of the tip of the catheter against the atrial septum is confirmed by echocardiography, indenting the atrial septum. The needle of the transseptal assembly is gently advanced across the atrial septum; a slight "give" indicates entry into the left atrium. The cannula is advanced slightly and the needle removed and the pressure recorded, confirming the left atrial position. Once confirmed to be in the left atrium, the dilator and the sheath assembly is advanced into the left atrium and the dilator removed leaving the sheath in place.

In the radiofrequency (RF) perforation procedure, the technique is similar except that the RF wire is used instead of the needle. Once in the left atrium, static dilatation of the atrial septum as described in the preceding section or stent implantation (to be described in the next section) is performed.

#### **Stent Implantation**

Because of tendency for closure of dilated atrial septal openings, stents to keep the defects open may have to be used [25,26]. Stents should be implanted only across the highly restrictive PFO or after septal perforation by one the above described techniques. Initially the guide wire is positioned into the left lower pulmonary vein and an appropriately sized delivery sheath is positioned across the atrial septum. The stentmounted balloon catheter (large coronary or small peripheral stent of approximately 10 to 15 mm length) is advanced over the wire, but within the sheath. Based on boney land marks the stent is positioned in such way that it is centered across the atrial septum. The delivery sheath is withdrawn into the right atrium while keeping the stent catheter in place. A combination of test injection via the side arm of the sheath and echocardiogram (trans-thoracic or transesophageal) is used to ensure correct position of the stent. The balloon is inflated at the manufacturer's stated pressure, thus expanding the stent. The balloon is deflated and removed, taking care not to dislodge the stent. This is followed by the removal of the guide wire. Recording of pressures across the atrial septum, via the stent, is not recommended to reduce the probability of stent dislodgement. Echo is used to record Doppler flow velocity across the stent. Low velocity flow indicates good result.

#### CONCLUSIONS

There are a number of cardiac defects in which an atrial septal defect is beneficial. But, the naturally occurring PFO undergoes spontaneous closure, causing poor mixing and/or obstruction to systemic or pulmonary venous flow. In such situations, the PFO may be enlarged or an atrial septal defect created by transcatheter methods. The selection of the method used is largely based on the atrial septal anatomy and left atrial size. In the vast majority of the patients, the septostomy procedures are successful in creating an appropriate sized opening. In the rare cases, surgical septostomy may be required.

#### References

- Rubio-Alvarez V, Limon-Lason R, Soni L. Valvalotomias intracardiacas por medico de un cateter. Arch Inst Cardiol Mexico 1953; 23: 183-92.
- Rubio V, Limon-Lason R. Treatment of pulmonary valve stenosis and of tricuspid valve stenosis using a modified catheter [abstr]. Second World Congress of Cardiology, Program Abstract 1954; 11: 205.
- Dotter CT, Judkins MP. Transluminal treatment of arteriosclerotic obstruction: description of a new technique and a preliminary report of its application. Circulation 1964; 30: 654-70.
- 4. Rashkind WJ, Miller WW. Creation of an atrial septal defect without thoracotomy. J Am Med Assoc 1966; 196: 991-2.
- Porstmann W, Wierny L, Warnke H. Der Verschluss des Ductus arteriosus persistens ohne thorakotomie (1, Miffeilung), Thoraxchivargie 1967; 15: 109-203.
- Porstmann W, Wierny L, Warnke H, et al. Catheter closure of patent ductus arteriosus: 62 cases treated without thoracotomy. Radiol Clin North Am 1971; 9: 203-18.
- King TD, Thompson SL, Steiner C, Mills NL. Secundum atrial septal defect: non-operative closure during cardiac catheterization. J Am Med Assoc 1976; 235: 2506-9.
- 8. Mills NL, King TD. Non-operative closure of left-to-right shunts. J Thorac Cardiovasc Surg 1978; 72: 371-8.
- Kan JS, White RJ, Jr, Mitchell SĚ, Gardner TJ. Percutaneous balloon valvuloplasty: a new method for treating congenital pulmonary valve stenosis. New Engl J Med 1982; 397: 540-2.



### Division of Pediatric Cardiology Saint Louis University School of Medicine Cardinal Glennon Children's Medical Center

Saint Louis University, a Catholic, Jesuit institution dedicated to student learning, research, health care, and service is seeking additional pediatric cardiologists to join an established group within the Division of Cardiology and the Department of Pediatrics at Cardinal Glennon Children's Medical Center. Applicants will be considered at the Assistant/Associate Professor rank, and must be board certified/ eligible in Pediatric Cardiology. General responsibilities will include clinical care, teaching, and research.

### Interventional Cardiologist

We are seeking an invasive cardiologist to assist our current faculty with the growing number of cardiac catheterizations and interventional cases. A complete spectrum of pediatric interventional procedures is currently being performed. An interest in clinical research is encouraged. Academic rank will be commensurate with qualifications and experience.

### **Non-Invasive Cardiologist**

We are also seeking a non-invasive cardiologist to participate in inpatient and outpatient cardiology care as well as echocardiography services. Interest and experience in cardiac MRI techniques a plus, but not required. Significant participation in and direction of an active outreach clinic program is possible if desired. An interest in clinical research is encouraged. Academic rank will be commensurate with qualifications and experience.

The cardiology division is in a state of rapid growth, with construction of a dedicated Congenital Heart Center within Cardinal Glennon Children's Medical Center projected to begin in 2008. An active congenital heart surgery program exists, and a new addition to the hospital recently opened housing state-of-the-art operating rooms and a new 60-bed neonatal intensive care unit. A new 10-story tower that will serve as the Health Sciences Center Research Building will be completed in 2007.

Interested candidates must submit a cover letter, application, and current CV to **http://jobs.slu.edu**. Other correspondence regarding this position can be sent to Kenneth O. Schowengerdt, MD, Wieck-Sullivan Professor and Director of Pediatric Cardiology, Saint Louis University School of Medicine, 1465 South Grand Blvd, St. Louis, MO 63104. Telephone: (314)-577-5633; Fax: (314)-268-4035; email: schowko@slu.edu.

Saint Louis University is an Affirmative Action, Equal Opportunity Employer, and encourages nominations of and applications from women and minorities.

- 10.Gruntzig AR. Transluminal dilatation of coronary artery stenosis. Lancet 1978; 1: 263.
- 11.Rao PS. Transcatheter management of heart disease in infants and children. Pediat Rev Comm. 1987; 1: 1-18.
- 12.Rao PS. Medical Progress: Balloon valvuloplasty and angioplasty in infants and children. J Pediat 1989; 114: 907-14.
- Rao PS. Balloon Angioplasty and Valvuloplasty in Infants, Children and Adolescents. Current Problems in Cardiology. Yearbook Medical Publishers, Inc. Chicago, 1989; 14(8): 417-500.
- 14.Rao PS (Ed). Transcatheter Therapy in Pediatric Cardiology. Wiley-Less, Inc., New York, 1993.
- 15.Rao PS. Interventional pediatric cardiology: state of the art and future directions. Pediat Cardiol 1998; 19: 107-24.
- 16.Singer MI, Rowan M, Dorsey TJ. Transluminal aortic balloon angioplasty for coarctation of the aorta in the newborn. Am Heart J 1982; 103: 131-2.
- 17.Tynan M, Jones O, Joseph MC, et al. Relief of pulmonary valve stenosis in first week of life by percutaneous balloon valvuloplasty. Lancet 1984; 1: 273.
- Lababidi Z, Weinhaus L. Successful balloon valvuloplasty for neonatal critical aortic stenosis. Am Heart J 1986; 112: 913-6,
- 19.Rao PS. Balloon angioplasty for coarctation of the aorta in infancy. J Pediat 1987; 110: 713-8.
- 20.Wren C, Sullivan I, Bull C, Deanfield J. Percutaneous balloon dilatation of aortic valve stenosis in neonates and infants. Br Heart J 1987: 58: 608-12.
- 21.Rao PS. Static balloon dilation of atrial septum (Editorial). Am Heart J 1993; 125: 1826-7.
- 22.Park SC, Neches WH, Zuberbuhler JR, et al. Clinical use of blade septostomy. Circulation 1978; 58: 600-6.
- 23.Shrivatsava S, Radhakrishnan S, Dev V, et al. Balloon dilatation of atrial septum in complete transposition of great arteries - a new technique. Indian Heart J 1987; 39: 298-300.
- 24.Rao PS. Static balloon dilation of restrictive atrial septal defects (Editorial). J Saudi Heart Assoc 1992; 4: 55-8.
- 25.Gewillig D, Boshoff L, Mertens L. Creation with a stent of an unrestrictive lasting atrial communication, Cardiol Young 2002; 12: 404–7.
- 26.Eicken H, Gildein, C, Schreiber C, et al. Stenting of a restrictive foramen ovale in a patient with hypoplastic left heart syndrome. Internat J Cardiol, 2006; 113: 254-6 A.

- 27.Atz AM, Feinstein JA, Jonas RA, et al. Preoperative management of pulmonary venous hypertension in hypoplastic left heart syndrome with restrictive atrial septal defect. Am J Cardiol 1999; 83: 1224–8.
- 28.Justino H, Benson LN, Nykanen DG. Transcatheter creation of an atrial septal defect using radiofrequency perforation. Catheter Cardiovasc Intervent. 2001; 54: 83-7
- 29.Sakata Y, Feldman T. Transcatheter creation of atrial septal perforation using a radiofrequency transseptal system: novel approach as an alternative to transseptal needle puncture. Catheter Cardiovasc Intervent. 2005; 64: 327-32.
- 30.Hill SL, Mizelle KM, Vellucci SM, et al. Radiofrequency perforation and cutting balloon septoplasty of intact atrial septum in a newborn with hypoplastic left heart syndrome using transesophageal ICE probe guidance, Catheter Cardiovasc Intervent 2005; 64: 214–7.
- 31.Rao PS. A unified classification for tricuspid atresia. Am Heart J 1980; 99: 799-804.
- 32.Rao PS. Terminology: tricuspid atresia or univentricular heart? In: Rao PS (Ed): Tricuspid Atresia, Mount Kisco, NY, Futura Publishing Co, 1982: 3-6.
- 33.Dick M, Fyler DC, Nadas AS. Tricuspid atresia: clinical course in 101 patients. Am J Cardiol 1975; 36: 327-37.
- 34.Rao PS. Left-to-right shunting in tricuspid atresia. Br Heart J 1983; 49: 345-9.
- 35.Rao PS. Cardiac catheterization in tricuspid atresia. In: Rao PS (Ed): Tricuspid Atresia, Mount Kisco, NY: Futura Publishing Co, 1982: 153-78.
- 36.Rao PS, Covitz W, Chopra PS. Principles of palliative management of patients with tricuspid atresia. In: Rao PS (Ed): Tricuspid atresia, ed 2. Mount Kisco, NY: Futura Publishing Co, 1992: 297-320.
- 37.Rashkind WJ, Waldhausen JA, Miller WW, et al. Palliative treatment of tricuspid atresia: combined balloon atrial septostomy and surgical alteration of pulmonary blood flow. J Thorac Cardiovasc Surg 1969; 57: 812-8.

- 38.Rao PS. Transcatheter blade atrial septostomy. Cathet Cardiovasc Diagn 1984; 10: 335-42.
- 39.Rao PS. Natural history of the ventricular septal defect in tricuspid atresia and its surgical implications. Br Heart J 1977; 39: 276-88.
- 40.Rao PS, Liebman J, Borkat G. Right ventricular growth in a case of pulmonary stenosis with intact ventricular septum and hypoplastic right ventricle. Circulation 1976; 53: 389-94.
- 41.Rao PS. Comprehensive management of pulmonary atresia with intact ventricular septum. Ann Thorac Surg 1985; 40: 409-13.
- 42.Siblini G, Rao PS, Singh GK, et al. Transcatheter management of neonates with pulmonary atresia and intact ventricular septum. Cathet Cardiovasc Diagn 1997; 42: 395-402.
- 43.Rao PS. Pulmonary atresia with intact ventricular septum. Current Treatment Options in Cardiovasc Med 2002; 4: 321-36.
- 44.Shams A, Fowler RS, Trusler GA, et al. Pulmonary atresia with intact ventricular septum: report of 50 cases. Pediatrics 1971; 47:370-6.
- 45.Rao PS, Striepe V, Merrill WM. Hypoplastic left heart syndrome. In: Kambam J (Ed), Cardiac Anesthesia for Infants and Children. St. Louis, Mosby, 1993: 299-309.
- 46.Rao PS, Turner R, Forbes TJ. Hypoplastic left heart syndrome. In emedicine – Pediatrics http://www.emedicine.com.
- 47.Mickell JJ, Mathews RA, Park SC, et al. Left atrio-ventricular valve atresia: clinical management. Circulation 1980; 61: 123-7.
- 48.Rao PS, Kulangara RJ, Moore HV and Strong WB. Syndrome of single ventricle without pulmonic stenosis but with left atrioventricular valve atresia and interatrial obstruction: palliative management with simultaneous atrial septostomy and pulmonary artery banding. J Thorac Cardiovasc Surg 1981; 81: 127-130.
- 49.Serrato M, Buchelers HG, Bicoff P, et al. Palliative balloon atrial septostomy



### Interventional Pediatric Cardiologist

The Department of Pediatrics at Louisiana State University Health Sciences Center in conjunction with Children's Hospital, New Orleans, LA, is seeking a BE/BC interventional pediatric cardiologist to join the faculty of a busy academic clinical and surgical heart program. This will be an open rank position, (Assistant -Full Professor, Clinical), with rank determined by the candidates' credentials and experience. Currently, approximately 300 cardiac catheterizations and 400 cardiothoracic surgeries are performed in infants, children and young adults each year. Opportunities are available for both clinical and basic research.

Interested applicants should submit c.v. and 3 letters of recommendation to:

Robert J. Ascuitto, M.D., Ph.D. Professor, Department of Pediatrics Director, Pediatric Cardiology LSU Health Sciences Center, N.O. Children's Hospital 200 Henry Clay Ave. New Orleans, LA 70118 FAX: (504) 896-3952 E-Mail: nrossa@lsushc.edu

LSUHSC is an AA/EOE



 17<sup>th</sup> Utah Conference on Congenital Cardiovascular Disease February 24 – 27, 2008
 Intermountain

 The Cliff Lodge at Snowbird Ski Resort, Snowbird, Utah
 Intermountain

Please visit our web site: www.primarychildrens.com/medicalclasses and download a PDF file of the registration form.

### One-Year Clinical Lecturer Appointment for Advanced Experience in Pediatric Cardiac Intensive Care

The University of Michigan Congenital Heart Center is pleased to offer a 1-year clinical lecturer appointment for advanced experience in pediatric cardiac intensive care at C.S. Mott Children's Hospital, University of Michigan. The primary focus of this position is to provide the clinician with additional experience in the preand post-operative management of all forms of congenital and acquired heart disease in children. The majority of the training will take place in a state-of-theart 15-bed pediatric cardiothoracic intensive care unit located in C.S. Mott Children's Hospital, one of only a handful of intensive care units in the country dedicated entirely to children with cardiac disease. The applicant will also have an opportunity for additional experience in cardiac anesthesia, pediatric cardiac surgery (and perfusion techniques), advanced pediatric critical care medicine, adult cardiac intensive care, extracorporeal life support (including cardiac ECMO, ventricular assist devices, etc.), and pediatric cardiac transplantation. Candidates should be board-certified in pediatrics and board-certified/board-eligible in pediatric cardiology.

The University of Michigan Congenital Heart Center is a world-renowned treatment facility within C.S. Mott Children's Hospital, University of Michigan. Care is provided by a team of over 20 highly-specialized pediatric cardiologists, 4 cardiac surgeons, 15 pediatric cardiology fellows, pediatric residents, nurses and nurse practitioners, anesthesiologists and allied health professionals. The University of Michigan is an Affirmative Action/ Equal-Opportunity Employer. Minorities and women are encouraged to apply.

Interested physicians may contact or mail their CVs and 3 letters of reference (including one from their fellowship Program Director) to:

John R. Charpie, MD, PhD Medical Director, Pediatric Cardiothoracic Intensive Care Unit University of Michigan Congenital Heart Center C.S. Mott Children's Hospital L1242 Women's, Box 0204 1500 East Medical Center Drive Ann Arbor, Michigan 48109-0204 Tel: 734-936-6703 FAX: 734-936-9470 e-mail: jcharpie@umich.edu



- 50.Abinader E, Zeltzer M, Riss E. Transumbilical atrial septostomy in the newborn. Am J Dis Child 1970; 119: 354-5.
- 51.Hurwitz RA, Girod DA. Percutaneous atrial septostomy in infants with transposition of the great arteries. Am Heart J 1976; 91: 618-22.
- 52.Sunderland CO, Nichols GM, Henken DP, et al. Percutaneous cardiac catheterization and atrial balloon septostomy in pediatrics. J Pediat 1976; 89: 584-7.
- 53.Baker EJ, Allan LD, Tynan M, et al. Balloon atrial septostomy in the neonatal intensive care unit. Br Heart J 1984; 51: 377-8.
- 54.Bullaboy CA, Jennings RB, Jr, Johnson DH. Bedside balloon atrial septostomy using echocardiographic monitoring. Am J Cardiol 1984; 53: 971-2.
- 55.Park SC, Neches WH, Mullins CE, et al. Blade atrial septostomy: collaborative study. Circulation 1982; 66: 258-66.
- 56.Park SC, Neches WH. Blade atrial septostomy: In: Rao PS (Ed). Transcatheter Therapy in Pediatric Cardiology. Wiley-Liss, Inc., New York, 1993: 17-27.
- 57.Mitchell SE, Kan JS, Anderson JH, et al. Atrial septostomy: stationary angioplasty balloon technique. Pediatr Res 1986; 20: 173A.
- 58.Sideris EB, Fowlkes JP, Smith JE, et al. Why atrial septostomy and not foramen ovale angioplasty? Annual Symposium of Texas Heart Institute, Sept 28 – Oct 1, 1988:36.
- 59.Ross J, Jr, Braunwald E, Morrow AG. Transseptal left atrial puncture: New technique for the measurement of left atrial pressure in man. Am J Cardiol 1959; 3: 653-5.
- 60.Brockenbrough EC, Braunwald E, Ross J, Jr. Transseptal left heart catheterization: A review of 450 studies and description of an improved technique. Circulation 1962; 25: 15-22.
- Duff DF, Mullins CE. Transseptal left heart catheterization in infants and children. Cathet Cardiovasc Diagn. 1978; 2: 213-23.
- 62.Ali Khan MA, Mullins CE, Bash SE, et al. Transseptal left heart catheterization in infants, children, and young adults. Cathet Cardiovasc Diagn. 1989; 17: 198-201.

### ССТ

P. Syamasundar Rao, MD Professor and Director Division of Pediatric Cardiology Univ. of Texas/Houston Medical School 6431 Fannin, MSB 3.130 Houston, TX 77030 USA (P) 713-500-5738; (F): 713-500-5751

P.Syamasundar.Rao@uth.tmc.edu

Heart Valves • Cannulae Oxygenators & Filters • RVOT Conduits Ablation Technologies • Pacemakers • ICDs



### Interview with Richard Van Praagh- Part 2: Family Life

By Bradley W. Robinson, MD

This is the last in a series of two articles drawn from an interview by Dr. Bradley W. Robinson with a pioneer in the field of pediatric cardiology, Richard Van Praagh. The first article "Academic Career," appeared in the October 2007 edition. It is available on the web in a PDF file.

## **Q:** So you grew up in Toronto? Tell me about your family.

I was actually born in London, Ontario and moved to Toronto at the age of 2 with my family... and grew up in Toronto. I went to school there and went to ... the University of Toronto.

My father was a stockbroker from London, England. His dad had a seat on the London Exchange and... my Dad came to Canada in 1920 and was doing very well as a stockbroker. Then 1929 came along, which was the great crash. I was born in 1930 and my brother Ian in 1931. Ian is also a physician. He is the sensible member of our family. He is an obstetrician-gynecologist in New York.

Well, in order to go to university, this is very likely to require a little bit of money, so we started working in the summertime. I started probably when I was 12, and ... I started to work in Toronto's biggest cemetery, Mount Pleasant Cemetery, raking grass then cutting grass with a hand mower. We would walk, oh, 15-20 miles every day. It was hard work and these were early post Depression times. If you were caught talking on the crew, you would be fired immediately, for example. Then I got a power mower after that and then I became a gravedigger, which was when it was done by pick and shovel and not a backhoe digger the way it now is done. Then they decided that they were going to raise the pay of everybody in our cemetery except for high school kids like us.

So that's when I started working on construction. And again, it was a non-union crew and started often around 6-6:30 am in the morning and we worked until 8 or 9 pm at night doing pick and shovel labor building roads, bridges, dams, sewers. Then I became a dump truck driver and Ian was the roller operator. I also

ran a small bulldozer, something called a front-end loader and this was right after the war. We got to know people from all over the world. I felt like Homer listening to the heroes come home from the war.

### Q: What else did you do in the Summer?

Oh, we were able to make our fees, books, clothes, instruments a lot more and able to save money too doing this and for us it was an escape from the prison of privilege. We went to wonderful schools. We went to the Conservatory of Music. But we always worked incognito. Nobody ever knew we were medical students. We wanted to be treated just like everybody else, as we were, and English was seldom spoken on the crew. That's when Ian and I started to use our high school French.

### Q: So how did this help you in medicine?

Well, it was a real world experience out of a very privileged home, with excellent schools, the Conservatory of Music, and so on. The combination was the thing. A physician's job, as much as anything else is people, and this was a marvelous introduction to people from all over the world. For example, we had people on our crew who fought on both sides at Stalingrad.

### **Q:** I understand you sold brushes?

I became a Fuller Brush man selling Fuller brushes door-to-door and I became the Canadian champion at selling Fuller brushes. My stakes were probably quite a bit higher than most other peoples' were, but this was all business. You know, you did your own sales. You handled your own complaints. You did your own deliveries. I would find myself telling people, "No, no, don't buy this. Get this. It's cheaper, and it is better," you know?

# **Q:** Did you show an interest in science in undergrad and college?

Yes, I have always loved science...[and] people. Our neighbor said, "Dick, it is no surprise that you went into pediatrics. Children and dogs all love you." Oh I love all kinds of sports too such as skiing and swimming and sailing. I used to be a mem-

Committed to providing more options for the lifetime care of patients with congenital heart disease



### CONGENITAL CARDIOLOGY TODAY



# PEDIATRIC CARDIOLOGIST San Antonio, Texas

Well-established, full service pediatric cardiology practice in San Antonio seeks an additional pediatric cardiologist to join our five-member team. We have hospital affiliations with excellent cardiac catheterization labs, state-of-the-art imaging facilities, and an open-heart surgical program with a PICU covered 24/7 by board certified intensivists. This is an outstanding opportunity for someone who is a team player. Must be BC/BE in pediatric cardiology with an interest in all clinical aspects of the practice, including servicing outreach clinics within South and West Texas. Recent and forthcoming fellowship graduates are encouraged to apply. San Antonio, located on the edge of the Texas Hill Country, is a modern, historic, vacation city with many cultural and recreational attractions.

Pediatrix offers physicians competitive salaries and excellent benefits, including professional liability insurance, CME allowance, comprehensive health/life benefits, stock purchase plan and 401(k), EOE.

For more information contact Karen Everest, Pediatric Cardiology Associates, 210.614.3264, ext. 122, 210.614.3921(fax). karen\_everest@pediatrix.com

### **Pediatric Cardiology Associates**

an affiliate of Pediatrix Medical Group, Inc. 4499 Medical Drive, Suite 272, San Antonio, TX 78229 www.pediatrix.com ber of the baseball team. I got a broken thumb, and as a result it was a bad catch.

### Q: How did you get interested in medicine?

14

My great-grandfather was a speech therapist. He was the Dutchman. His name was Willem, which is William, and Willem van Praagh was the person who introduced the system of teaching the congenitally deaf, and consequently dumb, how to speak by lip reading and vibrations, rather than by signing. In other words, how to do what Helen Keller did. This was known as the Continental Method. He wasn't a physician. He was a speech therapist who came to Britain and came to know the royal family quite well because they had a terrible problem with speech impediments, which has run in their family. For example, George VI, King George VI, during World War II, would make broadcasts to the Empire, to the world, etc., and his stuttering was so bad that it was really painful to listen to.

My grandfather and father were both stockbrokers. Then after 1929, my dear father went into the car business. He was the best Chrysler products salesman in Canada for years and years and years. He was a wonderful salesman who had renounced stocks and bonds for the reasons I mentioned. I wanted to do the best thing that I possibly could do with my life. So at late 14 or thereabouts, I came home and announced to my parents that I was going to become a minister, an Anglican minister, church of England. My dear mom, who was a Scottish Canadian, born a Presbyterian, said, "Well now, Richard, that's very interesting. But don't you think it might be a good idea, before you make a final decision, to go to university, study a little philosophy, a little history, a little comparative religion, and then make up your mind." So I did that, and then I finally decided that I would go in to medicine, in which I would be able to minister to the body as well as the soul. And here I am now, a cheerful agnostic and terribly pleased because of what my dear mother ever so gently and perceptively suggested to me.

# **Q:** So by taking those different courses in the university, you developed an interest in medicine?

Oh, in our family, we had always regarded medicine as the noblest of the professions, the finest thing that one could do was the feeling. You know that one of the tragedies in life is that you have to make decisions - I decided to apply to medical school. Amazingly, in retrospect, I only applied to the University of Toronto, not to any other medical school. If I hadn't been accepted, I don't know what I would have done.

### **Q:** What did your mother do?

Well, she was the second university graduate in her family. She graduated in 1924 from University College at the University of Toronto in liberal arts and then she taught French and art in St.





Working Together to Develop a Better Tomorrow

### CONGENITAL CARDIOLOGY TODAY

15

Catherine's for about two years before getting married to this entrancing Englishman.

My mom had me singing French songs before I went to kindergarten. She was charming, highly educated, sophisticated...one of our very best friends, to my brother lan and me.



# **Q:** Where did your parents get married and why did they move to Toronto?

They got married in St. Catherine's and then lived in London, Ontario for a while, and then moved to Toronto in 1932 when I was two. Even at that time, it was obvious that Toronto was going to become the economic capitol of Canada, which it now is, with all due respect to Montreal, which is still a marvelous place, but Toronto is now economically preeminent in Canada.

# **Q:** What were your favorite extracurricular activities?

Oh gosh, I was on the gym team for example. Gymnastics. I think my height of gymnastic accom-



## **PEDIATRIC CARDIOLOGISTS**

HCA is the largest healthcare company in the US, with over 170 hospitals in 21 states. Whether you are looking for a place to start, or somewhere to complete a successful career in pediatric cardiac care, chances are we can help you find it.

We have opportunities in big cities and rural areas, near sandy beaches and skyward reaching mountains. Wherever your internal compass points, it's likely that we can help guide you to a destination that will match your personal preferences while fulfilling your career dreams.

As our co-founder, Dr. Thomas Frist, Sr. said, "Bettering the human condition is the greatest good any individual can achieve." You are doing your part by taking care of our most precious commodity, the children who will be our future. Now let us do our part by helping you find your dream job.

Call or e-mail me today for more information on opportunities in your sub-specialty.



If you are a fellow in training, please be sure to ask about our excellent stipend program!

For more information on available positions contact Kathleen L. Kyer, Manager, National Pediatric Subspecialty Recruitment at: 888-933-1433 or kathleen.kyer@hcahealthcare.com

### Do you Want to Recruit a Pediatric Cardiologist?

Advertise in Congenital Cardiology Today, the only monthly publication totally dedicated to pediatric and congenital cardiology.

For more information send an email to: TCarlsonmd@Gmail.com

### Director of Pediatric Cardiovascular Critical Care

Major Medical Center in Dallas seeks a Pediatric Cardiovascular Critical Care Director. Preferred candidate will possess a charismatic personality, leadership attributes with evidenced experience, strong clinical skill set for a complex patient population and board certification in Pediatric Cardiology and Critical Care. Candidates with board certification in one discipline and solid experience in the alternate subspecialty should also apply. The incoming Director will serve as the Medical Director of the existing 10 bed Pediatric Cardiovascular ICU and the new, state-of-the-art unit due for completion in late 2008. Additional responsibility includes coordinating a collegial collaboration with pediatric cardiology physicians/subspecialists and nursing staff. Incoming physician will be provided an outstanding financial package and the opportunity to advance their medical and/or research career.

The Congenital Heart Surgery program performs more than 300 surgeries each year. Two thirds of the surgeries are pump cases. The program provides care to neonates (approximately 30%) and children under 2 yrs of age (approximately 70%).

A team of nine pediatric intensivists and eleven pediatric cardiologists cover the congenital heart surgery unit. A dedicated 10-bed pediatric cardiovascular intensive care unit opened in 2004. Construction has begun on a newer unit with a completion date in late 2008.

The program participates extensively in research initiatives and i-Rounds, a web based informatics system allowing second to second tracking of clinical data and shares information with outside referring physicians. The center employs all of the latest technologies for monitoring patients and performing point of care testing. The Medical Center operates a very busy research entity in which the Pediatric Cardiovascular Surgery Director, actively participates.

> Call or inquire by email today: Kathleen Kyer, Manager, Pediatric Subspecialty Recruitment, 888-933-1433 or Kathleen.Kyer@HCAHealthcare.com

plishments was doing a front flip, a running front flip, which I thought was pretty interesting. Of course it is fairly rudimentary for real gymnasts, which I never was. This was in high school. Scouting was fun, particularly the camping bit. I love camping and canoeing, canoe tripping. I love anything that has to do with the out of doors. I love skiing. My parents had the strange idea that I shouldn't play football, which I would have loved to do, because they thought I should be careful not to hurt my hands if I was thinking about going into medicine. I used to play the piano but I am rusty now.

### **Q:** What would you discuss at the dinner table?

Everything and anything. We were intensely interested in international affairs. Well, you know that World War II really began in 1939, not in 1941 as many of my friends here think. We were listening to the declaration of war over the radio and my brother lan and I were very excited. We thought it was just wonderful. But I'll never forget looking into my dad's face and he looked so incredibly sad because, as a native Englishman, he understood what this meant, that the Great War was being resumed and that this was a tragedy.

### Q: Would your family all eat together?

We would all eat together and my mom was a wonderful cook. Early on, there were also my grandfather and grandmother Anderson, Alexander and "Teeny" (her name was Clementine and she was quite small, so they called her "Teeny"). We lived in their home in Toronto during the early years of the Depression to save money because there were no jobs. So, those grandparents were very important and our dad's mother, whom we called 'Grandma with the Biscuits'. Her name was Gertrude (my paternal grandmother who made nice biscuits). She was from southern Ireland, from Cork City actually. She was, again, fluent in French, knew Europe well and she was Roman Catholic and we were raised as Protestants. That was, for a period of time, quite a problem because of the Protestant-Catholic divide. You know the story of Northern Ireland.

At that time, it was a very real problem because since my mom and dad were not married in a Roman Catholic Church. We were Episcopalians. For a period of time, Ian and I were regarded by that side of the family as little bastards. Now we may have been, in a certain sense, but not in a religious sense. I am joking of course.

### Q: So what religion did your parents practice?

My mom is a Scottish Canadian who was Presbyterian, which would make her a Protestant, you see. My dad became an Anglican, Episcopalian - Church of England, which was the church

### **ViewFlex** Catheter

EP MedSystems' ViewFlex Catheter is an affordable one-piece, single-use, disposable catheter intended for use within the heart to accurately visualize cardiac structures and blood flow.

**EP**MedSystems

www.epmedsystems.com



that his dad had belonged to, in order to be able to marry my mom. That's why Ian and I were raised as Protestants, which wasn't very popular with the other side of our family. But this was something that we all outgrew and that became less and less and less important as everybody, I think, matured. It is now a complete non-issue, but it was quite real during our early childhood, unfortunately.

### Q: Anything else you care to comment on?

I was the president of our graduating medical year in 1954. This was a huge honor. It was just another fascinating experience. I had to arrange the farewell banquet, you see, and we did it at one of the big downtown hotels. Apparently, every year all kinds of liquor would be stolen by the hotel employees and I listened to these stories and said, "By George, it's not going to happen to us!" I assigned various people to look after our liquor very carefully and sure enough, apparently during the height of the festivities, attempts were made to take whole cases of scotch, etc., and replace them with empties. Our guys immediately jumped on this and that didn't happen.

Sir Robert Watson Watt agreed to be our graduation speaker. He was the inventor of radar. He told us the story of the Battle of Britain, in the summer of 1940. How, with a string of radar stations across the south coast of England, they could spot the Luftwaffe air fleets, as soon as they had cleared the French Coast line and were heading across the channel. Then he told us, the story of the battle of the Atlantic where we used sonar, or ASDIC, which is of course a radar modification, to expose the U-boat packs. As soon as the U-boats became other than invisible, we started to win the Battle of the Atlantic. That made possible the invasion of Europe.

### **Q:** When did you get married?

Nineteen hundred-sixty-two. We had three children in three years because neither of us was terribly young and as pediatricians both, we wanted to avoid nondisjunction and things like that, and we were very pleased. Everybody was beautiful and acyanotic and everything worked out nicely.

### **Q:** What are your children doing now?

Well, our son Andrew is a PhD microbiologist working for a pharmaceutical company here called Cubist, which is named in honor of Picasso. You know, cubism and Picasso were famous for looking at things in a different way and this is what they have been doing. They have a compound called Cubicin, which is the trade name, and daptomycin is the generic name, and it is the only thing that we presently have against otherwise totally resistant Staphylococcus aureus. This is hugely important and if you look at say the Journal of Thoracic and Cardiovascular Surgery, sometimes it comes wrapped in advertisements for this, and Andy is part of the team who is developing that.

Then, our second child, dear Helen, was a teacher of French and a mother of two beautiful boys, and she, alas, died of malignant melanoma in 2001. Our third child, Alexander, is an MIT architect. He graduated from MIT and he is now helping to plan a whole health-care city in Dubai. So everybody is gainfully employed and each of them has two children; so we now have six grandchildren, which means that Granddad has heavy babysitting responsibilities.

### **Q:** Is there something else you would like to do after your medical book?

The most, I think, urgent human problem at the present time is not congenital heart disease. It is mass murder. It is war, particularly, given our thermonuclear toys. Well you see, I developed the segmental approach to the diagnosis of congenital heart disease that is now used worldwide, the method of diagnosis. Now, if we can get a similar understanding, that, all living human beings are in fact members of the same family. We are all brothers and sisters, which the molecular geneticists have proved. If we can taboo murder, mass murder, like we have managed to taboo, oh say, cannibalism, which used to be very common, then we will have a much better chance of surviving to inherit our future. In other words, a modern approach to diagnosis and management in the field of politics and economics that is nonideologic and data based and that taboos mass murder, this is what we need most urgently.

### **Q:** So how would you plan to pursue that?

I wonder, I am open to suggestion. It is an extraordinarily important idea. Given time and strength, perhaps I can write another book but it is very important that all kinds of people get involved in order to avoid the extinction of the human family – and many other life forms. You remember Bernie Lown who did this wonderful thing called "Physicians Against Nuclear War." There was only one thing wrong with that. It has got to be more than physicians. It has got to be politicians, economists; it has got to be people; it has to include all ethnic and religious subgroups.

### **Q:** It strikes me how children in the Middle East are taught so early to hate their enemies.

Trying to distinguish between education and indoctrination is very, very, very important. Coming to understand mythology, religion, and culture is also very, very, very important. Mythology should be understood, not believed. Only when that happens is it likely for us to be able to take the step beyond ideology, to a data-based, evidence-based approach to things. Let people be-







### Opportunity in Pediatric Echocardiography and Non-invasive Imaging

The Heart Center at Nationwide Children's Hospital (NCH, formerly Columbus Children's) the pediatric teaching facility for the Ohio State University, is seeking a board certified/eligible non-invasive imaging pediatric cardiologist to join its' faculty. The NCH Echocardiography Laboratory is ICAEL accredited and fully digital. Our echocardiography laboratory team made up of six attending physicians and nine sonographers performs more than 9000 studies annually using state-of-the-art transthoracic, transesophageal, fetal, intracardiac, intravascular, and 3D techniques. We have an active cardiac MRI/CT service. There are numerous opportunities to engage in translational research, quality assurance initiatives and development of cutting edge informational services. The candidate would participate in programmatic growth encompassing all aspects of The Heart Center's mission including clinical service, education and research. The Heart Center at NCH has excellent programs in adult CHD, cardiac intensive care, electrophysiology, interventional catheterization, thoracic surgery, hybrid strategy and outreach clinics. The fellowship program includes pediatric and combined pediatricadult cardiology and fourth year specialty fellowships.

# Candidates are encouraged to submit their curriculum vitae to:

John Kovalchin, MD, Director of Echocardiography and Non-Invasive Imaging ED628 700 Children's Drive Columbus, OH 43205 PH 614:722-2530 or John.Kovalchin@nationwidechildrens.org

The Ohio State University is an Equal Opportunity, Affirmative Action Employer. Women, minorities, veterans, and Individuals with disabilities are encouraged to apply.





### Opportunity for Adult Congenital Heart Disease Cardiologist

The Heart Center at Nationwide Children's Hospital (NCH, formerly Columbus Children's) and the Ohio State University Medical Center (OSUMC) is seeking a third adult congenital heart disease (ACHD) specialist at the assistant or associate professor level to join our active established program. Candidates with pediatric and/or adult cardiology training who demonstrate proficiency in the medical management of the ACHD patient will be considered. Candidates would be expected to provide consultation to in-patient services at both NCH and the OSUMC campuses as well as to conduct regular ACHD clinics. The ability to perform independent and collaborative laboratory or clinical research in this field is expected. Our ACHD program includes two dually trained & board certified pediatric and adult cardiology physicians, ACHD nurse practitioners, nurse clinicians, research nurse and dedicated technical staff. Additionally, we have an accredited ACHD five-year fellowship program currently with three fellows in training. Our program actively manages nearly 2000 ACHD patients and has an emerging transitional ACHD program. State of the art imaging, interventional catheterization & surgery, electrophysiology, intensive care, CHF and heart, heart-lung and lung transplant programs are integrated with the ACHD program.

### Candidates may submit their curriculum vitae to:

Curt Daniels, MD, Nationwide Children's Hospital 700 Children's Drive Columbus, Ohio 43205 or via e-mail to curt.daniels@nationwidechildrens.org

The Ohio State University is an Equal Opportunity, Affirmative Action Employer. Women, minorities, veterans, and individuals with disabilities are encouraged to apply.



### Watch LIVE CASES from Congenital Cardiology Today

www.CHDVideo.com Presented courtesy of:

The Pediatric and Adult Interventional Therapies for Congenital and Valvular Disease (PICS)

and The 6th International Workshop on Interventional Pediatric Cardiology (Workshop IPC)

Watch a Percutaneous Valve Implantation, a Perimembranous VSD Closure or both

www.CongenitalCardiologyToday.com

lieve whatever they want in private but they shouldn't force it on their neighbors, or kill them if they don't accept their little stories, their beliefs, whatever they happen to be. You know, this is, again, the separation of church and state idea, but with the understanding that, the whole idea of right versus left is absurd. It should be right versus wrong. Correct versus incorrect. Can you imagine if the politicians of the world were to approach things like, say, cardiologists do. And we'll have a meeting in six months or a year to see how this or that is working out. How can we do this better?

In other words, the medical model of how major human problems can be diagnosed and managed is both scientific and humanistic. We have to understand other peoples' cultures, religions, fears, and biases. Science won't do it alone. This is why we need a medical approach. You see, a biomedical approach is much more than science. There is a lot of art and humanism involved.

### **Q:** What books are on your night-stand?

"The Journey of Man" by Spencer Wells. Then there is another one called "Before the Dawn", by Nicholas Wade which is, again, the early history of homo sapiens. In my bag is "The Story of Civilization" by Will Durant, who wrote an 11 volume account of The Story of Civilization. I finished reading recently a book called "Parallel Worlds, A Journey Through Creation Higher Dimensions, and The Future of the Cosmos" by Michio Kaku. It's enthralling, particularly if you like Physics (which I do). So, I hope you see a pattern here. These interests are both scientific and humanistic - not either one alone. At its best, this is what modern medicine is: both scientific and humanistic.

We physicians have a huge and very important "SECRET." By trial and error, we have learned how to understand and manage successfully many important potentially fatal human problems. Once understood, the "medical model" is very widely applicable to major human problems.

# **Q:** And you are working at the Registry now?

No, I am working at 21 Autumn Street. That is where they send retired gentlemen, Autumn Street. I am wondering where Winter Street is. Actually, bad jokes aside, the two buildings on Autumn Street, 1 and 21, which used to be Deaconess buildings, have been purchased by Children's Hospital and the floor I am on now, the fourth floor, is entirely cardiology, which is nice. But unfortunately, I am separated from the lab. But that is appropriate because my successor is there, you see.

# **Q:** How has life been for you without Stella?

Very, very different. My cooking, by comparison with my bride's culinary savoir faire, can only be described as embryonic. It is scarcely fetal. But I manage. I make my own lunches and I make my own dinners. I am not as good at dusting as I should be. Dear Stelly, you know they are dedicating this whole meeting in Washington to her. I have to give a memorial lecture in her honor, which is also not going to be easy for me. I often ask myself, what would Stella say or think about this? Or that? Because we discussed everything of course.

# **Q:** So do you have any advice for young pediatric cardiologists or young surgeons?

I think that one should get as well-trained as possible, and that one should adhere to the truth. For us, this is the only thing that matters. Everything else is just conversation. Getting well-trained at the best places by the best people and trying to do, literally, the very best that one possibly can in all situations. It is the old Boy Scout motto, "Do your best." It really is, and it comes to that. Not just in every case, in your training, in each and every paper.

## **Q:** What is your most important paper?

The paper that we published in 1972 about the segmental approach to diagnosis, many people tell me they think that is our most important paper because it turned out to be the diagnostic approach that people use worldwide, with or without minor modifications.

All of the others are as good as we could do too. So I would say to people, just do the very best that you possibly can. In other words, for me, science is not an occupation. It is a way of life. It is a way of looking at everything. It is a way of thought. It is a way of living. What a privilege it is to be a physician.

ССТ



Bradley W. Robinson, MD Pediatric Cardiologist Department of Pediatrics Division of Pediatric Cardiology Nemours Cardiac Center - AIDHC 1600 Rockland Road Wilmington, DE 19803-3607 USA Phone: (302) 651-6600 Fax: (302) 651-5345

bwrobins@NEMOURS.org



### Introducing the iE33 intelligent echo system ... for better visualization of congenital heart defects

Learn more at: www.medical.philips.com/iE33 or call 800-229-6417



The Foundation for The Gator Nation

# Congenital Heart Center at the University of Florida

The Congenital Heart Center at the University of Florida, is recruiting a Board Certified Pediatric Cardiologist for the faculty position of Clinical Assistant Professor with experience in echocardiography. This faculty position will be focused in the area of congenital heart disease serving as a general pediatric cardiologist and being primarily responsible for performing echocardiography services. This role includes opportunities to participate in research and teaching efforts of residents, fellows, medical students and other health care professionals.

The appointment will be non-tenure-accruing. This position will remain open until an appropriate candidate is selected

Applicants should send a letter of application, a C.V., and three letters of recommendation referencing LP#00017673 to:

> Barry J. Byrne, M.D., Ph.D. The Congenital Heart Center University of Florida College of Medicine P.O. Box 100296 Gainesville, FL 32610-0296

The University of Florida is an Equal Opportunity Institution.



20

### Faculty Member for the Congenital Heart Center at the University of Florida

The Congenital Heart Center at the University of Florida is recruiting a Board Certified Pediatric Cardiologist with fourth year interventional catheterization training for faculty position of Pediatric Interventional Cardiologist. This position will assist with coordinating all aspects of pediatric interventional cardiac catheterization services and provide general pediatric cardiology care. This role includes teaching of residents, fellows, medical students and other health care professionals and participation in a strong clinical research program and excellent clinical practice.

The appointment will be at the nontenure accruing level of Clinical Assistant/Associate Professor based upon experience. This position will remain open until an appropriate candidate is selected.

Applicants should send letter of application, C.V., and three letters of recommendation referencing LP# 00023005 to:

Randal M. Bryant, M.D. Search Committee Chair Congenital Heart Center University of Florida College of Medicine P.O. Box 100296 Gainesville, FL 32610-0296

The University of Florida is an Equal Opportunity Institution.



### Faculty Member for the Congenital Heart Center at the University of Florida

The Congenital Heart Center at the University of Florida is recruiting a Board Certified Pediatric Cardiologist for the faculty position of Director, Non-invasive Imaging. This position will lead a cohesive unit for advancing research and clinical care in congenital heart disease imaging through coordination of echocardiography, MRI and CT angiography. The position will also provide general pediatric cardiology services and includes teaching residents, fellows, medical students and other health care professionals.

The appointment will be at the nontenure or tenure accruing level of Associate Professor or Professor based upon experience. This position will remain open until an appropriate candidate is selected.

Applicants should send letter of application, C.V., and three letters of recommendation referencing LP# 00023002 to:

Barry J. Byrne, M.D., Ph.D. Medical Director Congenital Heart Center University of Florida College of Medicine P.O. Box 100296 Gainesville, FL 32610-0296

The University of Florida is an Equal Opportunity Institution.

### CSI 2008 Congenital and Structural Interventions Frankfurt, Germany June 25-28 2008

Congenital and Structural Interventions with live case demonstrations from Frankfurt and Hands on Workshops. For more information: http://www.csi-congress.org/



Structural Interventions

### New Products, News and Information

#### Adding Rapid Response Medical Team in a Children's Hospital Helps Reduce Risk of Death, Rates of Cardiac and Respiratory Arrest

CHICAGO – A children's hospital that added a rapid response medical team for patients not in the intensive care unit saw an 18% decrease in the death rate, and about a 70% decline in the rate of cardiac and respiratory arrests, according to a study in the November 21, 2007 issue of JAMA.

Introduction of a rapid response team (RRT; medical emergency team) has been shown to decrease death and cardiopulmonary arrests outside of the intensive care unit (ICU) in adult inpatients, according to background information in the article. An RRT is a multidisciplinary team frequently consisting of ICU-trained personnel who are available 24 hours per day, 7 days per week for evaluation of patients not in the ICU who develop signs or symptoms of clinical deterioration.

"The RRT intervention was developed in response to research that revealed adult patients on general medical and surgical hospital units often have evidence of physiological deterioration several hours before cardiopulmonary arrest, and that after a cardiac arrest occurred, survival rates were poor. Given that there appears to be a window of opportunity to identify and proactively treat 'pre-arrest' adult inpatients effectively, the Institute for Healthcare Improvement recommended RRTs be implemented nationwide in an effort to decrease inpatient mortality rates," the authors write. Limited data exist evaluating the effectiveness of RRT implementation in pediatric inpatients.

Paul J. Sharek, MD, MPwww.H, of Stanford University School of Medicine, Palo Alto, Calif., and colleagues evaluated whether RRT implementation was associated with decreases in hospitalwide mortality rates and code rates (respiratory and cardiopulmonary arrests) outside of the ICU in pediatric inpatients at a 264bed academic children's hospital. Pediatric inpatients who spent at least one day on a medical or surgical ward between January 2001 and March 2007 were included. A total of 22,037 patient admissions were evaluated pre-intervention (before September 1, 2005), and 7,257 patient admissions were evaluated postintervention (on or after September 1, 2005).

A significant decrease of 18% occurred in the hospital-wide mortality rate after implementation of the RRT. The rate of codes outside of the ICU setting per 1,000 eligible admissions declined by 71.7%, with pre-intervention and post-intervention rates of 2.45 vs. 0.69, respectively. The rate of codes outside of the ICU per 1,000 eligible patient-days decreased by 71.2% after RRT implementation.

The RRT intervention, using statistical modeling, was associated with a decrease of 0.178 deaths per 100 discharges or 1.78 deaths per 1,000 discharges. During the 19-month post-intervention period, the RRT intervention is estimated to have resulted in 33 lives saved at this hospital.

"Implementation of an RRT in our free-standing, quaternary care academic children's hospital was associated with statistically significant reductions in hospital-wide mortality rates and code rates outside of the ICU setting. These reductions cannot be explained by differences in patient characteristics or severity of illness between the control and post-intervention populations," the authors write. For more information, see (JAMA. 2007;298(19):2267-2274.

#### The Heart Center at Nationwide Children's Hospital Opens First Hybrid Congenital Cardiac Operating Suite in the US



The innovative spirit of the cardiothoracic surgery and interventional cardiology teams in The Heart Center at Nationwide Children's Hospital have come together to bring technology first developed in its Hybrid Catheterization Suites to a state-of-the-art operating suite - the first Hybrid Congenital Cardiac Operating Suite in the U.S.

This operating suite is designed specifically around the cardiac surgeon and cardiac surgical patient, offering new advanced access and imaging in an operating room environment. The suite accommodates the collective catheterization and surgery team, and facilitates rapid information sharing with new technology from TOSHIBA Medical Systems and Maquet Surgical Workplaces.

The Hybrid Congenital Cardiac Operating Suite features:

- More than 900 square feet of working space to accommodate the team involved in a hybrid case (surgeons, interventional cardiologists, electrophysiologists, echocardiographers, cardiac anesthesiologists, advanced level nurse practitioners, physicians' assistants, cath/O.R. nurses, technicians and perfusionists).
- A surgical theater with a fixed base surgical table, modutec booms and surgical lighting from Maquet Surgical Workplaces, allowing for more precise and consistent imaging, as well as greater flexibility without taking up critical floor space.
- Flat Panel Detectors (FPD) from TOSHIBA America Medical Systems for a better quality picture which facilitates more accurate assessment and diagnosis.



University Hospitals Case Medical Center



22

### Rainbow Babies and Children's Hospital / University Hospitals/Case Medical Center

The Division of Pediatric Cardiology seeks interested faculty candidates for:

Pediatric Electrophysiology – responsibilities would include directing and developing the Pediatric Electrophysiology Program at UHCMC / Rainbow Babies and Children's Hospital. This person would be expected to foster and develop an invasive EP program in the northeast Ohio region. Clinical and academic collaboration with two busy affiliated adult EP divisions is available and encouraged.

Non-invasive Cardiology / ECHO Lab Director – cardiologist with imaging interest and expertise to help develop comprehensive imaging services to include fetal, MRI, TEE, 3-D, and regional telemedicine program. Would join three other full time non-invasive cardiologists in providing clinical services at Rainbow Babies and affiliated institutions.

Critical Care Cardiology / Director – responsibilities will include establishing, developing, and directing cardiovascular critical care services in collaboration with the Divisions of Cardiothoracic Surgery / Cardiology and Pediatric Intensive Care.

The Pediatric Cardiovascular Program at UHCMC/ Rainbow's is provides full-service congenital cardiovascular care to the Northeast Ohio region, and currently consists of 6 full time cardiologists, 2 cardiothoracic surgeons, 3 certified NPs, 2 full-time investigators and comprehensive subspecialty and fellowship programs. We are academically affiliated with Case Western Reserve University, and interest in clinical and/or basic research is encouraged. Rainbow Babies and Children's Hospital is consistently ranked among the top children's hospitals nationally.

Interested applicants should forward cover letter and CV to:

Ernest S. Siwik, MD Interim Director, Pediatric Cardiology RBC 380C – 11100 Euclid Avenue, Cleveland, OH 44106 ernest.siwik@uhhospitals.org

"In Employment, as in education, UHCMC/CWRU are committed to Equal Opportunity and world class diversity. Applications from qualified women and minorities are encouraged."

- Technical capabilities that allow staff members the ability to transmit information anywhere in the suite, at remote sites within the hospital, or even to countries around the world, enabling Nationwide Children's Heart Center faculty to consult with physicians in distant countries and to serve as a teaching hospital.
- Nine interconnected, flat panel, high-resolution display screens for superior visualization and better information exchange among the team members; at least two monitors are visible to all team members at any time, from any location.
- A non-sterile control/observation room to serve as an educational and training resource for students, staff and faculty.

The Hybrid Congenital Cardiac Operating Suite can accommodate any cardiac surgical case, catheterization or collaborative hybrid procedure. Cardiothoracic surgeons and interventional cardiologists working together in this suite will reduce the amount of time required to correct a heart problem and the amount of emotional and physical stress placed on a patient or their family which translates into less pain, less scarring and a faster recovery time for patients.

The Nationwide Children's Hospital Heart Center was established in 2002 when all Cardiology and Cardiothoracic Surgery services were consolidated into a comprehensive "one-stop" model of patient/family convenience and excellence in patient care, education and research. The Heart Center opened the nation's first Hybrid Cardiac Catheterization Suites in 2004 in an innovative new approach to cardiac care, whereby the benefits of cardiothoracic surgery and interventional cardiology were combined in two suites to help critically ill children and adults with congenital heart disease achieve improved therapeutic outcomes. These facilities harness state-of-the-art technology that allows surgeons and interventionalists to collaboratively perform a wide range of innovative hybrid procedures.

The new Hybrid Congenital Cardiac Operating Suite and the Hybrid Cardiac Catheterization Suites deviate from traditional cardiac care delivery by allowing surgical and interventional catheterization procedures to be performed by a single interdisciplinary team in one setting. This approach supports enhanced patient outcomes through shorter and safer procedures that require less recovery time and rehabilitation. The expansive suites accommodate the collective catheterization and surgical teams, and facilitate rapid information.

The first procedure performed in the Hybrid Congenital Cardiac Operating Suite occurred in November 2007 and the first procedure in the Hybrid Cardiac Catheterization Suites took place in May 2004. Both procedures were conducted to treat Hypoplastic Left Heart Syndrome. For more information, visit: www.NationwideChildrens.org/heartcenter.

### <u>PFO Detection</u> <sup>with</sup> Power M-Mode Doppler

Higher Sensitivity than TEE\*
 Non-invasive method enables a measured
 valsalva strain
 Easy to perform, office-based testing
 'I of Neuroimania Vol 14 No 4 Ot 2004





The Division of Pediatric Cardiology at the University of Utah School of Medicine and Primary Children's Medical Center is recruiting 3 Pediatric Cardiologists with expertise in the following areas to join a 20-member division. All candidates should have a strong clinical background in all areas of pediatric cardiology, must be BE/ BC in pediatric cardiology, and eligible for wwwww in the State of Utah.

**Noninvasive Imaging:** The candidate should have expertise in echocardiography (TTE and TEE). The candidate should also have expertise in fetal echo and/or cardiac MRI.

**Transplant/Heart Failure:** This candidate should have expertise in caring for children with heart failure and those awaiting and undergoing transplantation.

**Electrophysiology:** The candidate should have expertise in pediatric EP, including medical management of inpatients and outpatients with rhythm abnormalities, invasive EP studies, pacemakers, and some exposure to implanted EP devices.

The successful candidates will receive faculty appointments at the University of Utah. These positions may be at the rank of Instructor through Associate Professor. For all positions, there will be protected time for clinical research with mentoring available within the Division. The Division has a very active clinical research program and is one of the participating centers in the Pediatric Heart Network funded by the NIH.

The Division is based at Primary Children's Medical Center, a tertiary referral center for a 3-state area located on the hills overlooking Salt Lake City. The area offers an excellent quality of life with immense cultural and recreational opportunities close and available. The University of Utah is an Equal Opportunity Employer and welcomes applications from minorities and women and provides reasonable accommodations to the known disabilities of applicants and employees.

### Interested individuals should contact:

Lloyd Y. Tani, M.D. Chief, Division of Pediatric Cardiology Professor of Pediatrics University of Utah School of Medicine Email: Iloyd.tani@intermountainmail.org Phone: (801) 662-5400



Would you like to get your issue of Congenital Cardiology Today in a PDF file?

If, so, send an email to: PDF@CCT.bz

### **CONGENITAL CARDIOLOGY TODAY**

© 2007 by Congenital Cardiology Today ISSN: 1544-7787 (print); 1544-0499 (online). © 2007 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499online). Published monthly. All rights reserved.

#### Headquarters

9008 Copenhaver Dr. Ste. M Potomac, MD 20854 USA

#### Publishing Management

Tony Carlson, Founder & Editor TCarlsonmd@mac.com Richard Koulbanis, Publisher & Editor-in-Chief RichardK@CCT.bz John W. Moore, MD, MPH, Medical Editor/ Editorial Board JMoore@RCHSD.org Jeffrey Green, Contributing Editor

#### **Editorial Board**

Teiji Akagi, MD Zohair Al Halees. MD Mazeni Alwi, MD Felix Berger, MD Fadi Bitar. MD Jacek Bialkowski, MD Philipp Bonhoeffer, MD Mario Carminati, MD Anthony C. Chang, MD, MBA John P. Cheatham, MD Bharat Dalvi, MD, MBBS, DM Horacio Faella, MD Yun-Ching Fu, MD Felipe Heusser, MD Ziyad M. Hijazi, MD, MPH Ralf Holzer, MD Marshall Jacobs, MD R. Krishna Kumar, MD, DM, MBBS Gerald Ross Marx, MD Tarek S. Momenah, MBBS, DCH Toshio Nakanishi, MD, PhD Carlos A. C. Pedra, MD Daniel Penny, MD James C. Perry, MD P. Syamasundar Rao, MD Shakeel A. Qureshi, MD Andrew Redington, MD Carlos E. Ruiz, MD, PhD Girish S. Shirali. MD Horst Sievert, MD Hideshi Tomita, MD Gil Wernovsky, MD Zhuoming Xu, MD, PhD William C. L. Yip, MD Carlos Zabal, MD

#### FREE Subscription

Congenital Cardiology Today is available free to qualified professionals worldwide in pediatric and congenital cardiology. International editions available in electronic PDF file only; North American edition available in print. Send an email to Subs@CCT.bz. Include your name, title, organization, address, phone and email.

#### **Contacts and Other Information**

For detailed information on author submission, sponsorships, editorial, production and sales contact, current and back issues, see website or send an email to: INFO@CCT.bz.

# **REVISED DATE**

# PICS-AICS 08 July 20-23, 2008 Bellagio, Las Vegas

PEDIATRIC AND ADULT INTERVENTIONAL CARDIAC SYMPOSIUM



Course Directors: Dr. Ziyad M. Hijazi, Dr. William E. Hellenbrand, Dr. John P. Cheatham, & Dr. Carlos Pedra

- FOCUSING ON THE LATEST ADVANCES IN INTERVENTIONAL THERAPIES FOR CHILDREN AND ADULTS with congenital and structural heart disease, including the latest technologies in devices, implantable valves, stents and balloons. Special sessions will provide an in-depth focus on septal defect closure, coarctation stenting, hybrid intervention for HLHS and fetal intervention.
- SPECIAL SESSIONS will be dedicated to the care of adults with congenital heart disease.
- HOT DAILY DEBATES between cardiologists and surgeons on controversial issues in intervention for congenital and structural heart disease.
- The popular session of "MY NIGHTMARE CASE IN THE CATH LAB"

- LIVE CASE DEMONSTRATIONS featuring approved and non-approved devices will be transmitted daily from many cardiac centers around the world. During these live cases, the attendees will have the opportunity to interact directly with the operators to discuss the management options for these cases.
- BREAKOUT SESSIONS for cardiovascular nurses and CV technicians.
- MEET THE EXPERT SESSION, held on Sunday, will give attendees the opportunity to discuss difficult cases with our renowned faculty.

**ACCREDITATION** The Society for Cardiovascular Angiography and Interventions is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to sponsor continuing medical education for physicians.

Abstract Submission Deadline is March 15, 2008. For registration and abstract submission go online to **www.picsymposium.com** 



### **O** RUSH UNIVERSITY MEDICAL CENTER