News and Information for Pediatric and Congenital Cardiovascular Physicians and Surgeons

Vol. 4 / Issue 11 November 2006 North American Edition

INSIDE THIS ISSUE

Tetralogy of Fallot (TOF) With Anomalous Left Coronary Artery From Pulmonary Artery (ALCAPA) as a Rare Presentation of Neonatal Heart Failure by Mashail AL-Obaidan, MD; Zuhair AL-Halees, MD, RCSC, FACS, FACC, FACCP ~Page 1

Highlights from PICS 2006, Las Vegas, Nevada by Ziyad M. Hijazi, MD ~Page 10

Berry Syndrome in a Neonate: Report of a Case Successfully Repaired in a One-Stage Operation by Marci Marcello, MD; Calvaruso Davide, MD; Cipriani Adriano, MD; Ocello Salvatore, MD; Salviato Nicoletta, MD; and Marcelletti F. Carlo, MD ~Page 16

DEPARTMENTS

Ziyad M. Hijazi Receives the PICS Founder Award ~Page 14

Medical News, Products and Information ~Page 15

Medical Conferences ~Page 18

CONGENITAL CARDIOLOGY TODAY 9008 Copenhaver Drive, Ste. M Potomac, MD 20854 USA Tel:+1.301.279.2005 Fax: +1.240.465.0692

www.CongenitalCardiologyToday.com

© 2006 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499online). Published monthly. All rights reserved. Congenital Cardiology provides timely news and information for pediatric and congenital cardiologists. Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors, and are not necessarily the views of Congenital Cardiology Today.

Recruitment ads See pages 11, 13, 15, 19 WWW.CONGENITALCARDIOLOGYTODAY.COM

TETRALOGY OF FALLOT (TOF) WITH ANOMALOUS LEFT CORONARY ARTERY FROM PULMONARY ARTERY (ALCAPA) AS A RARE PRESENTATION OF NEONATAL HEART FAILURE

By Mashail AL-Obaidan, MD; Zuhair AL-Halees, MD, RCSC, FACS, FACC, FACCP

Abstract

Coronary artery anomalies are common among patients with tetralogy of Fallot, however, the origin of the left coronary artery from the main pulmonary artery in a patient with TOF is rarely seen and not yet reported in the literature. The patient reported here presented at three weeks of age with impaired cardiac function and was found to have this rare anomaly.

Introduction

Tetralogy of Fallot is considered one of the most common cyanotic congenital heart defects. It is characterized by VSD, hypoplastic stenotic pulmonary valve and main pulmonary artery, hypoplastic outlet septum (infundibulum), and overriding of the aorta.

Associated coronary abnormalities are quite common, including RCA from LCA, LAD from RCA with LAD crossing the RV outflow tract, and single coronary ostium. There is also a single case report of RCA arising from the MPA in TOF associated with Pulmonary Atresia. The patient in this case report had TOF with ALCAPA. This specific combination of abnormalities has not to our knowledge, been previously reported. We report the initial presentation and management of the patient, and review the literature.

Case report

A 1.5 month old boy weighing 3.2 Kg was referred from the periphery with congestive heart failure. He had two previous admissions due to chest infection and cardiomegally. There was a history of low grade fever, cough and shortness of breath ten days prior to the first admission associated with blue discoloration of the lips. After complete blood work and screening for sepsis, he was treated for seven days with antibiotics and was started on anti-failure medications (digoxin, lasix, and captopril). He was referred to our institute because of a heart murmur to rule out congenital heart disease.

On examination, he had no dysmorphic features. He was pale and tachypneic, had nasal flaring, and his saturation in room air was 84%. His lung examination was not remarkable, but his abdominal examination revealed a liver edge 4 centimeters below the costal margin. His cardiovascular examination showed normal S1, single S2, and a

AMPLATZER® "Cribriform" Septal Occluder

Now Available in the U.S.



A Single Solution for Multi-Fenestrated Atrial Septal Defects

Multi-fenestrated Atrial Septal Defects require an occlusion device with a narrow waist for atraumatic placement in the central defect, and broad contact surface to occlude the peripheral "windows".

The AMPLATZER® Cribriform Septal Occluder completes AGA Medical Corporation's line of ASD Occlusion Systems, to provide a transcatheter closure solution for the majority of atrial septal defects.

Come visit us at: AHA Booth 2247



Multi-fenestrated atrial septal defects represented by a multitude of small holes in the atrial septum. This is also referred to as a "Swiss cheese" ASD.



The AMPLATZER Cribriform Occluder enables occlusion of the defect by covering the holes with a single device.



For More Information, Visit: www.amplatzer.com

888-546-4407 (toll free, US) 763-513-9227

W711: Turbre to Indicadorans for Tube for a complete follow of indicatoria and image, continuencedores, scienceps and prevaillosis, Gorda E AGA Mindicad Corporation for Turbice information.

Bonly C€....

O AVA Modual Conjunctions 2008. ABPL/VEDE is a singulared transmark of AVA Modual Conjunctions. Not in any way commotivel with modual gam in suppresent solid under the "AVA". Second by AVA-301 or the secondary.

\$88841171 (HS



Figure 1. ECG of the patient showing deep Q wave, ST elevation and RVH.



Figure 2. Chest X-ray of the patient which revealed cardiomegally and oligaemic lung fields.

systolic ejection murmur most prominent at the left uppersternal border. The remainder of his examination was normal for his age.

Laboratory investigation showed: normal CBC with Hemoglobin of 14 gm/dl. Serum electrolytes and liver function tests were within normal limits. Urine analysis was negative. Blood and urine cultures were negative. Serum virology studies were also negative. Serum carnitine was normal. LDH was 564 (elevated).

ECG: S/R, RAD, RVH, deep Q wave especially in lead V6, ST elevation (Figure 1).

Chest X-ray: cardiomegally, oligaemic lung field (Figure 2).

Echocardiography: Showed TOF with malaligned VSD, overriding aorta, infundibular and pulmonary valve stenosis with maximum velocity of 4.4 m/s (peak gradient 76 mm Hg), dilated RCA, myocardial coronary collaterals, LCA from MPA, markedly impaired left ventricular systolic function (ejection fraction of 15%, shortening fraction < 10%), PFO, mild mitral valve regurgitation (Figures 3-6).

Cardiac Catheterization: Proved the presence of TOF and ALCAPA (Figure 7-9).

Hospital Course

After completing these investigations and confirming the diagnosis, the patient was scheduled for surgical repair. Surgery included repair of TOF using the transatrial approach with VSD closure and resection of the infundibular stenosis and pulmonary valvotomy. Direct reimplantation of the LCA to the ascending aorta was also performed. Cross-clamp time was 90 minutes,





Working Together to Develop a Better Tomorrow



Figure 3. Echocardiography of the patient showed malalinged VSD & overriding of aorta.



Figure 4. M-mode revealed impaired LV systolic function.

and bypass time was 140 minutes. The patient's chest was left opened for 36 hours, and the patient required standard inotropes (dopamine, dobutamine, and milrinone). The patient remained in sinus rhythm, and his ECG normalized on the 3rd post-operative day. He required ventilation for 6 days, and stayed for an additional 11 days in the hospital. Echocardiography performed seven days after surgery revealed no residual VSD and no evidence of residual pulmonary stenosis. Left ventricular function was improved with ejection fraction of 30%.

Follow Up

Three months after hospital discharge, the patient was thriving and had no cyanosis. He was taking anti-failure medications. Echocardiography revealed further improvement in left ventricular function with ejection fraction of 45%.

Discussion

Coronary artery anomalies are often associated with tetralogy of Fallot. The left anterior descending coronary artery arising from the right coronary artery and crossing the RVOT is the commonest of them with reported incidence of 2-9% (1,2,3).

Additional anomalies include single coronary ostium, origin of the RCA from the LCA (4), a significant conal branch from the RCA (or a dual LAD), paired anterior descending arteries originating from left coronary artery and right coronary artery as well as RCA from LAD (5).



10th Annual Update on Pediatric Cardiovascular Disease New and Evolving Practices February 21-25, 2007 Disney's Yacht & Beach Club Resort[®] Lake Buena Vista, Florida USA

www.chop.edu/cardiology2007



Figure 5. parasternal short axis view showing malalinged VSD with infundibular and pulmonic valve stenosis v max @ 4.5 m/s.



Figure 6. 2-D and color-Doppler showing pulmonic valve stenosis with LCA arising from the pulmonary artery.

Anomalous origin of the LCA from the pulmonary trunk (ALCAPA) is a well described, although rare, congenital anomaly in humans, with an incidence as an isolated anomaly of about one case for each 300,000 live birth. It represents 0.5% of the total cases of congenital heart disease. This condition is often characterized by a single origin of the whole left coronary artery

from the posterolateral wall of the pulmonary trunk. The majority of patients present in infancy after 2 months of age (the time of normalizing pulmonary vascular resistance) with a picture of congestive heart failure and angina. Almost all die in the first year of life if they are not treated (6,7). A few reports describe older individuals with the disease (7).



PediHeartNet is a free internet-based discussion group exclusively for health care professionals interested in Pediatric Cardiology and Cardiac Surgery.

For information or to sign up please visit:

www.Pediheart.net



www.CongenitalCardiologyToday.com



Figure 7. Aortogram (RAO and LAO) revealed RCA filling from aorta & LCA filling from RCA through collaterals and Rt aortic arch (arrow indicate LCA).





Figure 8. Aortogram revealed dilated RCA.

Figure 9. Selective RCA angiogram revealed filling of LCA from collaterals arising from the RCA.

The association of tetralogy of Fallot and anomalous left coronary artery from the main pulmonary artery has not been previously reported. There are scattered reports of TOF with RCA origin from the MPA (8), and TOF and PA with MPA arising from the RCA or LCA wherein the coronary arteries arise from the aorta and are the sole blood supply to the pulmonary artery (9,10). It is also worth mentioning that there are a few reported cases of coronary to pulmonary artery collaterals in which the coronary arteries arise normally from the aorta in patients with TOF and PA (11).

The clinical presentation of our patients was earlier than the typical presentation of patients with isolated ALCAPA. Our

patient presented in the neonatal period with congestive heart failure and markedly impaired left ventricular function.

The earlier presentation of congestive heart failure in patients with TOF and ALCAPA may be explained by lower pulmonary vascular resistance in the neonate born with pulmonary stenosis and infundibular stenosis. In contrast patients with isolated ALCAPA may have a more delayed fall in pulmonary vascular resistance. This delays the steal phenomenon, the ischemic insult to the left ventricle and the development of congestive heart failure.

Our patient underwent successful total repair of tetralogy of Fallot using a transatrial approach with VSD closure and resec-

-SOMANETICS

REVEAL TISSUE PERFUSION

WITH SITE-SPECIFIC SATURATION DATA



Augment *systemic* data with noninvasive *regional* oxygen saturation (rSO₂) from up to four sites. Only the INVOS[®] Cerebral/Somatic Oximeter monitors oxygenation data from the brain and body simultaneously and continuously. This site-specific tissue perfusion can enhance your clinical assessment in a meaningful way and help detect ischemic problems earlier than traditional measures. This additional vital sign lets you intervene and put things right — before they escalate.

800-359-7662 www.somanetics.com REFLECTING THE COLOR OF LIFE"





© 2006 Somanetics Corporation. Somanetics and INVOS are registered trademarks of Somanetics Corporation. "Reflecting the color of life" is a trademark of Somanetics Corporation. US federal regulations restrict the sale of this device to, or on the order of, licensed medical practitioners.

NOVEMBER 2006

8

"Coronary artery anomalies are common among patients with tetralogy of Fallot, however, the origin of the left coronary artery from the main pulmonary artery in a patient with TOF is rarely seen and not yet reported in the literature."

tion of the infundibular stenosis and pulmonary valvotomy, as well as direct reimplantation of the anomalous left coronary artery to the ascending aorta (12).

The patient had a favorable immediate and early result as well as in the first follow up period with improvement of the ventricular function.

From this case, we learned that any type of coronary artery anomaly can present in conjunction with tetralogy of Fallot, including ALCAPA. Furthermore the combination of TOF and AL-CAPA may result in an early and devastating patient presentation, requiring prompt diagnosis and treatment.

Acknowledgement

To Mrs. Farida AL-Hazmi the senior echocardiographer for her grateful help in retrieving the best echo, images.

References

1. The Science and Practice of Pediatric Cardiology. 2nd edition by Garson, Bricker, Fisher, Neish 1998.

2. Echocardiography in Pediatric Heart Disease 2nd edition by Snider, Serwerard Ritter 1997.

3. Husain AK, Khadeparkar JM, Tendolkar AG, Mulay AV, Surgical Management of Tetralogy of Fallot with Anomalous Coronary Artery, Journal of Postgraduate Medicine JPGM 1991; 37 (3):148-51.

4. Hekmat M, Rafeyian S, Detal M, Associated Coronary Anomalies in 135 Iranian Patients with Tetralogy of Fallot. Asian Cardiovascular & Thoracic Annals 2005; 13:307-310.

5. Ruzmetov M, Jimenez MA, Pruitt A, et al: Repair of Tetralogy of Fallot with Anomalous coronary arteries Coursing Across the Obstructed Right ventricular Outflow Tract, Pediatric Cardiology 2005; 26(5):537-542.

6. Amaral F, Carvalho JS, Joao A et al, Anomalous Origin of the Left Coronary Artery from the Pulmonary Trunk. Clinical Features and Midterm Results after Surgical Treatment, Arq. Bras. Cardiol. 1999; 72(3):766-82.

7. Alabdulgader AA, Bulbol ZR, Anomalous Left Coronary Artery Arising from the Pulmonary Artery in an Adolescent, Normalization of the LV Dimension and Function Following Repair 2001; 21(3):204-205.

8. Moss RL, Backer CL, Zales VR, et al, Teralogy of Fallot with Anomalous Origin of the Right Coronary Artery, Ann Thorac Surg 1995; 59:229-231.

9. Wu QY, Yang XB, Anomalous Origin of the Pulmonary Artery from the Right Coronary Artery, Ann Thorac Surg 2001; 72:1396-1398.

10. McMahon CJ, Nihill MR, Origin of the Main Pulmonary Artery from the Left Coronary Artery in Complex Pulmonary Atresia, Pediatric Cardiology 2001; 22(4):347-349.

11. Amin Z, McElhinney DB, Reddy VM, et al: Coronary to Pulmonary Artery Collaterals in Patients with Pulmonary Atresia and Ventricular Septal Defect, Ann Thorac Surg 2000; 70:119-123.

12. Cardiac Surgery 2nd edition by John W. Kirklin, Brian G. Barratt-Boyes 1996.

~CCT~

Corresponding Author:

Mashail AL-Obaidan, MD Consultant Pediatric Cardiology Paediatric Cardiology Dept. PSCC PO BOX 999111 Riyadh 11625 KSA Fax/Tel: 9661473-0049 Tel: 4777714 EXT: 8795, Pager: 4110

Email: mashaila@hotmail.com

Zuhair AL-Halees, MBBS, MD, FRCSC, FACS, FACC, FACCP Senior surgeon KFSH & RC





Meet the second-smallest thing to enter your ICU.

MICROMAX



In the crowded pediatric ICU, where every second counts,

the MicroMaxx[®] ultrasound system from SonoSite gives you the power of your echo lab in an 8-pound, hand-carried package for immediate bedside analysis. Our P10/8-4 transducer is ideal for pediatric cardiology, and a wide range of built-in cardiac calculations plus high image quality helps you make effective, efficient diagnoses. The future of portable echo is here. Just in time to improve the futures of even your tiniest patients.



www.sonosite.com

HIGHLIGHTS FROM PICS 2006, LAS VEGAS, NEVADA

By Ziyad M. Hijazi, MD

With just about 1000 attendees from 60 countries in addition to the USA, PICS (Pediatric and Adult Interventional Therapies for Congenital and Valvular Heart Disease) was a huge success. 60% of the attendees were from the US, and 40% from abroad. I was told many of our international colleagues could not get a visa to the US!

For the first time, this year's meeting included combined and breakout pediatric and adult sessions for interventional therapy of congenital and structural heart disease. Live cases from 13 sites were transmitted via satellites to the venue at the Bellagio Hotel. The response we received so far is overwhelming with excellent feedback about the educational contents and the venue.

PICS started Sunday Sep. 10th, 2006 at 9:00 AM with an industry-sponsored symposium (AGA Medical Corp. Annual Meeting). This was very well attended. In this symposium, the Sim Suite was featured for PFO and ASD closure. Dr. Cheatham and Dr. Ruiz went over the cases with the attendees to demonstrate the protocol and closure steps. The addition of the Sim Suite to the tools available to the interventional cardiologists will greatly enhance the training and certification of physicians with the use of the Amplatzer devices.



The afternoon was marked by an exciting workshop on ASD's and PFO's. In this workshop, the majority of devices/techniques used (investigational and approved) for ASD and PFO closure were discussed. A total of 14 speakers discussed various devices and topics related to PFO and ASD closure. In addition to the devices used for closure, the use of ICE was discussed as well as the use of TCD to detect right to left shunt.

At the same time as the workshop, there were simultaneous sessions of oral abstract presentations. A total of 30 abstracts were selected out of 145 for oral presentations in congenital and valvular heart disease. This was followed by "meet the expert sessions." Again, this was a combined session between adult and pediatric cardiologists to discuss difficult cases that were presented by the attendees.

At the end of the day, a welcome reception was held in the exhibit area that held more than 30 exhibits representing different manufacturers of devices, catheters, etc.

The following day, Monday, September 11th was a combined day between the adult and pediatric cardiologists. The symposium was officially inaugurated and the activities started with live cases from Milan, followed by live cases from Frankfurt, Sao Paulo and Toronto. In these cases the operators discussed vari-



ous cases of congenital heart disease in adult patients. From Milan, Drs. Mario Carminati and Gianfranco Butera operated together and demonstrated excellent cases; from Frankfurt, Dr. Horst Sievert demonstrated two cases of PFO closure using different devices; Dr. Carlos Pedra, from Sao Paulo, demonstrated three excellent cases of various interventions and Drs. Lee Benson and Eric Horlick from Toronto demonstrated excellent two cases including a percutaneous pulmonary valve implantation.

The first lecture of the day was by Howard Herrmann and David Nykanen about the technique of trans-septal puncture in adults and children. This was followed by an excellent lecture by Bill Hellenbrand on coarctation stenting.

During the entire day, nine cases were transmitted live from the above mentioned centers, which demonstrated the latest techniques for adults with congenital heart disease. Further, ten



DON'T BUY ANY MORE MEDICAL EQUIPMENT - YET ...

MD Advantage Inc. will work with you to analyze your needs and the marketplace to help you make a well informed and cost effective decision for your practice. Spend more time with your patients and less time with sales people. Call MD Advantage today!

MD Advantage, Inc. 🔶 1-888-224-2234 🔶 www.MDAdvantageMedical.com

lectures from the course faculty were given addressing various topics including as mentioned above, trans-septal puncture techniques, coarctation stenting and a series of talks about the right ventricle outflow tract (given by Charlie Kleinman, Dianna Bardo, James Wilkinson, Kevin Walsh, Philipp Bonhoeffer and Ziyad Hijazi)! Finally, Drs. Ted Feldman and Mike Mullen gave a nice presentation on how to develop a successful program for adults with congenital and valvular heart disease.

The last activity of the day was the annual PICS achievement award. This year the award went to Dr. Philipp Bonhoeffer for his innovative work in congenital heart disease, specifically, for his development of percutaneous pulmonary valve implantation that paved the way for the percutaneous valve replacement/repair era we are witnessing.

Tuesday, September 12th, 2006 was the second full day of the meeting. There were two parallel sessions in two different ballrooms, one for congenital heart disease and the other one for valvular/structural heart disease. To each room, there were live cases transmitted via satellite from around the world. To the adult arena, cases were transmitted from Chicago and Vancouver. For the first time in a US course, a percutaneous aortic valve implantation was transmitted live from St. Paul's Hospital performed by Dr. John Webb, the leader in the world in this technique. This was followed by a case of Mitral valve repair also from Vancouver by Dr. Webb. Again, this case was the first to be transmitted live to a US course. From Chicago, I transmitted to the adult arena two cases: an ASD and a PDA device closure in two patients demonstrating the imaging techniques and the devices used.

To the Pediatric Arena, live cases were transmitted from Toronto Sick Kids with Dr. Benson who performed a percutaneous pulmonary valve implantation; from Columbus, OH where Drs. Cheatham and Galantowicz transmitted a case of hybrid therapy for a neonate with HLHS, while Dr. Amin performed three cases of LPA angioplasty, large ASD closure in a small child under TEE and ICE guidance to demonstrate the two techniques, and a coronary artery fistula that the attendees and the moderators felt that Dr. Amin should wait and not do the case, which he did! I finally transmitted a case of RVOT stent implantation in a child with severe conduit obstruction after a Rastelli operation.

That day, a total of 12 cases were transmitted to both arenas.

In terms of lectures: the adult arena focused Tuesday on the aortic valve where over 12 lectures/debates/discussions addressing the various issues related to the aortic valve. The list of the speakers was very impressive including Drs. Bob Bonow, Michael Mack, Allan Stewart, Ted Feldman, Marty Leon, Jeff

Interventional Catheterization

The Division of Pediatric Cardiology at the University of Utah School of Medicine is recruiting a pediatric cardiologist at the Assistant or Associate Professor level with a major interest in interventional catheterization. The candidate must have advanced training and/or experience in interventional catheterization. Our division is very busy clinically and academically productive with all subspecialty areas represented. In 2004, we performed approximately 650 catheterization procedures including a full spectrum of interventions. We expect to open 2 brand new cath labs in 2007 ensuring the continued growth of our interventional program. Opportunities exist for wide ranging clinical and bench-top collaboration both locally and nationally within both academics and industry.

In addition to expertise in interventional catheterization, applicants should have a sound clinical background in all areas of pediatric cardiology. The current division staff is composed of individuals with significant professional experience and expertise. There will be protected time for clinical research, with mentoring available within the division for clinical research studies. The division has a very active clinical research program and is one of the participating centers in the Pediatric Heart Disease Clinical Research Network funded by the NIH.

The successful candidate will receive a faculty appointment at the University of Utah. The pediatric cardiology division is based at Primary Children's Medical Center, a tertiary referral center for a three state area, which is located on the hills overlooking Salt Lake City. The area offers an excellent quality of life with immense cultural and recreational opportunities close and available. The University of Utah is an Equal Opportunity Employer and welcomes applications from minorities and women and provides reasonable accommodations to the known disabilities of applicants and employees.

Interested individuals should contact: Robert E. Shaddy, Professor of Pediatrics, Division Chief of Pediatric Cardiology, University of Utah School of Medicine, 100 North Medical Drive, Salt Lake City, Utah 84113. E-mail: robert.shaddy@ihc.com

The future of portable echo is here. Or there. Depending on where you need it.

11

Visit www.sonosite.com

MKT00836 © 2005 SonoSite, Inc. All rights reserved.





Moses, Alain Cribier, George Hanzel, Eberhard Grube, Craig Smith, Donna Buckley and a roundtable discussion that included Marty Leon, Gregg Stone, Peter Block, Larry Wood, Amr Salahieh, Jacque Seguin and Raoul Bonan. Dr. Webb gave his aortic valve talk via satellite from Vancouver.

In the afternoon sessions in the adult arena, the focus was on the PFO/ASD closure. Dr. John Carroll talked about the use of imaging modalities during PFO and ASD closure; this was followed by Bob Sommer who talked about the basics of PFO closure; Jon Tobis talked about strokes and PFO; Jose Suarez de Lezo talked about closure of ASD in adults and Horst Sievert talked about new PFO and ASD closure devices. Of note, on Monday, Horst transmitted live cases from Frankfurt, and traveled the whole day to come and give his talk. It was most certainly, appreciated by all. The last activity of the day in the adult arena was an excellent debate between Gregg Stone and Paul Kramer about PFO closure...I was told that Gregg won that debate calling for randomized trials before we can recommend PFO closure to prevent stroke recurrence.

The pediatric arena also had a very busy schedule of lectures and debates. The list of speakers and topics included: A mini





symposium on stents given by Felix Berger, Frank Ing, Chuck Mullins, Evan Zahn and David Bichell. Another mini symposium on ASD and PDA closure was also organized and speakers discussed various issues related to ASD and PDA including: echocardiographic evaluation in children by Charlie Kleinman, the Helex device by Geoff Lane; the Amplatzer device by Jozef Masura; closure of large PDAs by Krishna Kumar and closure of PDAs in premies by Neil Wilson.

Donna Buckley, MD from the FDA gave a talk about the appropriate trial design to seek device approval in the pediatric age group. This was followed by an interesting debate between Evan Zahn and Emile Bacha on coarctation angioplasty in children... this was certainly a lively debate.

During lunch there were some parallel sessions: the popular "my nightmare case in the cath lab." Needless to say, the moderators of this popular session (Neil Wilson and Mike Mullen) made this the most crowded session. Another session was on fetal intervention that featured the pioneering work of Wayne Tworetzky and Achi Ludomirsky.

Also, in the afternoon, there were parallel sessions for nurses and techs organized by Ms. Sharon Hill. On Tuesday, the focus was on Adult Congenital Heart Disease. In this session, Dr. Chuck Mullins, Dr. Stephen Cook, Ms. Elizabeth Sparks and Mr. Russel D'Sa gave excellent presentations addressing various issues related to the adult with CHD.

The last day, September 13th, was equally busy in both arenas. In the adult arena, live cases were transmitted from Columbia Medical Center in New York City and from Evanston, IL. Dr. Bob Sommer performed three excellent cases and Dr. Ted Feldman also performed 3 excellent cases of valvular and structural heart disease intervention. To the Pediatric Arena, Drs. Tom Jones, Mark Reisman performed three cases from Seattle and Dr. David Balzer performed also three cases from St. Louis. All cases that day were excellent and educational and the attendees enjoyed them tremendously.





Working Together to Develop a Better Tomorrow

NOVEMBER 2006

From lecture point of view, in the adult arena the focus that day was on the mitral valve and its diseases and the various new technologies available to treat MR/MS.

Drs. Allan Schwartz, Roberto Lang, Marc Gillinov, Francesco Maisano, Steve Bolling, Igor Palacios, Howard Herrmann, Gregg Stone, Maurice Buchbinder, Peter Block, John Webb, George Hanzel, Stephen Ellis, and Allan Stewart gave excellent talks addressing the various aspects of mitral valve disease and therapeutic tools. This was followed by a roundtable discussion featuring Marty Leon, Gregg Stone, Peter Block, Don Bobo, Ferlyn Powell and David Reuter. Discussion focused on the industry and medical community, and what can be done to bring these tools to patient side. A debate between Allan Schwartz and Michael Mack regarding mitral valve therapeutic techniques took place a day before. The debate was excellent and friendly. The last session for the adult arena was focused on the VSDs in adults given by Jo De Giovanni, PDAs in adults by Ziyad Hijazi, LAA imaging and closure by Roberto Lang and David Holmes, coronary fistulas by Mazeni Alwi and paravalvar leaks and their management by Jean Francois Piechaud.



In the pediatric arena, there were many topics discussed including a mini symposium on Hybrid intervention featuring John Cheatham and Mark Galantowicz who



discussed their pioneering work in HLHS, David Bichell and Emile Bacha talked about intra operative stent implantation and VSD closure. Then a mini symposium on VSD closure was organized and this featured the echocardiographic evaluation of VSD by Norm Silverman, techniques and results of VSD closure using the Amplatzer device by John Bass, closure of muscular VSD by Basil Thanopoulos, use of Nit Occlud for VSDs by TP Le.

Then, a very interesting and lively debate between Joaquim Miro and Redmond Burke about PmVSD closure, this was indeed a fun debate!

The last two talks of the meeting were given by John Cheatham on cutting balloons and Fontan Completion in the cath lab by Lee Benson.

The parallel sessions that day included a luncheon session on critical aortic valve disease beyond fetal life. This session featured Norm Silverman who discussed the echocardiographic features of this disease, Gil Wernovsky who discussed the ICU management of such patients, Shak Qureshi discussed the catheter option technique and Redmond Burke discussed the surgical management of these children. The other parallel session was focused on "Challenges in the sick neonates". In this symposium moderated by Gil Wernovsky and Sharon Hill, speakers touched on cardiac catheterization and







he Division of Pediatric Cardiology at OHSU Doernbecher Children's Hospital has a faculty position available at the assistant or associate professor level. The position requires excellent clinical skills in general pediatric cardiology. Subspecialty interest or training in heart failure, pulmonary hypertension and the care of cardiac transplant patients (currently performed elsewhere) is required. The candidate needs to be BE/BC in pediatric cardiology. In addition, the candidate should have excellent clinical judgment, a good work ethic, and the ability to collaborate with staff. This position will include ample opportunities for teaching and time to perform clinically focused research.

OHSU is located in Portland, Oregon, and is the state's only medical school. Doernbecher Children's Hospital is part of OHSU and includes 12 (soon to be 16) intensive care beds, 48 general pediatric beds, 12 (soon to be 24) oncology beds, and 48 neonatal intensive care beds. A well-established pediatric residency and pediatric cardiology fellowship program also exist. Portland is the largest city in Oregon and provides wonderful access to the Pacific Northwest.

Please send CV's to Mark Reller, M.D.

Division Head, Pediatric Cardiology • <u>rellerm@ohsu.edu</u> OHSU, Mail code CDRCP 707 S.W. Gaines Road





11th Vail Symposium on Pediatric Cardiac Diseases Vail Cascade Resort, Vail, Colorado March 4-7, 2007 Contact: dixon.jeannie@tchden.org or patilla.renee@tchden.org

www.CongenitalCardiologyToday.com

interventions in the critically ill neonate by Evan Zahn; perfusion modifications for cardiac catheterization while on ECMO by Vince Olshove; device closure in the neonate and small infants by Ziyad Hijazi, and premounted stents and hybrid procedures to palliate critically ill neonates by Sharon Hill.

After full three and a half days of intense lectures and live cases, the attendees were invited to the Gala night party that was held Wednesday September 13th, 2006 at the Bellagio Hotel. During the Gala, raffle ticket (30 GB iPOD[®] courtesy of Congenital Cardiology Today) was given to the person who traveled the longest to come to PICS. Also, AGA Medical and B. Braun held their annual raffle to give four attendees free four registrations for PICS-2007! Then the band which was excellent, entertained the attendees until midnight!!

So, we hope to see you all at PICS-XI to be held again at the Bellagio, in Las Vegas July 22-25th, 2007.

~CCT~



Ziyad M. Hijazi, MD, MPH, FACC, FAAP, FSCAI George M. Eisenberg Professor of Pediatrics and Medicine Staff Cardiologist, Section of Pediatric Cardiology University of Chicago 5841 South Maryland Ave., MC4051 Chicago, IL 60637 USA Tel: (773) 702.6172 Fax: (773) 834.4100

zhijazi@peds.bsd.uchicago.edu

Ziyad M. Hijazi Receives the PICS Founder Award



By CCT Staff

Ziyad M. Hijazi, MD, MPH, Professor of Pediatrics and Medicine at the University of Chicago received the PICS Founder Award at the Pediatric and Adult Interventional Therapies for Congenital and Valvular Heart Disease - PICS 2006, held in the Bellagio Resort in Las Vegas, September 10-13, 2006. Dr. William Hillenbrand, a Course Co-Director with Hijazi since the first PICS, presented the award. PICS, now marking its first decade, has become an annual institution, and is the premier international forum for the advancement and teaching of interventional procedures for congenital heart disease.

The hallmark of PICS is numerous live interactive case demonstrations from multiple international venues. PICS attendance has grown to approximately 1000 international congenital cardiologists and surgeons from its relatively humble beginning in Boston with about 80 American pediatric cardiologists attending the first PICS in 1997.

Pediatric Cardiology Services PC.



Pediatric Cardiology Services is a "state of the art" private practice in Atlanta. We are expanding in the metro Atlanta and surrounding area and are seeking a board certified/ eligible non invasive pediatric cardiologist.

Atlanta offers all the cultural amenities and educational institutions of a rapidly growing cosmopolitan city as well as an extremely favorable climate for outdoor activities in an extremely scenic suburbia.

We offer a fantastic salary coupled with comprehensive benefits including CME, Health insurance, Life insurance, malpractice insurance etc.

Our offices are equipped with state-of-the-art diagnostic equipment.

We provide comprehensive services including:

- fetal cardiology
- echocardiography
- electrocardiography
- ambulatory rhythm monitoring
- exercise stress testing
- tilt-table testing

Neill Videlefsky, M.D. Shiva Sharma, M.D. Pediatric Cardiology Services, PC.

email to: nvidele@mindspring.com tel. no.: 770 995 6684



DON'T BUY ANY MORE MEDICAL EQUIPMENT - YET ...

MD Advantage Inc. will work with you to analyze your needs and the marketplace to help you make a well informed and cost effective decision for your practice. Spend more time with your patients and less time with sales people. Call MD Advantage today!

MD Advantage, Inc. ♦ 1-888-224-2234 ♦ www.

www.MDAdvantageMedical.com

MEDICAL NEWS, NEW PRODUCTS AND INFORMATION

Medtronic Announces CE Mark Approval of the World's First Transcatheter Valve Replacement System For Structural Heart Disease

Signaling a new frontier in the use of transcatheter valve technology, Medtronic, Inc. has announced that it has received CE Mark approval for its Melody[™] Transcatheter Pulmonary Valve and Ensemble[™] Transcatheter Delivery System. The system is the first of its kind worldwide to treat congenital patients with structural heart disease requiring pulmonary heart valve replacement.

The Medtronic Melody[™] Transcatheter Pulmonary Valve and Ensemble™ Transcatheter Delivery System is designed to benefit patients with congenital heart defects involving the connection between their right ventricle and pulmonary artery. Oftentimes, this type of defect requires open heart surgery early in life to implant a prosthetic valvedconduit to establish adequate blood flow from the heart to the lungs. However, the functional life span of these conduits is relatively limited, and as a result most patients with this type of defect are committed to multiple open heart surgeries over their lifetime. The Melody valve and Ensemble system provide a non-surgical means to restore effective valve function and prolong the functional life of prosthetic conduits, thereby reducing the number of open heart surgeries for these patients throughout their lifetime.

"While the population that will benefit from this new technology is relatively small, Medtronic is committed to patient-centered lifetime management of congenital heart disease," said Oern Stuge, MD, senior VP and president of the Cardiac Surgery business at Medtronic. "The Melody Valve and Ensemble System are examples of Medtronic's commitment to evolving the science and focus of structural heart disease. This focus further demonstrates our dedication to delivering innovative valve technology that addresses unmet patient needs. By providing less invasive options for physicians and therefore eliminating even one open heart procedure for a patient, we improve that patient's quality of life immeasurably."

Medtronic has partnered with Professor Philipp Bonhoeffer, professor and chief of cardiology and director of the Catheterization Laboratory, Great Ormond Street Hospital for Children (GOSH) in London, and NuMed, a pediatric catheter company in the development of the Melody Valve and Ensemble System. A pioneer in transcatheter valve technology, Professor Bonhoeffer implanted the world's first transcatheter valve in September 2000. To date, more than 150 patients worldwide have been implanted with the Melody Transcatheter Pulmonary Valve.

Transcatheter valve technology represents a less invasive means to treat heart valve disease and is designed to allow physicians to deliver replacement valves via a catheter through the body's cardiovascular system, thus eliminating the need to open the chest. Traditionally, open heart surgery has been required to correct the problem and it is not unusual for a patient to undergo multiple, open-heart surgeries during their lifetime. Patients with this condition tire very easily, as the heart over-exerts itself trying to get oxygenated blood throughout the body.

"To reopen the chest many times is obviously uncomfortable for the patient but also comes with risk. By using the Melody Valve and Ensemble System, we can now avoid the open chest procedure and instead perform the same procedure safely, quickly, and more comfortably," said Prof. Bonhoeffer.

CE Mark Approval gives Medtronic the ability to train and educate physicians in Europe in preparation for implanting the Melody[™] and Ensemble[™] System so that more patients can benefit from this innovative treatment. With proper training in place, Medtronic will work with hospitals across Europe in gaining access to this new technology.

While the Melody Valve and Ensemble System are not currently available in the United States, Medtronic is working with the U.S. Food and Drug Administration to pursue plans to apply this new approach in transcatheter valve therapy for patients in the U.S.

For more information about the product and treatment visit www.melody-tpv.com.

The International Children's Heart Foundation Needs Volunteer Pediatric Cardiac Specialists for Upcoming Missions

William M. Novick, MD of the University of Tennessee Health Sciences Center, and Founder and Medical Director of the International Children's Heart Foundation is looking for volunteers for the following missions:

- Armed Forces Institute of Cardiology/ National Institute of Heart Disease, in Rawalpindi, Pakistan; Dec. 2-16, 2006: Pediatric Cardiac Anesthesiologist (1) and pediatric cardiac perfusionist (1).
- Beijing Children's Hospital, Beijing, China; Jan. 6-20, 2007: Pediatric Cardiac Anesthesiologist (1).
- Armed Forces Institute of Cardiology/ National Institute of Heart Disease, in Rawalpindi, Pakistan; Jan. 27- Feb. 10, 2007: Pediatric Cardiac Anesthesiologist (1), Pediatric Intensivist (2), Pediatric Cardiologist (1).

Interested individuals should contact Martina Pavanic, Volunteer Coordinator, at martapavanic@aol.com.



11th Vail Symposium on Pediatric Cardiac Diseases Vail Cascade Resort, Vail, Colorado March 4-7, 2007 Contact: dixon.jeannie@tchden.org or patilla.renee@tchden.org

BERRY SYNDROME IN A NEONATE: REPORT OF A CASE SUCCESSFULLY REPAIRED IN A ONE-STAGE OPERATION

By Marcì Marcello, MD; Calvaruso Davide, MD; Cipriani Adriano, MD; Ocello Salvatore, MD; Salviato Nicoletta, MD; and Marcelletti F. Carlo, MD

Case Report

A severely ill infant, weighting 3,1 Kg, was referred to our Institution soon after the birth for severe cyanosis, metabolic acidosis (pH = 7,1) and anuria (creatininemia =3,5 mg/dL). Physical examination revealed tachypnea and a systolic murmur while femoral pulses were weak. Radial and femoral arterial pressures, invasively monitored showed a difference of about 30 mmHg. Similarly oxygen saturation was higher in upper limbs. Echocardiography showed interruption of the aortic arch, type A, according to the classification of Celoria and Patton. Thus flow in the descending aorta was provided by a patent ductus arteriosus. Furthermore, aortopulmonary window and anomalous origin of the right pulmonary artery from ascending aorta were associated.

Correction of acidosis, PGE1 and dopamine infusion and diuretics allowed rapid recovery with stabilization of hemodynamic conditions and balance of oxygen saturation and arterial pressure in upper and lower limbs, with subsequent recovery of renal function.

Cardiac catheterization and angiocardiography confirmed the presence of aortic arch interruption associated with PDA, aorto-pulmonary septal defect and anomalous origin of RPA. At 5 days of age the baby underwent total repair with deep hypothermic circulatory arrest a 14°C nasopharingeal.

After aortic cross-clamping distal to the defect, cardioplegia was infused through an arterial cannula into the ascending aorta that was incised transversely showing a very large aorto-pulmonary window. The coronary artery origins were normal, the pulmonary and aortic valve leaflets were identified. The right pulmonary artery originated directly from the right-lateral wall of the ascending aorta. After the great vessels were divided, both aortic and pulmonary defects were closed using two different pericardial patches with continuous 6/0 polypropylene suture. The PDA was divided and all ductal tissue was removed. Right pulmonary artery was excised from aorta and then anastomosed to the main pulmonary artery. The mobilized descending aorta was end-to-end anastomosed to aortic arch with an 8 mm goretex prosthetic tube. Wean from cardiopulmonary bypass was early obtained.

The recovery from operation was uneventful and the patient was discharged home on the 20th postoperative day.

Comment

Aorto-pulmonary septal defect is a rare malformation: its incidence ranges from 0.2% to 0.6 % in patients with congenital heart diseases, according to Kutsche and Van Mierop.

It may be associated in about 25% of patients with interruption of aortic arch, usually type A of Celoria and Patton classification.

"The association of aortopulmonary septal defect, PDA and anomalous origin of RPA from ascending aorta is extremely rare and was reported in 1982 by Berry et al. This potentially lethal lesion may be a tough diagnosis for the cardiologist, because an accurate echocardiographic identification is not always obvious. Often a cardiac catheterization is needed to confirm the diagnosis."

Etiological factors involved in the development of aortic arch interruption (AAI) and aorto-pulmonary defect (APW) are not well defined. Though both abnormalities involve the aorta, they have different embryological pathogenesis. APW should be caused by lack of development of aorto-pulmonary septum in the aortic sac, which appears late in embryogenesis and is believed to be caused by reduction of blood flow toward the descending aorta during intrauterine life because of the frequent association with intra-cardiac lesions with obstruction to the left ventricular outflow tract and/or VSD. This hypothesis has been confirmed by experimental observations in animal embryos, in which aortic interruption was caused by re-



The Barth Syndrome Foundation

P.O. Box 974, Perry, FL 32348

Tel: 850.223.1128 info@barthsyndrome.org www.barthsyndrome.org **Symptoms:** Cardiomyopathy, Neutropenia, Muscle Weakness, Exercise Intolerance, Growth Retardation

striction of the conotruncus. According to this hypothesis AAI should be the consequence of reduction of blood flow into the ascending aorta. In addition, a genetic disorder could play a role in development of interruption of aortic arch, as demonstrated by the association of this abnormality with Di George syndrome. Whereas, aortopulmonary septal defect has never been reported in patients with chromosomal delection in 22g11. In fact, development of aortic arch and of aortopulmonary outflow tract are influenced by migration of neural crest cells to the branchial arches and to the aortic sac.

Removal of the cardiac neural crest RPA or hemitruncus arteriosus. Acregion in the animal embryo can determine truncus arteriosus and type A interruption, but not defect of aorto-pulmonary septum. RPA or hemitruncus arteriosus. According to Richardson's classification, Berry syndrome would be a combination of two different types of APW. Actually, hemitruncus is not consid-

17

Several classifications of APW have been proposed according to the size and location of the defect. In Richardson et al's classification three types of aorto-pulmonary defect are described: type I, the most common, is a defect localized between ascending aorta and pulmonary trunk, in the type II the defect extends distally toward the bifurcation of the pulmonary artery, the type III is an anomalous origin of the

RPA or hemitruncus arteriosus. According to Richardson's classification, Berry syndrome would be a combination of two different types of APW. Actually, hemitruncus is not considered a form of aorto-pulmonary defect by other authors. Mori et al. for example, distinguish three forms of APW: "proximal defect" is just above the semilunar valves, "distal defect" is near the bifurcation of pulmonary artery, and "total defect" involves most of the ascending aorta and of the main pulmonary artery.

In the majority of cases, aortopulmonary septal defect is frequently associated to



Arkansas Children's Hospital, Little Rock, AR Clinical Fellowship – Pediatric Cardiac Intensive Care

The Divisions of Pediatric Cardiology and Pediatric Critical Care Medicine at the University of Arkansas for Medical Sciences College of Medicine solicit applications for a clinical fellowship in Pediatric Cardiac Intensive Care for the academic year 2006-2007 and 2007-08.

This training is available for a period of 6-12 months to individuals who are BC/BE in either Pediatric Cardiology or Pediatric Critical Care Medicine. All clinical activities will take place at the Arkansas Children's Hospital Heart Center. Care is provided in a 15 - Pediatric Cardiac Intensive Care Unit 24/7 by a dedicated and highly specialized team of intensivists, cardiologists, cardiac surgeons, cardiac anesthesiologists, advanced nurse practitioners, nurses and allied health professionals. The Cardiac Intensive Care Unit admits approximately 600 patients per year. The pediatric and congenital cardiovascular surgery team led by Dr Robert DB Jaquiss performs approximately 550 congenital heart surgeries yearly with approximately one half being cardiopulmonary bypass cases. The program has a very strong neonatal cardiac surgery, heart transplant, ventricular assist device and mechanical support focus. Arkansas Children's Hospital maintains a very busy ECMO program, with one of the Nation's only mobile ECMO programs, and provides formal training in ECMO support of cardiac patients. The fellows will also have the opportunity of working closely with dedicated pediatric cardiac anesthesiologists during their training. The hospital also houses the Donald W. Reynolds Pediatric Intensive Care Unit, a state-of-the-art 26-bed facility which receives approximately 1100 admissions annually, where the fellow will receive training in lung-protective ventilator strategies and support of multi-organ dysfunction.

The program presents a very conducive environment for clinical training in various aspects of care of critically ill children with congenital heart disease. In general, this position will entail approximately 85% clinical time and 15% time for clinical research. There are numerous research opportunities and the fellow is encouraged to carry out meaningful clinical research during the course of the fellowship. The clinical training and responsibilities will be tailored for the individual candidate, depending on prior fellowship experience (i.e. Critical Care or Cardiology) and personal interests.

Clinical fellows will be appointed to the Department of Pediatrics, UAMS at the Instructor level in a non-tenure track position with faculty benefits including health and liability insurance. Applicants accepted for this clinical fellowship will be required to obtain an Arkansas medical license prior to starting training. The University of Arkansas Medical Sciences is an Equal Opportunity Employer. Little Rock, the capital of Arkansas is a warm and friendly cosmopolitan community with affordable housing, is well connected to other parts of the country and offers an abundance of recreational activities.

Interested individuals please contact:

Parthak Prodhan, MD Pediatric Critical Care and Cardiology University of Arkansas for Medical Sciences 800 Marshall Street Slot 512-3 Little Rock, AR 72202-3591 Tel: 501-364-1479; Fax: 501-364-3667 ProdhanParthak@uams.edu truncus arteriosus. Ac-

NOVEMBER 2006

NOVEMBER 2006

other lesions, such as type A interruption of aortic arch, anomalous origin of right coronary artery from pulmonary artery, tetralogy of Fallot, transposition of great arteries, right aortic arch and anomalous origin of the RPA. Moreover association with ventricular septal defect, d-transposition, pulmonary or aortic atresia have also been described. Patients affected by APW and associated lesions have a worse prognosis. The association of aortopulmonary septal defect, PDA and anomalous origin of RPA from ascending aorta is extremely rare and was reported in 1982 by Berry et al. This potentially lethal lesion may be a tough diagnosis for the cardiologist, because an accurate echocardiographic identification is not always obvious. Often a cardiac catheterization is needed to confirm the diagnosis. An early surgical treatment is recommended to avoid the development of irreversible obstructive pulmonary vascular disease.

Boonstra et al. reported the first surgical repair of such syndrome in 1992. To our knowledge, since then, only a few other cases of Berry syndrome, surgically treated, have been reported in literature.

Summary

We report a case of Berry syndrome, a rare combination of aorto-pulmonary malformations, in a neonate successfully treated with primary surgical repair.

See authors for reference information.

~CCT~

Corresponding Author:

Dott. Marcello Marcì, MD Via Annibale 6 B- 90149 Palermo, Italy

marcellomarci@katamail.com

Calvaruso Davide, MD

Department of Cardiac Surgery ARNAS Ospedale Civico Palermo, Italy

Cipriani Adriano, MD Department of Cardiac Surgery, ARNAS Ospedale Civico Palermo, Italv

Ocello Salvatore, MD Department of Cardiac Surgery ARNAS Ospedale Civico Palermo, Italy

Salviato Nicoletta, MD Department of Cardiac Surgerv ARNAS Ospedale Civico Palermo, Italy

Marcelletti F. Carlo, MD Department of Cardiac Surgery ARNAS Ospedale Civico Palermo, Italy

Would you like to share your interesting stories or research in congenital cardiology?

Submit a brief summary of your proposed article to Congenital Cardiology Today at Article@CCT.bz

The manuscript may be between 400-3,500 words, and contain pictures, graphs, charts and tables.

MEDICAL CONFERENCES

Neonatal Nursing Excellence: Knowing and Growing Nov. 8-11, 2006; Nashville, TN USA www.NANN.org

Fourth Annual Symposium on Advances in Perinatal Cardiology Nov. 16-18, 2006; St. Petersburg, FL USA www.allkids.org/conferences/cme/ conferences/perinatalcard2006/ index.htm

5th Annual March of Dimes Prematurity Summit : Time to Take Action Nov. 17, 2006; Minneapolis, MN USA www.marchofdimes.com/minnesota

Second International Conference on Heart Failure in Children and Young Adults Nov. - Dec. 2, 2006; Laguna Niguel, CA USA

www.cbcbiomed.com

Combined Cardiovascular Pathology Nov. 29 - Dec. 1, 2006; London, UK mailto:Morphology@rbht.nhs.uk

Euroecho 10 Dec. 6-9, 2006; Prague, Czech Republic www.euroecho.org

International Congress for Genetics in Pediatrics Jan. 24-27, 2007; Luxor, Egypt www.genevent.org

The Conference on Neonatology Feb. 7-10, 2007; Lake Buena Vista, FL USA

www.pediatrix.com

Cardiology 2007 Feb. 21-25, 2007; Lake Buena Vista, FL USA www.chop.edu/cme/

Hands-on Cardiac Morphology Feb. 26-28, 2007; London, UK mailto:Morphology@rbht.nhs.uk

Cardiomyopathy Foundation

In support of infants, children and teens with pediatric cardiomyopathy

CHILDREN'S CARDIOMYOPATHY FOUNDATION

P.O. Box 547, Tenafly NJ 07670

Tel: 201-227-8852 info@childrenscardiomyopathy.org www.childrenscardiomyopathy.org

"A Cause For Today.... A Cure For Tomorrow"



19

PEDIATRIC CARDIOLOGIST

Montgomery, Alabama

We seek a non-invasive BC/BE Pediatric Cardiologist to join Pediatric Specialists of Montgomery, a respected private practice in Montgomery, Alabama. The practice is 90% office based and 10% NICU and inpatient consulting. Patients are followed from birth to college. The practice is affiliated with two Baptist hospitals with large, well staffed NICUs. Both hospitals have EMR. EMR is planned for the practice.

The practice is located in a spacious new building in a rapidly growing area near one of the hospitals and has in-house echo and stress test, Holter and Event Monitoring capability. Office hours are 7:30 - 4:00. Staff is in place to support two busy physicians.

Montgomery has a strong economy, low cost of living, vibrant arts community, minor league baseball team, excellent housing choices, public magnet schools and excellent private schools, and multiple colleges including Auburn University Montgomery.

The ideal candidate will be committed to practicing quality community medicine, have excellent relationship skills, a strong work ethic, and a collegial attitude toward physicians, office and hospital staff. He/ she will be interested in becoming an integral part of this dynamic growing community. Interest in Fetal Echoes or TEE will be a plus. Excellent compensation & benefits are offered.

Contact: Teresa Unser Resource Consulting Group, Inc. 30 Southridge Trail Cartersville, GA 30120 Toll free: 866-975-7399 Phone: 770-975-7399 Fax: 770-975-8838. Email: teresaunser@bellsouth.net



Asheville Cardiology Associates Recruiting BC/BE Pediatric Non-Invasive Cardiologist For Summer 2007

Known for clinical excellence, Asheville Cardiology Associates is seeking a third pediatric cardiologist with advanced skills in both inpatient and outpatient arenas. Those with expertise in new modality imaging, adult-congenital cardiology, or interest in exercise physiology or electrophysiology, are encouraged to contact us. Development of new programs within the practice is encouraged.

The Peds Division has a complex patient base, with active fetal and adult-congenital programs. Outpatient practice encounters ~3,000 clinic visits, ~2,000 echoes per year, with a growth rate of 5-7% annually. Inpatient is performed at Mission Hospitals, consistently rated in Top 100 Heart Hospitals in last six years. Its new children's outpatient facility is well represented by subspecialists and ancillary services.

Asheville is a beautiful city of 70,000 (county has 200,000 plus MSA of 391,000). Located in the Blue Ridge Mts. of North Carolina, it offers excellent public and private schools, universities and colleges. Amenities such as great restaurants, arts, music, & theater put it on the list of best places to live in many publications. A mild four season climate gives opportunities for mountain biking, hiking, camping, fishing, golf, whitewater activities and skiing.

Visit our website, www.avlcard.com for more information.

All inquiries remain confidential. Please send CV to:

James J. McGovern, MD, FACC Fax: (828) 277-6350 Email: jimm@avlcard.com

CONGENITAL CARDIOLOGY TODAY

© 2006 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-0499-online) Published monthly. All rights reserved

Publishing Management

Tony Carlson, Founder TCarlsonmd@gmail.com Richard Koulbanis, Publisher & Editor-in-Chief RichardK@CCT.bz John W. Moore, MD, MPH, Medical Editor/ Editorial Board JMoore@CHSD.org Editorial Board Teiji Akagi, MD Zohair Al Halees, MD Mazeni Alwi, MD Felix Berger, MD Fadi Bitar, MD Jacek Bialkowski, MD Philipp Bonhoeffer, MD Anthony C. Chang, MD, MBA John P. Cheatham, MD Bharat Dalvi, MD, MBBS, DM Horacio Faella, MD Yun-Ching Fu, MD Felipe Heusser, MD Ziyad M. Hijazi, MD, MPH Ralf Holzer, MD Marshall Jacobs, MD R. Krishna Kumar, MD, DM, MBBS Gerald Ross Marx. MD Tarek S. Momenah, MBBS, DCH Toshio Nakanishi, MD, PhD Carlos A. C. Pedra, MD Daniel Penny, MD James C. Perry, MD Shakeel A. Qureshi, MD Andrew Redington, MD Carlos E. Ruiz, MD, PhD Girish S. Shirali, MD Horst Sievert, MD Hideshi Tomita, MD Gil Wernovsky, MD Zhuoming Xu, MD, PhD William C. L. Yip, MD Carlos Zabal, MD To Contact an Editorial Board Member Email to: BOARD @CCT.bz. Place the Board Member's name in the Subject line. **FREE Subscription** Congenital Cardiology Today is available free to qualified professionals worldwide in pediatric and congenital cardiology. International editions available in electronic PDF file only; North American edition available in print. Send an email to Subs@CCT.bz. Be sure to include your name, title, organization, address, phone,

fax and email. Contacts and Other Information

For detailed information on author submission, sponsorships, editorial, production and sales contact, current and back issues, see website: www.CongenitalCardiologyToday.com

Do You Want to Recruit a Pediatric Cardiologist?

Advertise in the only monthly publication totally dedicated to pediatric and congenital cardiology. For more information: call +1.301.279.2005, or send an email to: TCarlsonmd@gmail.com

n the last 20 years, there has been a revolution in caring for children with congenital heart disease. Thanks to advances in diagnostics, surgery and treatment, many of these children are living into adulthood and enjoying active lives. With dedicated pediatric cardiology capabilities, the iE33 system answers your need for high performance, non-invasive diagnostic imaging for these patients. Now, the new X7-2 transducer combines xMATRIX technology with PureWave crystal technology for remarkable Live 3D Echo imaging in small patients.

The X7-2 transducer's footprint is small enough to allow intercostal access in pediatric applications and it uses an extremely lightweight, ultra-flexible cable. In addition to excellent 3D image quality, the X7-2 offers PW Doppler and superb color sensitivity in an impressively compact form factor.

The X7-2 transducer joins a full complement of cardiac transducers on the iE33 system, including pediatric and adult TEEs, to enable you to image your complete range of patient types – regardless of age, disease state or body composition. In addition, a unique analysis package truly follows the patient the way you do – so you don't have to fit your requirements into an analysis package designed for adult echocardiography.

Dedicated to your pediatric cardiology clinical and workflow requirements, our commitment goes beyond the system and its technologies. Global educational programs, outstanding service and worldwide centers of excellence help you get the most out of your system every day.





FOR MORE INFORMATION

Philips Medical Systems 22100 Bothell Everett Highway Bothell, WA 98021 +1 (425) 487 7000 1 (800) 285-5585

www.medical.philips.com/ultrasound

Expanding the clinical utility of echo with Live 3D specifically developed for pediatric imaging



The new X7-2 pediatric Live 3D transducer shown actual size.

