# ΟΝGΕΝΙ

RELIABLE INFORMATION I N CONGENITAL CARDIOLOGY

### NORTH AMERICAN EDITION

www.CongenitalCardiologyToday.com

VOL. 3, ISSUE 6 JUNE 2005

Formerly Pediatric Cardiology Today

### **INSIDE THIS ISSUE**

Long-Term Issues of Coronary Artery Sequelae in Kawasaki Disease by Teiji Akagi, MD, PhD	1
<b>The Pediatric Heart</b> <b>Network</b> <i>by Gail D. Pearson, MD, ScD</i>	5
Percutaneous Closure of a Residual Perimembranous Ventricular Septal Defect After Surgical Repair by Carlos A.C. Pedra, MD; Sérgio C. Pontes Jr, MD; Simone R.F. Pedra, MD; Juliana Neves, MD; M. Aparecida P. Silva, MD; M. Virginia T. Santana, MD; Valmir F. Fontes, MD	8
DEPARTMENTS	
Medical Conferences	4

September Conference	12
Focus (PICS/ENTICHS 2005	
(Pediatric Interventional	
Cardiac Symposium and	
Emerging New Technologies	
in Congenital Heart Surgery)	
September 15-18, 2005;	
Buenos Aires, Argentina)	
Medical News and	13
Information	

CONGENITAL CARDIOLOGY TODAY 9008 Copenhaver Drive, Ste, M Potomac, MD 20854 USA www.CongenitalCardiologyToday.com

© 2005 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-049 online). Published monthly. All rights reserved. Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and are not necessarily the views of Congenital Cardiology Today.

# LONG-TERM ISSUES OF CORONARY ARTERY SEQUELAE IN KAWASAKI DISEASE

By Teiji Akagi, MD, PhD

Kawasaki Disease is not really "congenital heart disease," but this acquired disease is of considerable interest to congenital cardiologists. The importance of Kawasaki Disease is not only the coronary artery aneurysm in the acute phase but also long-term cardiovascular sequelae, which may include risk of early atherosclerotic changes in adolescents or young adults.

Clinically, Kawasaki Disease is an acute, self-limited vasculitis of unknown etiology that occurs predominantly in infants and young children. This vasculitis frequently affects small to mid-size arteries especially coronary arteries. In Japan, 17 nationwide surveys have been conducted (every 2 years

since 1970) and more than 180,000 patients have been registered. Although no nationwide outbreak has been observed since the outbreak in 1986, the incidence rate has gradually increased over the past 16 years. The disease also has been reported in more than 60 countries around the world. In the United States, an estimated 4248 hospitalizations associated with Kawasaki disease occurred in 2000, with a median patient age of 2 years. Currently, the incidence of coronary artery abnormalities is about 3 to 5%. This means that nearly 200 children develop coronary artery abnormalities due to Kawasaki Disease in each year in the United States.

The most striking feature of coronary artery abnormalities in Kawasaki Disease is the change of size or shape of aneurysm. On the other hand, coronary descending artery.

artery stenosis occurs in 4% of patients, or in 20% with coronary aneurysms in the followup period. In general, coronary artery stenosis does not develop in patients after aneurysms regress .

### Does Kawasaki Disease increase the risk of atherosclerosis?

One unanswered question is whether the regressed coronary lesion becomes normal vessel or not. The major mechanism of regression of coronary aneurysms is intimal proliferation derived from the smooth muscle cells of media and regenerated endothelium. Thrombus formation in the aneurysm and calcification of arterial wall may combine in some instances. These findings are similar to atherosclerotic lesions.

Thus, recent long-term studies have focused



About 50% of coronary aneu- Figure 1. Fifteen-year old boy complained of mild chest pain. Coronary rysms regress within 2 years. angiography revealed the significant stenosis on his left anterior

CONGENITAL CARDIOLOGY TODAY Formerly Pediatric Cardiology Today

© 2005 by Congenital Cardiology Today (ISSN 1554-7787 - print; ISSN 1554-049 - online) Published monthly. All rights reserved.

Publishing Management
Tony Carlson, Founder TonyC @CCT.bz
Richard Koulbanis, Publisher & Editor RichardK@CCT.bz
John W. Moore, MD, MPH, Medical Editor & Medical Editorial Board jwmoore @mednet.ucla.edu
Editorial Board
Zohair Al Halees, MD
Mazeni Alwi, MD
Felix Berger, MD
Fadi Bitar, MD
Philipp Bonhoeffer, MD
Anthony C. Chang, MD, MBA
Bharat Dalvi, MD, MBBS, DM
Horacio Faella, MD
Yun-Ching Fu, MD
Felipe Heusser, MD
Ziyad M. Hijazi, MD, MPH
Ralf Holzer, MD
R. Krishna Kumar, MD, DM, MBBS
Gerald Ross Marx, MD
Tarek S. Momenah, MBBS, DCH
Toshio Nakanishi, MD, PhD
Carlos A. C. Pedra, MD
James C. Perry, MD
Shakeel A. Qureshi, MD
Andrew Redington, MD
Carlos E. Ruiz, MD, PhD
Girish S. Shirali, MD, MBBS
Horst Sievert, MD
Hideshi Tomita, MD
Gil Wernovsky, MD
Carlos Zabal, MD
To Contact an Editorial Board Member

Email to: NAME @CCT.bz. Place the Board Member's surname before @.

on functional abnormalities of coronary arteries, especially endothelium function. Dhillon and colleagues evaluated endothelial function after Kawasaki Disease using flow mediated dilatation of brachial artery. In response to reactive hyperemia, a measure of endothelium-dependent vasodilation, Kawasaki Disease patients had significant lower vasodilation compared to controls. In response to sublingual glyceryl trinitrate, a measure of endotheliumindependent vasodilation, the mean amount of dilatation was similar in the controls. This study suggested that endothelial dysfunction is present, possibly throughout the systemic arterial bed, in patients who have had Kawasaki disease. However, using a similar technique, a group from Toronto reported that there were no significant differences in either flow mediated or nitroglycerine mediated vasodilatation in brachial arteries of Kawasaki Disease patients.

We conducted a study of endothelial function evaluating intracoronary infusion of acetylcholine. The control or normal sites of coronary artery in Kawasaki Disease patients demonstrated significant vasodilatation, whereas the regressed aneurysms or persistent aneurysms revealed vasoconstriction or no changes. These findings suggest impaired endothelial function. Interestingly, patients having more than 4 mm aneurysms in the acute stage of Kawasaki Disease have significantly decreased coronary endothelial function, even if their coronary arteries appear to be completely normal by angiography or echocardiography. Evaluation of Kawasaki Disease patients without coronary artery lesions in the acute stage of illness using intravascular ultrasound imaging revealed normal findings at 10 to 20 years after the onset of disease. Patients whose original aneurismal size was less than 4 mm also had normal intravascular ultrasound findings. However, patients whose original aneurismal size was larger

"Clinically, Kawasaki Disease is an acute, self-limited vasculitis of unknown etiology that occurs predominantly in infants and young children."

than 4 mm, revealed thickened intima and media. From our studies of pathology with intravascular ultrasound and by pharmacological studies of vascular function, it appears that coronary artery lesions may be long-term coronary risk factors, even in regressed aneurysms. Other coronary risk factors such as hyperlipidemia, smoking or hypertension may accelerate these conditions. Furthermore, a recent study reported that CRP levels were significantly elevated in Kawasaki Disease patients with coronary artery abnormalities compared with those without coronary abnormalities. If the inflammatory response as well as endothelial dysfunction are present late after Kawasaki Disease, anti-inflammatory medication such as aspirin, beta-blocker or statin treatment may be considered in the future.

### New therapeutic strategy in Kawasaki disease

During the past decade, the clinical experience with interventional treatment in Kawasaki Disease has been gradually increasing. This includes balloon angioplasty, stent implantation, rotational ablation, and transluminal coronary revascularization. However, the experiences in Kawasaki Disease are still extremely limited compared to coronary intervention in adults. Coronary artery stenosis in Kawasaki Disease commonly involves severe calcifi-



### The SONOS 7500 system with Live 3D Echo ... for better visualization of congenital heart defects

Learn more at: www.medical.philips.com/live3D or call 800-229-6417







Figure 2. PTCRA using 2.15 mm burr.

Figure 3. Additional POBA using low pressure (4 mm, 8 atmosphere).

cation, in contrast to adult coronary artery lesions, which consist primarily of atherosclerosis. Therefore, the indications for catheter intervention for adult patients cannot be directly applied in Kawasaki Disease patients, a largely pediatric population. Currently, the fundamental therapeutic regimen for ischemic heart disease after Kawasaki Disease is coronary bypass surgery. However, the long-term coronary graft patency is not satisfactory, even with the use of intrathoracic arterial bypass graft, and especially if the operative age is under 12 years old.

Plain old balloon angioplasty (POBA) is a useful treatment of stenotic lesions with or without mild calcification within 6 years of the onset. Satisfactory acute results were obtained in patients with up to 6 years interval from the onset of the disease and the POBA. However, the incidence of restenosis after POBA was high. Approximately, one quarter of the patients developed restenosis or occlusion. Coronary arteries with thick intimal hyperplasia probably recoil significantly, even if the stenosis has been dilated sufficiently. Stent implantation may prevent the occurrence of restenosis in this situation.

The development of new coronary aneurysms after POBA is a significant clinical concern. The mechanism responsible for this phenomenon is unclear, however intimal and/or medial dissection due to high-pressure balloon inflation could play a role. In this regard, the maximum balloon pressure is recommended to be less than 8-10 atmospheres at the time of POBA in the pediatric population. In resistant coronary artery lesions, (which require greater than 10 atmospheres), rotational ablation or bypass surgery is advisable as alternative

procedures. Additionally, when POBA is performed in young children, shorter balloon lengths should be selected.

Experience with use of stent implantation for Kawasaki Disease is still limited. However, acute results with stent implantation have been excellent. Because this procedure relies on balloon dilation, the limitations are similar to those for POBA. The advantages of this procedure include high vascular patency rates in patients with segmental or relatively long stenosis, and the prevention of new aneurysm formation after POBA. Anticoagulation therapy with aspirin and clopidogrel is recommended after the procedure.

Excellent acute results for percutaneous rotational ablation (PTCRA) were observed in previous studies. Although use of this procedure is still limited,



Sponsored By

For information, please call 1-800-BBRAUN2 (227-2862) or visit www.bbraunusa.com

NORTH AMERICAN EDITION

### **CONGENITAL** CARDIOLOGY TODAY



Figure 4. Stenosis was completely resolved.

PTCRA may be the most appropriate catheter intervention for Kawasaki Disease. The advantage of PTCRA is the high success rate, even in patients with calcified coronary artery stenosis. The limitation is the need for larger arterial access for the metal burr. For this reason, PTCRA can only be performed in older patients. Using this procedure, the stenotic area can be dilated up to 2.5 mm. Anti-coagulation and antiplatelet medication should be continued at least 2 months after the procedure.

Based on our results, we make the following recommendations: POBA is effective in many situations, particularly in patients without severe calcification or in patients with a relatively short interval (within 6 years) between the onset of the disease and the intervention. POBA may be used even in small children. Therefore, POBA should be the first line procedure in younger children with significant coronary artery stenosis. Stent implantation is preferable to POBA, because it may prevent new aneurysm formation and restenosis. In adolescents and older patients, stent implantation should be considered instead of POBA alone. If patients have severely calcified coronary stenosis, PTCRA may be the only effective

treatment. Intravascular ultrasound imaging provides valuable information for the selection of the appropriate interventional procedure and early detection of vascular complications.

### Kawasaki Disease is the leading cause of acquired heart disease in children

Compulsive follow-up is required in patients with coronary aneurysms, especially if the original aneurismal size is larger than 4 mm. Early detection of stenotic lesions is essential Kawasaki disease is now a worldwide problem, which may cause life-threatening events in children. In Japan and in North America, Kawasaki Disease is the leading cause of acquired heart disease in children. Furthermore, it may develop into coronary artery disease in adults. To elucidate the etiology is essential. Long-term follow-up studies are also important.

> Email comments on this article to: JUNETA@CCT.bz

> > ~CCT~



Teiji Akagi, MD, PhD, FACC, FSCAI Associate Professor Cardiac Care Unit Okayama University Graduate School of Medicine and Dentistry 2-5-1, Shikata-cho Okayama, Japan 700-8558 Tel: 81-86-235-7682 Fax: 81-86-235-7683

t-akagi@cc.okayama-u.ac.jp

### MEDICAL CONFERENCES

Update on Cardiac Care in Neonates, Children, and Young Adults June 8-11, 2005; Shanghai, China www.congenitalcardiology.com/ 2005-ICU-Shanghai.pdf

PEDIRHYTHM-2 June 15-18, 2005; Antalya, Turkey www.pedirhythm.org

ASE 16th Annual Scientific Sessions June 15-18, 2005; Boston, MA, USA www.asecho.org

8th International Workshop Catheter Interventions in Congenital Heart Disease June 16-18, 2005; Frankfurt, Germany www.chd-workshop.org

5th International Pediatric Cardiovascular Symposium: Management of Complex Congenital Heart Disease from Infancy to Adulthood June 23-26, 2005; Amelia Island, FL, USA www.choa.org

12th World Congress on Heart Disease July 16-19, 2005; Vancouver, Canada www.cardiologyonline.com

SIS (Summer in Seattle) 2005 July 19-23, 2005; Seattle, WA, USA www.summerinseattle.com

New Concepts in Exercise and Pulmonary Hypertension in the Young: Their Interaction at Sea Level and Altitude

August 5-7, 2005; Aspen, CO, USA Taylor.Amy@tchden.org or www.congenitalcardiology.com/ Cardiac-Aspen2005.pdf

PICS-IX and ENTICHS-III (9th Pediatric Interventional Cardiac Symposium & Third Emerging New Technologies in Congenital Heart Surgery Sep. 15-18, 2005; Buenos Aires, Argentina www.picsymposium.com

The 4th World Congress of Pediatric Cardiology and Cardiac Surgery Sep. 18-22, 2005; Buenos Aires, Argentina www.pccs.com.ar/



### Saving Little Hearts, Inc.

5629 Barineau Lane • Knoxville, TN 37920 Dedicated to helping children with congenital heart defects and their families by providing financial and emotional assistance and educational information.

### www.savinglittlehearts.com

NORTH AMERICAN EDITION

### THE PEDIATRIC HEART NETWORK

By Gail D. Pearson, MD, ScD

When the National Heart, Lung, and Blood Institute (NHLBI) established the Pediatric Heart Network (PHN) in September 2001, it was the beginning of a new era in pediatric cardiovascular research. Relatively few multi-center trials had been undertaken previously, compared to the number in adult cardiovascular disease, and few pediatric academic centers had the infrastructure to conduct multiple trials simultaneously. This discouraging research landscape changed significantly with the advent of the PHN.

The mission of the PHN is to achieve public health advances for children, through the conduct and dissemination of collaborative research leading to evidence-based treatment options and improved outcome for pediatric patients with congenital and acquired heart disease.

The principal governing body of the Network is the Steering Committee, comprised of the 7 clinical center principal investigators, the data coordinating center principal investigator, and the Network Chair. The role of the Network Chair is to provide, with NHLBI, overall management and direction of Network activities.

The PHN had its origins in recommendations made to the NHLBI by pediatric cardiovascular researchers, and in the NHLBI Report of the Task Force on Research in Pediatric Cardiovascular Disease. To establish the PHN, the NHLBI released a Request for Applications – known as an RFA – to which there was a robust response. Applications to establish clinical centers and a data coordinating center were reviewed for technical merit by an independent group of experts convened specifically for this purpose. Clinical center applicants were judged on the scientific merit of the two clinical protocols proposed as well as on their ability to participate in a multi-center trials environment. Data coordinating center applicants were evaluated on the basis of their ability to coordinate complex clinical research, manage large databases, and provide analytical expertise in the execution of clinical protocols and analysis of resulting data. The decision as to which applications to fund, was made on the basis of technical merit, the National Heart, Lung and Blood Advisory Council's recommendations and funds available at the time.

As in other NHLBI Clinical Research Networks, standard oversight committees were established to assist the PHN. The Protocol Review Committee (PRC) is an independent group responsible for peer review of Network protocols. It is appointed by and reports to

# The Pediatric Heart Network (PHN) (with Principal Investigators)

### Network Chair:

University of Texas Southwestern Medical Center, Dr. Lynn Mahony

### **Clinical Centers:**

Boston Children's Hospital; PI, Dr. Jane Newburger

Children's Hospital of New York; PI, Dr. Welton Gersony

Children's Hospital of Philadelphia; PI, Dr. Victoria Vetter

Duke University Medical Center; PI, Dr. Page Anderson

Medical University of South Carolina; PI, Dr. Phillip Saul

Toronto Hospital for Sick Children; PI, Dr. Brian McCrindle

Utah Primary Children's Hospital; PI, Dr. LuAnn Minich

### Data Coordinating Center:

New England Research Institutes; PI, Dr. Lynn Sleeper

Table 1. The Pediatric Heart Network (PHN).

# AMPLATZER<sup>®</sup> Vascular Plug

### Simple. Reliable. Repositionable.

For more information visit: www.amplatzer.com or call: 763-513-9227 Toll free in the US: 888-546-4407



NORTH AMERICAN EDITION

**CONGENITAL** CARDIOLOGY TODAY

the NHLBI. Although the protocols proposed by the clinical center applicants are peer reviewed during the application process, protocols chosen for further development within the PHN may be considerably modified, and completely different protocols also may be developed. Therefore, an additional level of peer review provided by the PRC is an integral part of NHLBI Networks.

"The mission of the PHN is to achieve public health advances for children, through the conduct and dissemination of collaborative research leading to evidence-based treatment options and improved outcome for pediatric patients with congenital and acquired heart disease."

The other standing oversight committee is the Data and Safety Monitoring Board (DSMB). The DSMB is also an NHLBI-appointed independent group and is responsible for ensuring human subjects protections and data integrity as protocols are developed and during the conduct of clinical studies. Once studies are under way, a major role of the DSMB is to review adverse events, recruitment, interim findings, and protocol violations. The recent addition of a Medical Monitor to the Network will augment the review of adverse events and any study endpoints that require adjudication.

With this infrastructure in place, the PHN Steering Committee began the

arduous task of deciding what research to conduct in September 2001. The Steering Committee began by hearing presentations of the 14 protocols that its members proposed in their grant applications, and soon established a system of protocol evaluation including criteria such as benefit to pediatric patients, scientific merit, and feasibility. After careful deliberation, the Steering Committee settled on two initial protocols: The Relationship Between Functional Health Status and Laboratory Parameters of Ventricular Performance After the Fontan Procedure, and the Trial of Pulse Steroid Therapy in Kawasaki Disease.

The Fontan study was designed as a cross-sectional study to determine the interrelationships between health status and measures of cardiac performance in children 6-18 years old who have undergone a Fontan procedure. The Steering Committee originally set out to design a treatment trial in the Fontan population, but soon discovered that there were not enough published data to select clinically relevant endpoints. The goal of the Fontan study, therefore, was to acquire such data. When enrollment ended in 2004. 546 patients were enrolled. These patients underwent multiple evaluations, including exercise testing, MRI measures of ventricular function and



Figure 1. PHN Logo

"NHLBI will provide funds to support the Pediatric Heart Network for another five years, beginning in September 2006."

mass, echocardiographic assessment of ventricular mechanics, cardiac peptides, and health status using standard questionnaires for children, adolescents, and their families. Central interpretation of echocardiographic, MRI, and serology data was conducted. Several abstracts from this study have already been presented at national meetings, and the main results and additional papers are in preparation.

The Kawasaki trial was undertaken to determine whether a single high dose of methylprednisolone vs. placebo, on the background of standard therapy, affects coronary artery diameter at 6 weeks post-randomization, normalized for body surface area. Secondary endpoints include duration of fever and hospital stay, C-reactive protein levels, and adverse events. This trial also included central interpretation of echocardiographic and serology data. In 2 years, nearly 200 patients were randomized, with recruitment completed in December 2004. Data cleaning and analysis are underway, with the primary results expected soon.

Once these trials were launched, the PHN turned its attention to developing additional protocols. The ability to conduct multiple studies simultaneously is a strength of the Network infrastructure, but it also requires constant mental gear-changing and considerable hard work. To date, the Steering Committee has developed four more proto-



# The Fourth World Congress of Pediatric Cardiology and Cardiac Surgery

Buenos Aires, Argentina • September 18-22, 2005 www.pccs.com.ar

NORTH AMERICAN EDITION

JUNE 2005 PAGE 7

cols. Two are currently recruiting patients, and the remaining two are set to begin recruitment within the next month. These studies are:

- Trial of ACE Inhibition in Infants with Single Ventricle. The purpose of this trial is to determine the effect of ACE inhibition (enalapril) on somatic growth in infants with single ventricle physiology who have undergone the first staged surgical procedure.
- Trial of ACE Inhibition in Children with Mitral Regurgitation after Repair of an AVSD. This study is evaluating the safety and efficacy of enalapril therapy after repair of an atrioventricular septal defect in children who have at least moderate mitral valve regurgitation.
- Single Ventricle Reconstruction Trial. In this trial, the standard Norwood approach for neonates with hypoplastic left heart syndrome and variants will be compared to the right ventricle-topulmonary artery conduit surgical strategy.
- Ventricular Volume Variability Trial. This is a study of variability in echo measures of ventricular performance, assessed within patients, across time, and between echocardiographers. The purpose is to estimate, in a pediatric cardiac population, the variance of change in ventricular function measures to use for future clinical trial sample size calculations.

In addition to the benefits of the specific research performed, the Network has a number of other advantages. Participation in Network activities and meetings is providing experience in clinical research for many young nurse and physician investigators. PHN protocols offer fertile ground for ancillary studies, which can be proposed by investigators both within and outside the Network. The success of the Network is also fostering increased interest in multi-center clinical trials in pediatric cardiovascular disease.

NHLBI will provide funds to support the Pediatric Heart Network for another five years, beginning in September 2006. The RFA for the next funding cycle was published in early March, and can be found at: http:// grants.nih.gov/grants/guide/rfa-files/ RFA-HL-05-010.html. Applications are due September 23, 2005.

> Email comments on this article to: JUNEGDP@CCT.bz

> > ~CCT~



Gail D. Pearson, MD, ScD, FAAP, FACC, FAHA Leader, Heart Development, Function and Failure Scientific Research Group Division of Heart and Vascular Diseases National Heart, Lung, and Blood Institute National Institutes of Health/Department of Health and Human Services

pearsong@mail.nih.gov

Optimized for

Pediatrics

### RFA-HL-05-010 "Pediatric Heart Network"

This RFA solicits applications in an open competition for clinical centers and a data coordinating center for the next funding cycle of the Network. Current centers and new centers are both invited to apply.

### **Key Dates:**

- Letter of Intent Due: 8/26,/2005
- Applications Due: 9/23/2005
- Peer Review: February-March, 2006
- NHLBI Council Review: 6/5/2006
- Expected Start Date: 9/1/2006

# Direct questions about scientific/ research issues to:

Gail Pearson, M.D., Sc.D. Division of Heart and Vascular Diseases NHLBI 301-435-0510 pearsong@mail.nih.gov

Direct questions about financial or grants management matters to:

Ms. Marsha Mathis Grants Operations Branch NHLBI 301-435-0170 mathism@nhlbi.nih.gov

For complete details see http:// grants.nih.gov/grants/guide/noticefiles/NOT-HL-05-116.html



# MyLab30 5 PERFORMANCE AND PORTABILITY WITHOUT COMPROMISE

The Ultrasound Unit . High Frequency Phased Array Transducers

- All Modalities: 2D, PD&CW Doppler, CFM
- Superb Onboard Digital Management
- Weighs less than 20 lbs.



www.CongenitalCardiologyToday.com

## PERCUTANEOUS CLOSURE OF A RESIDUAL PERIMEMBRANOUS VENTRICULAR SEPTAL DEFECT AFTER SURGICAL REPAIR

By Carlos AC Pedra, MD; Sérgio C. Pontes, Jr., MD; Simone R.F. Pedra, MD; Juliana Neves, MD; M. Aparecida P. Silva, MD; M. Virginia T. Santana, MD; Valmir F. Fontes, MD

### Introduction

Isolated perimembranous (PM) ventricular septal defect (VSD) is one of the most common congenital cardiac malformations.<sup>1</sup> Surgery has been performed safely and effectively and has been regarded as the gold standard method of treatment for this disease.<sup>1,2</sup> Although residual leaks are observed in up to 5-10% of cases, most of them are restrictive and well tolerated.<sup>1,2</sup> Occasionally. they can result in significant left-to-right shunting with persistent left ventricular volume overload, which requires reintervention.<sup>2</sup> Recent publications have reported the feasibility, safety and efficacy of percutaneous closure of native PM VSDs with the use of the Amplatzer membranous VSD occluder (AGA Medical Corporation, Golden Valley, Minnesota).3-8 Controlled release coils made of a reinforced Nitinol wire (PFM, Cologne, Germany) has also been employed for transcatheter closure of such defects with encouraging results.9,10 In this paper, we describe a case in which a significant residual leak after surgical repair of a PM VSD was closed using transcatheter techniques.

### **Case report**

This patient was a non dysmorphic 4 year-old boy (weighing 15 kgs) with a large PM VSD who was referred from the countryside to our institution for surgical

repair. He was symptomatic and receiving digoxin, furosemide and captopril. Classical findings of significant PM VSD with pulmonary arterial hypertension were encountered on physical examination, chest radiograph and ECG. Transthoracic echocardiography revealed a large PM VSD with inlet extension, measuring 14 mm at its maximal dimension (Figure 1A). There was significant left atrial and ventricular volume overload. Pulmonary arterial pressure was estimated at systemic levels. The child was taken to the catheterization laboratory for further hemodynamic assessment. A routine left and right catheterization was carried out under general anesthesia. Hemodynamics showed the



Figure 1. Echocardiographic pictures of the defect. (A) Four chamber view (pre-operative TTE): Large PM VSD with inlet extension measuring 14 mm at its maximal diameter. (B) Modified four chamber view (post-operative TTE): There is a residual VSD (red arrow) measuring 5-6 mm at the superior portion of the patch (light blue arrow), near the crux of the heart and the AV valves. (C) Long axis view (TEE during the percutaneous procedure): The left ventricular loops of the PFM Nitinol coil are protruding into the LVOT. (D) Long axis view (TEE during the percutaneous procedure): The Amplatzer membranous VSD occluder is well positioned after final release. There is a small residual leak through the superior aspect of the device.

# Ask the Editorial Board

Would you like to ask the Editorial Board's opinion about a procedure, technology or product? Email your question to: ASK@CongenitalCardiologyToday.com

Selected questions and answers may be published in upcoming issues. Names will be withheld upon request.

following pressures (in mm Hg): RA: 5; RV: 88/6; MPA: 88/31 (mean 58); Ao: 90/58 (mean 71); LV: 90/12. Pulmonary vascular resistance (PVR), Qp/Qs and PVR/SVR were estimated at 3.6 Wood Units, 2.9 and 0.23, respectively. After the administration of NO (40 PPM), PVR was 1.6 U Wood X m-2, Qp/Qs 4.9 and RVP/RVS 0.11. Left ventriculogram in long axial view showed a large PM VSD, measuring 13 mm at its maximal diameter (Figure 2A). Pulmonary arterial angiogram demonstrated progressive tapering of the pulmonary arteries with satisfactory opacification of the peripheral vessels. The capillary phase was homogeneous and there was rapid return of the contrast media to the LA. The child was subsequently referred to surgery, which was carried out under cardiopulmonary bypass using standard techniques.<sup>2</sup> A pericardial patch was sutured at the edges of the defect with pledgets. The tricuspid valve was not removed for patch closure. Post-operatively, signs of persistent left-to-right shunting were detected invasively (SVC sat: 65%; PA sat: 82%) and confirmed by echocardiography. A 5-6 mm residual VSD was observed through the upper portion of the patch near the AV valves (Figure 1B). On the 8th post-op day the child was reoperated, however, the surgeon was unable to identify the site of the residual leak intra-operatively. After a month, a decision was made to attempt closing the residual leak using transcatheter techniques. Informed consent was obtained from parents. Under general anesthesia, vascular access was obtained with placement of a 5 Fr sheath in the left femoral artery and a 7 Fr sheath in the right femoral vein. Heparin sulfate (150 IU/kg) and Cefazolin (30 mg/kg) were given. Hemodynamics showed the following pressures (in mm Hg): RA: 8; RV: 50/10; MPA: 50/18: Ao: 90/50: LV: 90/15. The Qp/Qs was estimated at 1.8:1. Left ven-



Figure 2. Ventriculograms in long axial or hepatoclavicular views displaying the defect (A) Pre-operative findings: Large PM VSD with inlet extension and conspicuous left-to-right shunt, opacifying the dilated MPA. (B) Post-operative findings during the percutaneous procedure: Moderate left-to-right shunt through the superior portion of the patch, near the AV valves. The residual defect measured 5 mm on angiography. (C) The PFM Nitinol coil remained in a sub-optimal position with the left loops protruding into the LVOT, distant from the patch. (D) Adequate position of the Amplatzer membranous occluder immediately after release. There is a small residual leak at its superior portion.

triculogram confirmed the echocardiographic findings (Figure 2B). Using previously described techniques and under transesophageal echocardiographic guidance (TEE), the residual VSD was crossed retrogradely and an arterial-

# AMPLATZER Septal Occluder

# R

### Simple. Reliable. Repositionable.

For more information visit: **www.amplatzer.com** or call: 763-513-9227 Toll free in the US: 888-546-4407



venous loop established using a 260 cm Glide wire (Terumo Cardiovascular Systems Corporation, Ann Harbor, MI). A 7 Fr long, braided sheath (Flexor; Cook Cardiology, Bloomington, IN) was advanced to the ascending aorta across the VSD from the vein. The latest version of a pre-mounted PFM coil (with reinforced Nitinol wire and Dacron fibers) (12 X 6 mm) was advanced through the long sheath until the tip of the delivery catheter was about 1 cm out of the long sheath. By pushing the core wire, the loops of the coils were exteriorized in the ascending aorta with the last 2 loops remaining inside the delivery catheter. The whole system was carefully brought back as a unit across the aortic valve until the loops abutted the interventricular septum. Keeping a gentle traction on the system and the tip of the long sheath close to the tip of the delivery catheter, the last 2 loops were delivered in the RV by withdrawing the delivery catheter and pushing the core wire. Final position of the coil was assessed by TEE (Figure 1C) and angiography (Figure 2C). Because the left ventricular loops protruded into the LVOT and there was significant residual leak, the device was recaptured inside the long sheath and removed out of the body. A decision was then made to attempt closing the defect using an 8 mm Amplatzer membranous VSD occluder (AGA). The VSD was again crossed in a retrograde fashion and an arteriovenous loop established using a Rope wire (AGA) as described in previous published protocols (6-8). An 8 Fr long sheath (TorqVue, AGA) was advanced to the ascending aorta and subsequently positioned near the left ventricular apex as described before.(6-8) After removal of the Rope wire and dilator, the device was pushed through the long sheath and the left ventricular disc deployed within the left ventricular cavity with the radiopaque marker pointing downwards. The whole system was retracted as a unit until the left disc touched the patch. By pulling the long

sheath and advancing the delivery cable, the right disc was deployed on the right side. Good device position was confirmed by both TEE and angiography. Aortic and AV valve function were preserved. After device release, a tiny residual leak was seen through the superior portion of the device (Figure 1D and 2D). The child was awaken in the catheterization laboratory and had an uneventful recovery. He was discharged home the following day on aspirin (5 mg/kg/day). A transthoracic echocardiogram revealed complete closure of the defect after a month. There was no AI or significant TR. The child remained in sinus rhythm with no signs of left or right bundle branch block. Medications were gradually discontinued.

### Discussion

A significant residual VSD after surgical repair may occasionally require reoperation, resulting in increased morbidity and hospital stay. Although successful transcatheter closure of PM VSDs has been described recently,(3-8) there is limited experience with this approach for such residual defects. Since it has the potential to avoid a repeat cardiopulmonary bypass run, it may well be a safer therapeutic option. However, the issue of what type of residual defect that is amenable to transcatheter closure, including location, number and size, has yet to be clarified. This will only come with ongoing experience. In the case described herein, the residual defect was single and located at the superior edge of the patch, towards the crux of the heart. Despite being close to the AV valves, we felt there was enough room surrounding the defect to accommodate a device without interfering with AV valve function. Moreover, we decided to attempt the transcatheter approach because the child had undergone surgical repair twice unsuccessfully.

In regards to the types of devices used in this case, the PFM coil was employed initially under a study protocol to assess its safety and efficacy. Even acknowledging that clinical experience with it is still very limited 9,10 (about 20 implantation procedures with the latest version; unpublished data; Dr. Trong-Phi Le, personal communication), it has theoretical and potential advantageous features. It requires lower profile sheaths (6 or 7 Fr) for implantation; due to the flexible central portion of the device, it can be exteriorized in the ascending aorta and pulled back safely across the aortic valve, with no need to place the long sheath near the left ventricular apex; and it is secured within the defect without exerting radial forces. Although it seems to work well for small to moderate defects, especially those associated with aneurysm formations, it may be unsuitable for larger defects. In the case described herein, the PFM coil remained in an inadequate position, protruding into the LVOT, probably because it approached the septum (patch) from above, coming from the aorta. If we had delivered the LV loops close to the left ventricular apex, it might have been possible to engage it within the patch, in a more favorable course towards the defect itself. Possible entanglement with the mitral valve apparatus could be an issue employing this approach. Coil recapturing and removal was feasible and safe, keeping in mind that the long sheath had to be kept close to the delivery catheter tip to avoid entanglement with the tricuspid valve apparatus. Subsequent use of the Amplatzer membranous occluder was a natural choice. Initial clinical experience with this device has been encouraging.<sup>3-8</sup> The rate of complete closure is high (>90%) and aortic and tricuspid valve function are preserved, at least in the short-to-mid term.<sup>3-8</sup> However, complete heart block seems to occur with an incidence of 1-2%.<sup>11</sup> Whether this is related to the radial forces exerted by the central waist of the device onto the defect edges or



Fifth International Pediatric Cardiovascular Symposium: Management of Complex Congenital Heart Disease From Infancy to Adulthood The Ritz-Carlton, Amelia Island, June 23-26, 2005

www.choa.org/forprofessionals/cme or call Nancy Richardson 404-785-7843 or Kathy Murphy 404-785-6480

due to the endothelialization process is speculative. Other factors such as younger age, inlet extension of the VSD and crossing the VSD from the RV side may play a role in the development of this complication. In our experience, complete heart block did occur between 3-6 months after the procedure in a single case out of 33 implantation procedures (unpublished personal data). In the case presented herein, the Amplatzer device worked nicely because it approached the patch at a proper angle, coming from the LV apex. Besides, the longer inferior portion of the LV disc engaged well within the patch with the waist remaining inside the residual defect itself, which was probably responsible for complete closure. Improvements on the design of this device and delivery system have been made by placing a female screw in the center of the left ventricular disc and a male screw at the tip of the rope wire.<sup>12</sup> This enables to position the device more precisely because of the through-andthrough cable-to-device-to-rope wire system, creating traction from both the venous and arterial sides. We anticipate that this system can be useful in difficult cases. such as some residual VSDs after surgical repair.

In conclusion, percutaneous closure of a residual PM VSD after surgical repair using an Amplatzer membranous device was feasible, safe and effective in the selected patient presented herein. More experience is warranted before the widespread use of this technique is recommended.

### References

1. Tynan M, Anderson R.H. *Ventricular septal defect.* In: Anderson RH, Baker EJ, Macartney FJ, Rigby ML, Shinebourne EA, Tynan M, editors. Paediatric Cardiology. London: Churchill Livingstone, 2002: 983-1014.

2. Kouchoukos NT, Blackstone EH, Doty DB, Hanley FL, Karp RB. *Ventricular septal* 

*defect.* In: Kouchoukos NT, Blackstone EH, Doty DB, Hanley FL, Karp RB, editors. Kirklin/Barrat-Boyes: Cardiac Surgery. Philadelphia: Churchill Livingstone, 2003: 850-910.

3. Hijazi ZM, Hakim F, Haweleh AA, Madani A, Tarawna W, Hiari A et al. *Catheter closure of perimembranous ventricular septal defects using the new Amplatzer membranous VSD occluder: initial clinical experience.* Catheter Cardiovasc Interv 2002; 56(4):508-515.

4. Thanopoulos BD, Tsaousis GS, Karanasios E, Eleftherakis NG, Paphitis C. *Transcatheter closure of perimembranous ventricular septal defects with the Amplatzer asymmetric ventricular septal defect occluder: preliminary experience in children.* Heart 2003; 89(8):918-922.

5. Bass JL, Kalra GS, Arora R, Masura J, Gavora P, Thanopoulos BD et al. *Initial human experience with the Amplatzer perimembranous ventricular septal occluder device.* Catheter Cardiovasc Interv 2003; 58(2):238-245.

6. Pedra CA, Pedra SRFF, Esteves CA, Pontes Jr SC, Braga SLN, Arrieta SR et al. *Percutaneous closure of perimembranous ventricular septal defects with the Amplatzer device: technical and morphological considerations.* Catheter Cardiovasc Interv 2004;61: 403-410.

7. Pedra CAC, Pedra SRF, Esteves CA, Chamie F, Christiani LA, *Fontes VF. Transcatheter closure of perimembranous ventricular septal defects.* Exp Review Cardiovasc Therapy 2004; 2 (2): 253-264.

8. Hijazi ZM. *Device closure of ventricular septal defects*. Catheter Cardiovasc Interv 2003; 60(1):107-114.

9. Le TP, Vaessen P, Freudenthal F, Grabitz RG, Sievert H. *Transcatheter closure* of subaortic ventricular septal defect (VSD) using a nickel-titanium spiral coil

The Ultrasound Unit

Optimized for

Pediatrics

### Endowed Chair, Division of Pediatric Cardiology St. Louis University School of Medicine Cardinal Glennon Children's Hospital

St. Louis University, a Catholic, Jesuit institution dedicated to student learning. research, health care, and service is seeking candidates for a faculty position in the Department of Pediatrics at the Associate Professor/Professor rank as the Director of the Division of Cardiology at Cardinal Glennon Children's Hospital. A new Endowed Chair of Pediatrics will be established by Saint Louis University for the successful candidate. The Division Director will have opportunities to recruit both clinical and research faculty, to plan significant programmatic and facilities enhancements, and to develop a program of clinical and translational research. The candidate should have a proven record of academic excellence as well as a commitment to excellence in patient care and medical education.

Cardinal Glennon Children's Hospital is a 160-bed free standing hospital located in midtown Saint Louis, adjacent to Saint Louis University. It is currently in the midst of a 2-phase, \$100 million expansion and renewal program. The Hospital serves a diverse population from the inner city, the metropolitan area, and a 200-mile referral radius. The medical staff includes over 90 full-time St. Louis University School of Medicine faculty.

Interested candidates should forward correspondence and curriculum vitae to Robert E. Lynch, MD, PhD, Chair, Cardiology Search Committee, Professor and Director, Pediatric Critical Care, Department of Pediatrics, Saint Louis University School of Medicine, 1465 South Grand Blvd, St. Louis, MO 63104. Tel: (314)577-5395; Fax: (314) 268-6459; email: lynchre@slu.edu

Saint Louis University is an Affirmative Action, Equal Opportunity Employer, and encourages nominations of and applications from women and minorities.



# MyLab30 & PERFORMANCE AND PORTABILITY WITHOUT COMPROMISE

High Frequency Phased Array Transducers

All Modalities: 2D, PD&CW Doppler, CFM

- Superb Onboard Digital Management
- Weighs less than 20 lbs.



www.CongenitalCardiologyToday.com

© Copyright 2005, Congenital Cardiology Today. All rights reserved

(*NitOcclud*): animal study and initial clinical results. Progr Pediatr Cardiol 2001; 14: 83-88.

10. Ewert P, Kretschmar O, Peters B, Abdul Khaliq H, Nagdyman N, Schulze-Neick I et al. *Transcatheter closure of congenital ventricular septal defects*. Z Kardiol 2004; 93 (2): 147-55.

11. Holzer R, Hijazi ZM, Wilkinson J, DeGiovanni J, Hakim F, Al-Hroob A et al. *Transcatheter closure of perimembranous ventricular septal defects using the Amplatzer Membranous VSD Occluder: immediate results and mid-term results.* Circulation. 2004; 110: III-565.

12. Amin Z, Danford D. *Modification of Amplatzer device and delivery system to improve precise placement in the ventricular septal defects.* Catheter Cardiovasc Interv 2003; 60 [1]: 125.

Email comments on this article to: JUNECP@CCT.bz

~CCT~



Corresponding Author:

Carlos AC Pedra, MD Director, Catheterization Laboratory for Congenital Heart Disease Instituto Dante Pazzanese de Cardiologia São Paulo, SP, Brazil Tel: (55) (11) 5085-4114 Fax: (55) (11) 5085-4161

cacpedra@uol.com.br carlosacpedra@hotmail.com

### SEPTEMBER CONFERENCE FOCUS

PICS/ENTICHS- 2005 (Pediatric Interventional Cardiac Symposium and Emerging New Technologies in Congenital Heart Surgery) September 15-18, 2005; Hilton Buenos Aires Buenos Aires, Argentina www.picsymposium.com

PICS/ENTICHS-2005 will focus on the latest interventional catheter strategies and emerging technical advances in cardiac surgery for fetuses, children, and adults with congenital heart disease:

- Daily live case demonstrations including the latest technologies in devices, implantable valves, stents, balloons and more.
- Special didactic sessions including closure of all types of septal defects, percutaneous
- Live surgical cases focusing on hybrid intervention.
- Meet the Expert Sessions.
- "My Nightmare in the Cath Lab" case presentations.

The second annual meeting of The Congenital Cardiovascular Interventional Study Consortium (CCISC) will take place at PICS/ENTICHS 2005

**Course Directors:** Drs. Ziyad Hijazi; Emile A. Bacha; William E. Hellenbrand; and John P. Cheatham.

**Course Co-directors:** Drs. Horacio Faella; Mark Galantowicz; Miguel Granja; and Christian Kreutzer.

**Guest Faculty Includes:** Drs. Teiji Akagi; Luis Alday; B.G. Alekyan; Mazeni Alwi; Zahid Amin; Luigi Ballerini; John Bass; Lee Benson; Felix Berger; David Bichell; Philipp Bonhoeffer; Redmond Burke; Mario Carminati; Chin Chan; Qi-Ling Cao; Bharat Dalvi; Cesar Esteves; J.V. De Giovanni; Makram Ebeid; Jeffrey Feinstein; Craig Fleishman; Valmir Fontes; Thomas Forbes; Ronald Grifka, MD; Felipe Heusser; Frank Ing; Thomas Jones; Charles S. Kleinman; Krishna Kumar; Michael Landzberg; Trong-Phi Le; Geoffrey K. Lane; Larry Latson; Jose Suarez de Lezo; Achi Ludomirsky; Jozef Masura; Constantine Mavroudis; Joaquim Miro; John W. Moore; Charles E. Mullins; David Nykanen; Eustaquio Onorato, MD; Carlos Pedra; Jean-Francois Piechaud, MD; Alejandro Peirone; Shakeel A. Qureshi, MD; Wolfgang Radtke; P.S. Rao, MD; Mark Reisman; Carlos E. Ruiz; Satinder Sandhu; Horst Sievert; Basil (Vasilios) Thanopoulous; Hideshi Tomita; Alejandro Torres; Wayne Tworetzky, MD; Michael Tynan; Ross Ungerleider; Kevin Walsh; Gilbert Wernovesky; James Wilkinson; Neil Wilson; Carlos Zabal; and Evan Zahn.

For more details, abstract submission and registration, please visit the website.

Accreditation: The University of Chicago Pritzker school of medicine is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to sponsor continuing medical education for physicians



# PICS/ENTICHS- 2005

Pediatric Interventional Cardiac Symposium and Emerging New Technologies in Congenital Heart Surgery SEPTEMBER 15-18, 2005 At the Hilton Buenos Aires Buenos Aires, Argentina Immediately preceding the 4th World Congress of Pediatric Cardiology and Cardiac Surgery

www.picsymposium.com

### **MEDICAL NEWS AND INFORMATION**

### Study Points to Link Between Risks of Congenital Heart Defects and Alterations in Maternal Plasma Biomarkers

Findings from a study lead by Dr. Charlotte Hobbs, director of the Arkansas Children's Hospital Research Institute's Center for Birth Defects Research and Prevention, point to the link between babies with congenital heart defects and the presence of certain plasma biomarkers.

The study measured biomarkers of the folate-dependent methionine/homocysteine pathway among 314 women. Two hundred and twenty four of the subjects had children with congenital heart defects, and 90 had children who were unaffected. All of the women who had children with heart defects had higher average concentrations of homocysteine and lower average concentrations of methionine.

The findings were published in the January 2005 issue of American Journal of Clinical Nutrition, (www.ajcn.org) and while they will still need to be verified through subsequent tests, this study is the first to show the association between these biomarkers and congenital heart defects.

"This is important, because it is the first step in identifying a specific metabolic profile in women who have an increased risk of producing children with congenital heart defects," Hobbs said. 'If we can do that, we may be able to identify women who are at higher risk of having pregnancies affected by congenital heart defects before they even conceive. We could then prescribe a targeted clinical intervention that would increase their chances of having a healthy pregnancy."

Congenital heart defects are the most prevalent birth defect in the United States where 11 in 1,000 children are born with some form of the condition. While it is the most common birth defect, it also is one of the least understood. Only about 15% of heart defects can be attributed to a known cause.

At this time, the best tool to manage these defects is for

an expectant mother to have an ultrasound during her second trimester.

"Even with this tool, we can only identify that the problem exists and make plans for the birth to occur in a center where the defect can be treated immediately," Hobbs said. "These new findings could be used to prevent the defect from occurring in some cases."

The study is funded through a five-year, \$4.5 million grant from the National Institute of Child Health and Human Development. Dr. Hobbs is an associate professor Pediatrics, University of Arkansas for Medical Sciences College of Medicine.

For more information: www.archildrens.org or www.ajcn.org

U.S. Patent & Trademark Office Issues Patent No. 6,830,584 to CoreValve for Its 'Device for Replacing a Cardiac Valve by Percutaneous Route'

PARIS, FRANCE & IRVINE, CA, USA.--(BUSINESS WIRE)

CoreValve S.A. has announced that the United States Patent & Trademark Office has issued Patent No. 6,830,584 to CoreValve, thus protecting the Company's 'device for replacing a cardiac valve by percutaneous route'--i.e., the CoreValve Percutaneous ReValving System.<sup>™</sup>

With ReValving,<sup>™</sup> heart valve replacement can be performed 'non-surgically' in a cardiac 'cath lab' just like angioplasty and stenting, resulting in less trauma to the patient and substantial cost-savings to the healthcare system compared to open-heart surgery to replace diseased heart valves.

"U.S. patent protection for ReValving is critically important, of course--particularly in light of the fact that we already have clinically established that ReValving is feasible to non-surgically treat the widest possible range of diseased heart-valve patients, including traditional surgi-

# Sponsored By BBRAUN

For information, please call 1-800-BBRAUN2 (227-2862) or visit www.bbraunusa.com

# Making a difference, one patient at a time.

Are you looking for a well-respected technologically advanced health care system with the added henefit of a nationally recognized research facility? That's exactly what you'll find at Marshfield Clinic's 40 centers in communities throughout scenic central, western and northern Wisconsin.

Marshfield Clinic is a 700+ physician-directed multi-specialty group practice with over 80 medical specialties. Currently, we're seeking a third BC/BE Pediatric Cardiologist to join its tertiary facility in Marshfield, Wisconsin. The right candidate will join a large pediatric department of general and subspecialty trained pediatricians. Opportunities exist for teaching and research with our pediatric residency and the Marshfield Clinic Research Foundation. A 504-bed acute care facility with a Level I PICU and Level III NICU adjoins the clinic on the medical campus.

Marshfield Clinic offers its physicians an excellent salary and benefits, along with the opportunity for personal and professional growth. Interested candidates may contact either of our pediatric cardiologists, Dr. Kathy Finta or Dr. Julie Dietz at (800) 782-8581 to learn more, or may submit their CV and questions to: Mary Treichel, Physician Recruiter, **Marshfield Clinic**, 1000 N. Oak Avenue, Marshfield, WI 54449, or call (800) 782-8581 extension 19774, or fax to (715) 221-9779; E-mail: treichel.mary@marshfieldclinic.org

Check out our web site to learn more about Marshfield Clinic at: www.marshfieldclinic.org/recruit



Marshfield Clinic is an Affirmative Action/Equal Opportunity employer that values diversity. finorities, females, individuals with disabilities and veterans are encouraged to apply. Sony, not a health professional shortage area cal-candidates," said Jacques Seguin, MD, chairman and CEO of CoreValve. "Indeed, CoreValve's ReValving approach has potential universal applicability to percutaneously treat the two most prevalent diseases of the aortic valve--stenosis and regurgitation," added Professor Seguin.

### About CoreValve

Privately held CoreValve, S.A., headquartered in Paris, France, has developed a proprietary delivery system for percutaneous heart valve replacement, based on a novel catheter-and-self-expanding-stent approach on a beating heart, thus avoiding open-heart surgery. The CoreValve procedure--with the proprietary CoreValve Percutaneous ReValving System<sup>™</sup> --can be performed in a cardiac "cath lab" just like angioplasty and stenting, resulting in less trauma to the patient and substantial cost-savings to the healthcare system.

For more information: www.corevalve.com

Rare childhood genetic syndrome identified: Multiple problems include cardiac arrhythmias and atypical autism

Researchers at Children's Hospital Boston, Howard Hughes Medical Institute and the University of Utah have identified a rare, previously undiscovered genetic syndrome that is often fatal by the second year of life, but which may be treatable with calcium channel-blocking drugs. Findings were reported in the October 1, 2004 issue of the journal, Cell.

The disease, named Timothy Syndrome after one of the paper's authors, is characterized by a variety of problems including heart arrhythmias, congenital heart abnormalities, webbed hands and feet, a weakened immune system, cognitive abnormalities, and, surprisingly, autism. The researchers have identified 17 children with the syndrome, seven of whom were living.

Despite the complexity and severity of Timothy Syndrome, the researchers show that it arises from a single, spontaneous, very subtle gene mutation in the mother's egg or father's sperm - substitution of a single base pair. The reason so many body systems are affected is that mutation impairs a very fundamental molecule, a type of calcium channel, that is found in many tissues and organs.

# Do You Want to Recruit a Pediatric Cardiologist?

Advertise in the only monthly publication totally dedicated to pediatric and congenital cardiology. For more information: **Recruitment@CCT.bz** 

### CONGENITAL CARDIOLOGY TODAY

NORTH AMERICAN EDITION

Calcium channels control how much calcium can get inside a cell. Calcium is one of the body's most important signaling molecules, and normally, cellular calcium levels are tightly regulated. Dr. Mark Keating, senior author of the study and a Howard Hughes Medical Institute investigator at Children's Hospital Boston, likens the calcium channel to a screen door.

"After you go through the screen door, it automatically closes," he says. "This mutation dismantles the automatic closing mechanism, so the door just stays open."

As a result, cells are overwhelmed by an influx of calcium. Because calcium-channel blocking drugs can ameliorate calcium overload, these medications may be useful for treating arrhythmia and cognitive deficits in individuals with Timothy Syndrome, Keating says.

Experiments also showed that the gene encoding the calcium channel was active not only in heart muscle cells, but in tissues of the gastrointestinal system, lungs, immune system, smooth muscle, testes, and brain - including brain regions that are known to show abnormalities in autism. Keating notes, however, that autism is a complex disorder with many different causative factors.

The study was led by Igor Splawski, PhD, in the Cardiovascular Research Division at Children's Hospital Boston in collaboration with the University of Utah and the Boston University School of Medicine. The researchers will continue to treat patients with Timothy Syndrome and evaluate their response to calcium-blockers. They will also continue to look for arrhythmia genes and other calcium channels that might be involved in arrhythmia, and try to determine whether this calcium channel is involved in other forms of autism.

Children's Hospital Boston is the nation's leading pediatric medical center, the largest provider of health care to Massachusetts' children, and the primary pediatric teaching hospital of Harvard Medical School.

For more information: www.childrenshospital.org

### Submit an Article to Congenital Cardiology Today

Send an email with your name, title, organization, address, and email along with the title and a 50 word description to:

### Articles@CCT.bz



Positions Available

# PEDIATRIC CARDIOLOGISTS Phoenix, Arizona

We are seeking two BC/BE Pediatric Cardiologists to join a 12-member group practice with offices in the Phoenix and Tucson metropolitan areas. For the Phoenix area practice, we are recruiting two generalists with experience in fetal, transthoracic and transesophageal echocardiography. In 2004, this premier group performed over 400 catheterization cases, over 400 fetal echocardiograms and close to 12,000 echocardiograms, (over 5,000 were done in the office). The practice is engaged in clinical research and covers teaching rotations for residents and medical students.

While living in Phoenix enjoy 325 days a year of sunshine, 200 picturesque golf courses, world-class dining, exhilarating outdoor adventure, trendy shopping and enriching culture.

Receive a competitive income plus outstanding benefits including health, life and disability insurances, paid malpractice insurance and generous CME allowance.

For more information on these exciting opportunities please contact Lori Abolafia, Physician Relations Specialist e-mail: lori\_abolafia@pediatrix.com

> Pediatrix Medical Group 1301 Concord Terrace Sunrise, FL 33323

877.456.8686 toll free

877.780.4242 toll free fax www.pediatrix.com



### The SONOS 7500 system with Live 3D Echo ... for better visualization of congenital heart defects

Learn more at: www.medical.philips.com/live3D or call 800-229-6417



www.CongenitalCardiologyToday.com

9008 Copenhaver Drive, Suite M Potomac, MD 20854 USA

# Your FREE June Issue of CONGENITAL CARDIOLOGY TODAY

© 2005 by Congenital Cardiology Today (ISSN 1554-7787-print; ISSN 1554-049-online). All rights reserved. Photocopying, reproduction or quotation either in full or in part is strictly prohibited without the written permission of the publisher. For permission, send an e-mail to: *Permission* @*CCT.bz.* Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and are not necessarily the views of Congenital Cardiology Today. Postmaster: Send address changes to Congenital Cardiology Today, 9008 Copenhaver Dr., Suite M, Potomac, MD 20854 USA

# **CONGENITAL** CARDIOLOGY TODAY<sup>®</sup>

RELIABLE INFORMATION IN CONGENITAL CARDIOLOGY™

Article Submission Article @CCT.bz Information Info @CCT.bz

Medical Editor/Medical Editorial Board John W. Moore, MD, MPH, FACC *jwmoore* @*mednet.ucla.edu* 

**Contact the Editorial Board** See page 2 for a complete list of Editorial Board members and how to contact them.

New Product Submission PR@CCT.bz

Article Reprint Information Reprint@CCT.bz

Web Site Information Web@CCT.bz

Symposium Submission Symposium@CCT.bz

Medical Website Submission URL@CCT.bz

How to Reference CCT Reference@CCT.bz

### **FREE Subscription**

Congenital Cardiology Today<sup>®</sup> is published monthly and FREE to qualified professionals worldwide in congenital and pediatric cardiology, and related fields.

Print subscriptions are available in the U.S. and Canada for the North American edition.

Electronic subscriptions (in Adobe PDF format) are available in the International edition. To subscribe to either the International or North American edition, send an email with your name, title, organization, phone, address and e-mail to: *Subs*@*CCT.bz.* Or fax your request to +1.240.465.0692

Advertising Production Production@CCT.bz

Letters @CCT.bz

Sales & Marketing Sales @CCT.bz

Recruitment/Fellowship Advertising Recruitment@CCT.bz

### **Publishing Management**

Tony Carlson, Founder Tel: +1.301.279.2005 Fax: +1.240.465.0692 *TonyC@CCT.bz* 

Richard Koulbanis, Editor & Publisher *RichardK*@CCT.bz

Virginia Dematatis, Editorial Consultant VirginiaD@CCT.bz

Loraine Watts, Editorial Assistant

Caryl Cornell, Editorial Assistant

### Sales Office

CONGENITAL CARDIOLOGY TODAY 9008 Copenhaver Drive, Suite M Potomac, MD 20854 USA Tel: +1.301.279.2005 Fax: +1.240.465.0692

### **Editorial Office**

CONGENITAL CARDIOLOGY TODAY 19509 Pine Cone Court, Suite 100 Gaithersburg, MD 20879 USA *Edit*@ *CCT.bz* 

www.CongenitalCardiologyToday.com