



Table of Contents

- 1 **How I Do It: Tips, Tricks and Techniques – Radial Artery Access**
Weiyi Tan, MD, MPH & Jamil Aboulhosn, MD
- 8 **Highlights from PDA Stenting in Cyanotic Newborns: Comprehensive Management Strategies from Fetus to Toddler – First Annual PDA Stenting Symposium**
Howaida El Said, MD, PhD & Brent Gordon, MD
- 11 **2023 Pediatric Cardiology Research Fellowship Award \$35,000**
The Children's Heart Foundation
- 12 **Medical News**
 - PCA 500 – The World's Most Efficient and Versatile 12-lead ECG
 - ACTION and the Parent Project Muscular Dystrophy Develop Patient Registry to Improve Outcomes for Patients with Muscular Dystrophy
- 14 **Meeting Calendar**

How I Do It: Tips, Tricks and Techniques – Radial Artery Access

A PICS Society Education Series

Weiyi Tan, MD, MPH & Jamil Aboulhosn, MD

An Introduction to PICS "Tips and Tricks"

Need a quick and handy review of a novel technique or rare intervention? Want to know how leaders in pediatric and congenital cardiac interventions would approach the case?

The new PICS "Tips & Tricks" section provides congenital interventional cardiologists with nuanced techniques from experienced leaders in the field. These high yield articles highlight their approaches to these challenges, with particular attention paid to practical aspects of procedural techniques, challenges, and potential pitfalls.

Each topic is authored by a key opinion leader with specific procedural expertise and written in an accessible and succinct format with illustrative case examples and suggested references to make this an impactful educational platform.

In this issue of CCT, we present Dr. Tan's and Dr. Aboulhosn's approach to radial access. This duo has significant experience with complex radial access techniques that we are confident readers will appreciate, given the increase in teen and adult congenital case volumes.

We hope you enjoy this issue. We are confident PICS Society members of all experience levels will find this series to be educational and insightful. For the full list of available articles, which will be updated monthly, please visit https://www.picsymposium.com/tips_and_tricks.html.

GURUMURTHY "GURU" HIREMATH, MD, FPICS

Editor, PICS Tips and Tricks

SAROSH "SHAWN" BATLIVALA, MD, MSCI, FPICS

Editor, PICS Tips and Tricks

KAMEL SHIBBANI, MD

Associate Editor, PICS Tips and Tricks

JACQUELINE KREUTZER, MD, FPICS

Consulting Editor, PICS Tips and Tricks & PICS Board Member

Radial Artery Access

Introduction

Radial artery access can be an attractive form of alternative arterial access for cardiac catheterization, as it leads to reduced time for bed rest and a lower incidence of access site complications, such as bleeding, pseudoaneurysm, and arteriovenous fistula. In adults with acute coronary syndrome, radial access has been shown to improve mortality in patients undergoing cardiac catheterization with ST-elevation myocardial infarction, mainly driven by reduced bleeding.¹ In patients with congenital heart disease, radial artery access can also be attractive as many patients have femoral arterial occlusions or stenoses from prior procedures.




TABLE OF CONTENTS

- 1 **How I Do It: Tips, Tricks and Techniques – Radial Artery Access**
Weiyi Tan, MD, MPH & Jamil Aboulhosn, MD
- 8 **Highlights from PDA Stenting in Cyanotic Newborns: Comprehensive Management Strategies from Fetus to Toddler – First Annual PDA Stenting Symposium**
Howaida El Said, MD, PhD & Brent Gordon, MD
- 11 **2023 Pediatric Cardiology Research Fellowship Award \$35,000**
The Children's Heart Foundation
- 12 **Medical News**
 - PCA 500 – The World's Most Efficient and Versatile 12-lead ECG
 - ACTION and the Parent Project Muscular Dystrophy Develop Patient Registry to Improve Outcomes for Patients with Muscular Dystrophy
- 14 **Meeting Calendar**

D'VILL™

LONG-LENGTH HEMOSTASIS INTRODUCER



Optimal flexibility and kink resistance

Trusted by interventionalists worldwide.
Compatible with the CP Stent® family
of products and BIB® Catheters.



FIGURE 1 Radial Artery Cannulation Site and Ultrasound Image

Radial artery cannulation site and image of radial artery on ultrasound at this level. Contributed by Joshua Solano, MD (Wallace MW, Solano JJ. Radial Artery Cannulation. [Updated 2021 Jul 21]. In: StatPearls [Internet])

Anticipated Challenges of the Procedure

Challenges include confirming that the radial artery is an appropriate size, anatomical variation in radial artery anatomy that can preclude radial artery access, and radial artery vasospasm.

Tip 1 – Planning and Preparation

1. Patient Positioning and Setup: Try to use dedicated radial boards to position the hand/arm. Most adult cardiac catheterization labs carry arm boards for this purpose. Adjust the table so you can perform fluoroscopy of the wrist/arm during access.
2. Imaging: Review old imaging and notes. Did the patient have a prior classic BTT shunt that precludes radial artery access on that side? Does the patient have a CT angiogram that includes the subclavian arteries that shows significant tortuosity? Has radial access been attempted in the past with success?
3. Choice of Radial Artery: Traditionally, the right radial artery is chosen as a convenience to the operator, but consider left radial artery access in patients with history of bypass grafts, tortuous vascular anatomy, or right radial artery occlusions. Make sure the radial artery is an adequate size for cannulation (at least 2 mm in diameter).
4. Relative Contraindications: Vaso-occlusive disease (Raynaud disease, scleroderma), upper extremity vascular disease (subclavian artery stenosis, AV fistula for dialysis).



FIGURE 2 Terumo Slender Glidesheath for Radial Artery Access

This is a 6Fr Slender Glidesheath (Terumo Interventional Systems, Somerset, NJ). It has a 2.44 mm outer diameter compared to a regular 6Fr sheath with an outer diameter of 2.63 mm. It has a thin wall that allows for a better sheath to radial artery diameter ratio, which reduces spasm and radial artery occlusion, but it does make the sheath easier to kink.


Tip 2 – Tools Needed

1. Micropuncture Needle and Wire: There are two kinds of access kits, for an anterior wall stick or for the two-wall method.
2. Sheaths: The smaller the better. Terumo Slender Glidesheaths (**Figure 2**) are great for this purpose. A 5Fr Slender has an outer diameter of 2.14 mm, while a 6Fr Slender has an outer diameter of 2.44 mm.
3. Medicines: To prevent vasospasm of the artery. Many different cocktails exist, but I like to give a combination of heparin, nitroglycerin, and verapamil.
4. Catheters: Standard coronary catheters (Judkins left and right) are acceptable, but catheters specifically made for radial access include the Tiger and Jacky catheters. These allow for selective engagement of both coronary arteries with one catheter.
5. Imaging Devices: Ultrasound machine for access. While radial artery access can be successful without ultrasound guidance, studies have demonstrated that ultrasound guidance improves the success and efficiency of radial artery cannulation.²
6. Closure: Radial artery compression device. Use of a compression band, such as the TR Band (Terumo), allows for patent hemostasis and reduces the rate of radial artery occlusion after catheterization.

Tip 3 – How I Do It

1. Palpate the Radial Artery. Use ultrasound to identify the artery and its course if unable to confidently palpate the artery. Although assessment of the RA pulse is important, performing an Allen or Barbeau test to confirm the patency of dual arterial circulation to the hand and intact palmar arch system is only of historical interest. Recent studies of patients with normal and abnormal preprocedural Allen test who subsequently underwent radial access did not demonstrate differences in thumb capillary lactate, grip strength, or incidence of ischemia between the two groups.¹
2. Prep and Drape. Use a dedicated arm board. Sterilize a wide area and drape the area from the styloid process of the radius to about 4-5 cm proximally on the forearm. Adjust the table so the arm can be imaged if necessary.
3. Numb the area of entry with subcutaneous lidocaine. Be careful not to hit the vessel wall as this can cause spasm.
4. Puncture the skin 1-2 cm proximal to the radial styloid.³ Entering the radial artery over the bone can be painful for the patient (**Figure 1**).

RIGHT CHOICE.



Melody™
Transcatheter Pulmonary
Valve (TPV) System



Not intended to constitute medical advice or in any way replace the independent medical judgment of a trained and licensed physician with respect to any patient needs or circumstances. Melody TPV is not suitable for all patients and ease of use, outcomes, and performance may vary. See the Instructions for Use for indications, contraindications, precautions, warnings, and adverse events.

Restoring lives for
13
years and counting.

The only transcatheter pulmonary valve specifically designed for RVOT conduits and bioprosthetic valves. The longest studied transcatheter valve, with the largest body of clinical evidence at over 10 years.* More than 16,000 patients' lives have been changed over 13 years, and counting.

**Melody TPV — The Right Choice
for Your Patients**

*Melody Transcatheter Pulmonary Valve Study:
Post Approval Study of the Original IDE Cohort.
©2020 Medtronic. All rights reserved.
UC201809495b EN 11/2020

Medtronic
Further. Together

Melody™ Transcatheter Pulmonary Valve | Ensemble™ II Transcatheter Valve Delivery System

Important Labeling Information for the United States

Indications: The Melody TPV is indicated for use in the management of pediatric and adult patients who have a clinical indication for intervention on a dysfunctional right ventricular outflow tract (RVOT) conduit or surgical bioprosthetic pulmonary valve that has \geq moderate regurgitation, and/or a mean RVOT gradient \geq 35 mm Hg.

Contraindications: None known.

Warnings/Precautions/Side Effects

- **DO NOT implant in the aortic or mitral position. Pre-clinical bench testing of the Melody valve suggests that valve function and durability will be extremely limited when used in these locations.**
- DO NOT use if patient's anatomy precludes introduction of the valve, if the venous anatomy cannot accommodate a 22 Fr size introducer, or if there is significant obstruction of the central veins.
- DO NOT use if there are clinical or biological signs of infection including active endocarditis. Standard medical and surgical care should be strongly considered in these circumstances.
- Assessment of the coronary artery anatomy for the risk of coronary artery compression should be performed in all patients prior to deployment of the TPV.
- To minimize the risk of conduit rupture, do not use a balloon with a diameter greater than 110% of the nominal diameter (original implant size) of the conduit for pre-dilation of the intended site of deployment, or for deployment of the TPV.
- The potential for stent fracture should be considered in all patients who undergo TPV placement. Radiographic assessment of the stent with chest radiography or fluoroscopy should be included in the routine postoperative evaluation of patients who receive a TPV.
- If a stent fracture is detected, continued monitoring of the stent should be performed in conjunction with clinically appropriate hemodynamic assessment. In patients with stent fracture and significant associated RVOT obstruction or regurgitation, reintervention should be considered in accordance with usual clinical practice.

Potential procedural complications that may result from implantation of the Melody device include the following: rupture of the RVOT conduit, compression of a coronary artery, perforation of a major blood vessel, embolization or migration of the device, perforation of a heart chamber, arrhythmias, allergic reaction to contrast media, cerebrovascular events (TIA, CVA), infection/sepsis, fever, hematoma, radiation-induced erythema, blistering, or peeling of skin, pain, swelling, or bruising at the catheterization site. Potential device-related adverse events that may occur following device implantation include the following: stent fracture; stent fracture resulting in recurrent obstruction, endocarditis, embolization or migration of the device, valvular dysfunction (stenosis or regurgitation), paravalvular leak, valvular thrombosis, pulmonary thromboembolism, hemolysis.

"The term "stent fracture" refers to the fracturing of the Melody TPV. However, in subjects with multiple stents in the RVOT it is difficult to definitively attribute stent fractures to the Melody frame versus another stent.

For additional information, please refer to the Instructions for Use provided with the product or available on <http://manuals.medtronic.com>.

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician.

Important Labeling Information for Geographies Outside of the United States

Indications: The Melody™ TPV is indicated for use in patients with the following clinical conditions:

- Patients with regurgitant prosthetic right ventricular outflow tract (RVOT) conduits or bioprostheses with a clinical indication for invasive or surgical intervention, OR
- Patients with stenotic prosthetic RVOT conduits or bioprostheses where the risk of worsening regurgitation is a relative contraindication to balloon dilatation or stenting

Contraindications

- Venous anatomy unable to accommodate a 22 Fr size introducer sheath
- Implantation of the TPV in the left heart
- RVOT unfavorable for good stent anchorage
- Severe RVOT obstruction, which cannot be dilated by balloon
- Obstruction of the central veins
- Clinical or biological signs of infection
- Active endocarditis
- Known allergy to aspirin or heparin
- Pregnancy

Potential Complications/Adverse Events: Potential procedural complications that may result from implantation of the Melody device include the following: rupture of the RVOT conduit, compression of a coronary artery, perforation of a major blood vessel, embolization or migration of the device, perforation of a heart chamber, arrhythmias, allergic reaction to contrast media, cerebrovascular events (TIA, CVA), infection/sepsis, fever, hematoma, radiation-induced erythema, pain, swelling or bruising at the catheterization site. Potential device-related adverse events that may occur following device implantation include the following: stent fracture; stent fracture resulting in recurrent obstruction, endocarditis, embolization or migration of the device, valvular dysfunction (stenosis or regurgitation), paravalvular leak, valvular thrombosis, pulmonary thromboembolism, hemolysis.

"The term "stent fracture" refers to the fracturing of the Melody TPV. However, in subjects with multiple stents in the RVOT it is difficult to definitively attribute stent fractures to the Melody frame versus another stent.

For additional information, please refer to the Instructions for Use provided with the product or available on <http://manuals.medtronic.com>.

The Melody Transcatheter Pulmonary Valve and Ensemble II Transcatheter Delivery System has received CE Mark approval and is available for distribution in Europe.

medtronic.com

710 Medtronic Parkway
Minneapolis, MN 55432-5604
USA
Tel: (763) 514-4000
Fax: (763) 514-4879
Toll-free: (800) 328-2518

LifeLine
CardioVascular Technical Support
Tel: (877) 526-7890
Fax: (651) 367-0918
rs.structuralheart@medtronic.com




FIGURE 3 Radial Artery Access Kit

Radial artery access kit, with micropuncture needle, wire, and sheath/dilator.

5. If using the anterior-wall stick method, use a 4Fr micropuncture needle that is 2-3 cm in length (usually comes in the kit) at a 45-degree angle to the skin and advance the needle firmly and slowly until the front wall of the artery is punctured. I use ultrasound to guide my access. Tenting of the anterior vessel wall is usually visualized, and then the operator may feel a “pop” and see the anterior wall “bounce” as gentle pressure is applied with the needle to enter the vessel lumen. Confirm that the needle is in the middle of the lumen, visualize pulsatile blood flow from the needle, and then advance the wire. One may have to lower the angle of the needle by a few degrees to facilitate wire entry into the vessel.⁴ If using the two-wall method, use a 20-gauge Angiocath needle system to puncture the anterior wall. Once a flash of arterial blood is seen in the barrel of the Angiocath, then advance the entire system until the back-wall is punctured and the bleed-back stops. Remove the needle and slowly withdraw the plastic catheter until brisk, pulsatile blood flow is obtained, confirming that the catheter is in the arterial lumen. The guidewire can then be advanced through the Angiocath.
6. Use a 0.018-0.021-inch micropuncture wire (Figure 3) with a floppy tip to access the radial artery. Do not advance the wire if there is resistance, as this could signify the wire is in a small branch or is subintimal. Fluoroscopy of the wire tracking up the artery and past the elbow may be necessary in patients with tortuous radial arteries or radial artery loops.
7. Remove the needle and then advance a sheath over the wire. A skin nick should not be necessary as radial artery sheaths are gently tapered and designed to have minimal transition between the sheath and dilator.
8. Remove the dilator, aspirate the sheath, and flush the sheath with an anti-spasmodic “cocktail.”^{1,4} I like to use a combination of heparin (2000-5000 units depending on body weight, 50 units/kg), nitroglycerin (100-200 ug), and verapamil (2.5 mg). Dilute the mixture with 10-20mL of blood, as the mixture is acidic and can cause a burning sensation during infusion if injected too quickly. Flush the sheath with heparinized saline.
9. Secure the sheath with a transparent adhesive dressing.
10. At the end of the case, aspirate and flush the sheath one more time and then remove the sheath after placing a hemostasis device. I use an external compression device to maintain patent hemostasis, such as a TR band (Terumo Interventional Systems, Somerset, NJ), (Figure 4). Confirm patent hemostasis by using a pulse oximetry device placed on the ipsilateral thumb or index finger and document a normal waveform.
11. Excellent video example of radial artery access by Dr. Emmanouil Brilakis.⁵ (<https://www.youtube.com/watch?v=zQCx7wQSe-Q>)

Tip 4 – What Complications to Expect and How to Deal with Them

- Radial artery spasm: Make sure the patient is comfortable. Use topical lidocaine. Use of moderate sedation or general anesthesia leads to a reduction of radial artery spasm as well. Use of spasmolytic agents (nitroglycerin, verapamil) at the beginning of the case and between catheter exchanges is helpful. Anticoagulation with heparin also prevents radial artery occlusion.



FIGURE 4 Radial Artery Compression Device
The TR Band (Terumo Interventional Systems, Somerset, NJ) is a radial artery compression device designed to assist in selective hemostasis of the radial artery after a cardiac catheterization by providing enough pressure to maintain patent hemostasis, which prevents the risk of radial artery occlusion.

- Hypotension: In patients with severe aortic stenosis, left ventricular dysfunction, or cardiogenic shock, be careful in administering spasmolytic agents, as they may reduce preload significantly.
- Radial, Brachial, Aorto-subclavian Artery Tortuosity: There may be significant tortuosity that makes entering the aorta difficult. Use of a 0.035-inch 1.5 mm-radius J-tip wire or 0.035-inch hydrophilic wire may help. In short patients with a short aortic root, having the patient take a deep breath “lengthens” the aorta and may make advancing the wire and catheter into the ascending aorta easier. A stiff exchange-length wire can be used to also straighten tortuous arteries when performing catheter exchanges. Use balloon-assisted tracking (a slightly inflated coronary balloon) or catheter-assisted (smaller French multipurpose or pigtail catheter) tracking if there is resistance with advancing guide catheters through the artery over the wire.¹ Do not be afraid to convert to femoral access. Significant tortuosity may make catheter manipulation difficult, and extended catheter manipulation may lead to arterial vasospasm.
- Radial artery occlusion: It can occur in 3-5% of radial artery catheterizations. Use an external compression device to maintain patent hemostasis. Ipsilateral ulnar artery compression after the procedure can also reduce rates of radial artery occlusion. Fifty percent of radial artery occlusions recanalize in one to three months.¹
- Radial artery avulsion: Do not remove the sheath when there is severe



spasm. Use of more spasmolytics and deepening of anesthesia may be helpful.

- Radial artery hematoma: This is a rare complication, but if it occurs, adjust or reposition the external compression band to a more proximal location. It may be necessary to place a second band more proximal to the first one. If the hematoma is very large, use an inflated blood pressure cuff to provide hemostasis.

3. Wallace MW, Solano JJ. Radial Artery Cannulation [Internet]. StatPearls. Treasure Island, Florida: StatPearls Publishing; 2021. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK539796/>
4. Gupta BS, Nathan S, Perlowski AA. Basics of Radial Artery Access. Card Interv Today [Internet]. 2013;(July/August):25–36. Available from: <https://citoday.com/articles/2013-july-aug/basics-of-radial-artery-access>
5. Brilakis E. Radial access: step-by-step. [Internet]. Manual of PCI. 2020. Available from: <https://www.youtube.com/watch?v=zQCx7wQSe-Q>




Summary

The radial artery is an attractive location for alternative arterial access. When performed in the appropriate patient, radial artery access can lead to improved patient comfort due to reduced bed rest time as well as improved patient safety outcomes due to fewer bleeding complications. In patients with congenital heart disease, radial artery access may also be the only form of arterial access in those with a history of previous cardiac surgery or cardiac catheterizations leading to femoral artery occlusion.

References

1. Mason PJ, Shah B, Tamis-Holland JE, et al. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: A scientific statement from the American Heart Association. Circ Cardiovasc Interv. 2018;11(9):1–21.
2. Seto AH, Roberts JS, Abu-Fadel MS, et al. Real-time ultrasound guidance facilitates transradial access: RAUST (Radial Artery Access with Ultrasound Trial). JACC Cardiovasc Interv. 2015;8(2):283–91.



WEIYI TAN, MD, MPH

*Adult Congenital Cardiology
Assistant Professor, Department of Medicine
University of Texas Southwestern
Chair, PICS Adult CHD Committee
Dallas, TX, USA*



JAMIL ABOULHOSN, MD

*Ahmanson/UCLA Adult Congenital Heart Center
Professor, Departments of Medicine and
Pediatrics
David Geffen School of Medicine at UCLA
Chair, PICS Adult CHD Committee
Member, PICS Adult CHD Committee
Los Angeles, CA, USA*



CONGENITAL CARDIOLOGY TODAY

Timely News & Information for Congenital/Structural Cardiologists & Cardiothoracic Surgeons Worldwide



Recruitment Advertising

- ✓ In print and electronic monthly issue
- ✓ On our website
- ✓ In monthly Email Blast
- ✓ No cost for CCT to create the ad
- ✓ Multiple sizes available



Subscribe Electronically
Free on Home Page

www.CongenitalCardiologyToday.com



Highlights from PDA Stenting in Cyanotic Newborns: Comprehensive Management Strategies from Fetus to Toddler – First Annual PDA Stenting Symposium

Howaida El Said, MD, PhD & Brent Gordon, MD

The inaugural “PDA Stenting in Cyanotic Newborns: Comprehensive Management Strategies from Fetus to Toddler” Conference took place April 20th – 22nd, 2023, at the Loews Coronado Bay Resort in San Diego, California. The meeting was a tremendous success and hosted over 375 registrants from around the world who gathered in person and on zoom to participate in a discussion of this unique and critically important topic. Esteemed faculty from across the country were represented from neonatology, cardiology, intensive care, cardiothoracic surgery, hematology, neurology, bioengineering, and developmental pediatrics. Participants were challenged to discover how to not only enable these critically ill infants to survive, but to help them thrive after PDA stenting, as well.

The meeting kicked off with a welcome from course director Howaida El Said, MD, PhD, before reviewing fetal imaging, prenatal counseling, and how a fetal coordinator functions as the “quarterback” of the cyanotic newborn team. Dr. Amir Ashrafi from Children’s Hospital Orange County discussed optimal pre-procedural feeding strategies and the dangers of mechanical ventilation before Dr. Terri Inder (also from CHOC) gave an outstanding lecture on the complex interplay of prenatal brain development and anesthesia. She encouraged us to talk to our patients every time we examine them and how the human voice and music is critical for the developing brain. Dr. Ganga Krishnamurthy from Columbia New York Presbyterian highlighted the contribution of prematurity and weight to

mortality in congenital heart disease patients. She encouraged an individualized approach for each patient and suggested that prematurity is likely more important than weight. The first evening came to a close as the attendees were treated to an amazing sunset during the welcome reception.

The second day of the conference was jam-packed and focused on the peri-procedural aspects of PDA stenting. The session started with demonstrations of echocardiography and CT angiography and their role in planning PDA stenting. Dr. Timothy Slesnick from Children’s Hospital of Atlanta revealed how all CT scans at CHOA are performed without sedation or heart rate control and how this information helps to predict PDA stent length. Dr. Kanwal Farooqi (also from Columbia) showed a virtual reality PDA stent implant and the session closed with a spirited debate between Justin Ryan PhD and Frank Ing, MD on whether a pre-procedural CT should be required for all patients undergoing PDA stenting. Dr. Ing did a virtual “break-in” to Justin’s house to show 3D images on his walls while Justin demonstrated a live 3D segmentation and PDA stent planning case.

The interventional cardiologists then took to the stage and discussed the ins and outs of the stenting procedure. Dr. John Moore started the session by reviewing the history of PDA stenting and how his early work on lambs was eventually translated to the first attempt at PDA stenting in a patient by Dr.





Gibbs. We were honored by an in person visit from Dr. Alwi Mazeni from Malasia who shared his personal experiences in stenting symptomatic cyanotic newborns and how his pioneering work really opened the eyes of the rest of world to this important intervention. Attendees were then treated to talks on: stenting in dual versus single source pulmonary blood flow, how to manage ductal tortuosity, equipment and team considerations, if jailing pulmonary arteries was a good idea (it turns out it is), and what cardiac anesthesiologists really think of PDA stenting, especially when the baby is flipped! Additional talks included umbilical arterial and transvenous approaches for PDA stenting, and Dr. Dietmar Schranz joined us in person to highlight how PDA stenting in Germany has evolved over the past two decades. The crowd got to enjoy a stunning ocean view as they enjoyed lunch and were entertained by a few taped cases of challenging PDA stents. The afternoon kicked off with a talk on how a bioengineer thinks about a PDA stent and Dr. Lynn Peng from Stanford showed us animal work on the prototype of a custom-made PDA stent from Starlight Cardiovascular. We then learned the top 10 ways to avoid complications from Dr. Ing, and Dr. David Balzer imparted some of his hard-earned wisdom from doing this procedure for many years. Finally, we received an update on the COMPASS trial from one of the trial PIs, Jeffrey Zampi, MD, and how the information gathered from this trial will be so important for our community in many ways. The session ended with a debate on whether or not prostaglandin should be

discontinued before bringing the patient to the catheterization laboratory for a planned PDA stent.

After packing all of that information into a single day we were running behind, but the intensivists got us back on track as they always do! The last session of the day focused on how to take care of these fragile patients after a successful PDA stent. Artificial intelligence is now routinely used to predict trends in patients and alert practitioners to intervene before things get too far off track. Pulse ox and renal NIRS are utilized to predict oxygen delivery to the tissues and used to manipulate systemic and pulmonary resistance in the post-procedural period. Nutrition, specifically the importance of breastmilk, and the role and route of feeding, including timing after the procedure, was highlighted. Dr. John Kim from Colorado Children's Hospital discussed the use of bivalirudin to transition patients to dual anti-coagulation after ductal stenting. Finally, we heard about the PC4 outcomes for PDA stenting versus BTT shunts. After another amazing sunset, the attendees were treated to parent corner, where a family whose child had previously undergone a PDA stent, was interviewed and shared their experiences and provided suggestions for care providers. Their beautiful daughter humbled and reminded everyone why we work so hard to take care of these patients.

The last day kicked off with an in-depth discussion of the home monitoring program, including how to use



MyChart to track patient progress and neurodevelopmental outcomes after PDA stenting. Early intervention remains a key factor and many of the families face a significant amount of stress in the interstage period. While not statistically significant, babies with PDA stents tended to have better gross motor skills and less need for G-tubes as compared to those who received BTT shunts. Stent imaging and reintervention was touched on as well as a review of current literature on outcomes for PDA stenting versus BTT shunts. Dr. Andrew Glatz from St. Louis Children's Hospital and Dr. John Nigro from Rady Children's Hospital debated early primary repair of Tetralogy of Fallot versus PDA stenting, with Dr. Nigro pointing out that many of the patients who travel great distances or face significant challenges in access to care may do better with a primary repair.



FIRST ANNUAL PDA STENTING SYMPOSIUM

The final session of the meeting shifted gears and focused our attention on pulmonary atresia/intact ventricular septum and the unique challenges facing this population. Dr. Tara Karamlou from Cleveland Clinic reviewed how outcomes in these patients often hinges on the size of the tricuspid valve and how performing a right ventricular septal myectomy did not necessarily improve outcomes. The role of coronary artery interrogation and newer methods of non-invasive imaging such as echo VTI, CT angiography, and 4D flow to detect right ventricular dependent coronary circulation were also reviewed. Dr. Henri Justino from Rady Children's challenged the audience to consider transcatheter atrial restriction as an additional strategy to encourage the right ventricle to "grow" after right ventricular decompression. Finally, Dr. Eric Feins from Boston Children's shared their experience with this patient population and showed a recent surgery highlighting takedown of right ventricular muscle bundles, perforated patch closure of the ASD, and creation of a perforated patch in the right pulmonary artery to avoid hypertensive Glenn physiology. The session closed with a review from Laith Alshawabkeh, MD, who discussed what the future holds for these patients when they reach adulthood and how, even if we do get the right ventricle to grow, it may perform abnormally, especially during exercise. After the meeting concluded, interested attendees were treated to a tour of the Dickinson Image-Guided Interventional Center at Rady Children's Hospital, which features two Siemens ARTIS icono biplane catheterization labs with an embedded interventional CMR suite.

All in all, attendees gave the conference high marks and were especially pleased with the ability to focus on one topic in detail. The beautiful weather certainly helped and made us realize how lucky we are to have such wonderful colleagues from across the world. The meeting was recorded and will be made available in the upcoming weeks, so please stay tuned. Thanks again to everyone for their tireless efforts, and we look forward to seeing everyone in San Diego for our next conference!



HOWAIDA EL SAID, MD, PHD

*Professor of Pediatric Cardiology
Course Director
Rady Children's Hospital San Diego
San Diego, CA, USA
helsaid@rchsd.org*



BRENT GORDON, MD

*Associate Professor of Pediatric Cardiology
Rady Children's Hospital San Diego
San Diego, CA, USA*



*8th World Congress of
Pediatric Cardiology
and Cardiac Surgery*

Register Now! *Join the PICS Society at the
World Congress of Pediatric Cardiology and Cardiac Surgery for*

"PICS LIVE"

**at the Marriott Marquis convention center,
Monday, August 28th – Thursday, August 31st, 2023**

PICS LIVE will be concurrent with the **World Congress of Pediatric Cardiology & Cardiac Surgery (WCPCCS)**. Traditionally held every four years, the World Congress is a meeting of unparalleled depth, scope and size in pediatric cardiology/cardiac surgery. This year "PICS LIVE" will focus each morning with the live cases that PICS is renowned for, while enabling PICS LIVE attendees full access to the incredible programming of the entire World Congress.

Registration is Now Open for the World Congress of Pediatric Cardiology and Cardiac Surgery as well as the registration for "PICS LIVE" (you must register for PICS LIVE to be able to attend the live case sessions, in addition to the main World Congress program).

For Registration go to:
<https://www.wcpccs2023.org/Wnm7W7>

For Abstract Submission go to:
<https://www.wcpccs2023.org/bRR78E>

PLEASE VISIT THE WORLD CONGRESS WEBSITE
or the most up to date program information:
www.wcpccs2023.org



2023 Pediatric Cardiology Research Fellowship Award \$35,000

The American Academy of Pediatrics (AAP) and The Children's Heart Foundation (CHF) share a purpose...to promote understanding of the basic mechanism of cardiovascular disease in children. Thirty-five thousand dollars can provide a Training Fellow with invaluable experience in the basic science realm and encouragement to become a physician scientist. The submission site is now open for the 2023 Research Award with a deadline of July 1. You'll find guidelines, application, and timeline posted on the SOCCS Website.

Since 2003, this financial award has sparked the research work of and publications by Dr. Michael Gaies, Dr. Robert Webster, Dr. Thomas Dispenza, Dr. Katja Gist, Dr. Angela Lorts, Dr. Stephanie Nakano, Dr. Cammon Arrington, Dr. Aaron Olson, Dr. Michael-Alice Moga and Dr. Catherine Allan. The complete list of winners and the titles to their work can be found on the Section on Cardiology & Cardiac Surgery website. More recent winners have this to say about the honor and opportunity afforded to them by this award:



JOSHUA KURTZ, MD, FAAP, FACC

Assistant Professor, Director of Pediatric Cardiology Education Division of Pediatric Cardiology, University of Louisville
Congenital Interventional Cardiologist
Norton Children's Hospital



ANDREW L. CHENG, MD, FAAP, FACC

Attending Physician, Division of Pediatric Cardiology, Children's Hospital
Los Angeles
Assistant Professor of Clinical Pediatrics
Keck School of Medicine,
University of Southern California

"The AAP SOCCS Research Fellowship Award was instrumental for launching my career as a physician-scientist. It allowed me to successfully complete the aims of my fellowship research project during my 3rd and 4th years of training. I was then able to use the data from this project as a foundation for grants from the NIH and American Heart Association. Just as important, the award provided me with the unique opportunity to present my research to the entire SOCCS community at the AAP NCE meeting. This was a wonderful formative experience which provided me with sage advice from and connections with prominent pediatric cardiologists and cardiac surgeons from across the country."

"The AAP SOCCS research fellowship award is a great opportunity for pediatric cardiology fellows. As a fellow it allowed me to understand the full extent of what goes into applying for an extramural grant. It helped me solidify my research and career development plans, which are important parts of larger grant awards. It allows for a chance to obtain enough funding to collect data from small projects that can then be used to apply for larger career development awards, such as an NIH K award. This is such a career benefit and comes from an organization dedicated to pediatrics and congenital heart disease. Other society awards are geared towards adult cardiology and give preference to that population. This award is specifically for our population of congenital heart disease and children with heart disease."

If you're interested or know of a training fellow who would benefit from this financial opportunity, please visit the application page, <https://www.aap.org/en/community/aap-sections/cardiology-and-cardiac-surgery/soccs-awards/> for more details. If you have any questions, please contact Vivian Thorne at vthorne@aap.org.



NEONATOLOGY TODAY

Peer Reviewed Research, News and Information in Neonatal and Perinatal Medicine



PCA 500 – The World's Most Efficient and Versatile 12-lead ECG

Los Angeles, California /PRNewswire/ – QT Medical announced FDA clearance of PCA 500, a resting 12-lead electrocardiogram (ECG or EKG), for use in pediatric patients. First cleared in 2018 for professional and personal use by adults 18 years and older, PCA 500's new FDA clearance expands its indication for use to all pediatric populations, including: newborns, infants, children and adolescents.

The development of PCA 500's pre-positioned electrodes and compact recorder was initially funded by NIH for an easy-to-use ECG specifically designed for pediatric use. PCA 500 offers digital, mobile and cloud-based ECG management solution and is widely used by airlines, telehealth practices, clinics, urgent care centers, skilled nursing facilities, hospitals, schools, and in clinical trials, with many CRO partners anticipating the pediatric clearance. QT Medical will introduce PCA 500 to the pediatric market at the 2022 American Academy of Pediatrics National Conference.

"The potential of PCA 500 in improving the heart health of children is extremely exciting. As a pediatric cardiologist, I know how difficult it is to get an ECG on a child. With PCA 500, we can make ECG technology easily accessible to all children. We believe this will make a difference in many lives, which is the exact reason why I founded QT Medical," said Dr. Ruey-Kang Chang, CEO of QT Medical, Inc.

Two new initiatives will be announced at the pediatric market launch of PCA 500. First, Youth Xpress ECG, in partnership with Who We Play For, a non-profit organization dedicated to preventing sudden cardiac arrest of young people, is a mail delivery home 12-lead ECG testing service with interpretation by expert pediatric cardiologists. Second, Baby Xpress ECG, an at-home ECG screening service for newborns and infants with increased risks for Long QT Syndrome (LQTS). With Baby Xpress ECG, when a baby's ECG shows prolonged QT interval, a saliva test will be used to check for genetic mutations known for causing LQTS. Long QT Syndrome, occurring one in every 2000 babies, is a known cause for sudden death (including SIDS).



About QT Medical

QT Medical is a medtech company with a focus on high quality 12-lead diagnostic ECG for use by healthcare professionals and patients. With its simplicity, ease of use, mobile technology and cloud management, PCA 500 brings hospital-grade ECG to millions of patients for better cardiac care at home. More information at: www.qtmedical.com



CHIP NETWORK
CONGENITAL HEART INTERNATIONAL PROFESSIONALS



ACTION and the Parent Project Muscular Dystrophy Develop Patient Registry to Improve Outcomes for Patients with Muscular Dystrophy

ACTION MD Registry is the Largest Real-World Dataset Collecting Comprehensive Cardiac Data

Cincinnati, Ohio – A leading pediatric heart failure organization ACTION (the Advanced Cardiac Therapies Improving Outcomes Network in Cincinnati, OH), in conjunction with support of Parent Project Muscular Dystrophy (PPMD), has developed a patient registry to improve outcomes for the Muscular Dystrophy patient population. To date, this registry is the largest real-world dataset collecting comprehensive cardiac data.

"The care practices are rapidly evolving and there's an urgent need to understand how best to treat cardiac disease to improve the outcomes for Muscular Dystrophy patients," said Chet Villa, MD, ACTION, Cincinnati Children's. ACTION's initial MD study focused on Duchenne Muscular Dystrophy (DMD), the most common of muscular dystrophies. DMD is caused by the lack of a functional dystrophin protein and leads to progressive skeletal, lung and cardiac muscle weakness. The long-term survival of those living with DMD is increasing, leading to a growing number of people living with heart dysfunction.

As a result, clinicians and families are having to make significant decisions regarding cardiac care. This includes when to start cardiac medications and when to consider advanced therapies such as cardiac devices or transplantation. The burgeoning field of gene therapy and implications for the heart remains an important question for ACTION teams to answer. Additionally, ACTION's current DMD studies have highlighted the need to broaden its understanding to include individuals with Becker Muscular Dystrophy (BMD), as well as female carriers of the dystrophin mutation.

"Not only will ACTION improve the care and treatment of patients with dystrophinopathies, but this information

has the potential to impact all individuals with cardiomyopathy," said Linda Cripe, MD, ACTION, Nationwide Children's.

Until ACTION's MD initiative, patients, caregivers and providers have not had access to research and opportunities for evidence-based care. Research in a rare disease population poses unique challenges and requires multi-institutional data sharing. The enhanced ACTION MD Registry is the largest real-world dataset collecting comprehensive cardiac data.



Ricky, a Duchenne Muscular Dystrophy Patient, Orlando, Florida

As of April 10th, 2023, ACTION's MD registry includes:

- Institutions - 22 pediatric institutions have entered patient data.
- Patients - 275 patients have been enrolled in the MD Registry.

ACTION's infrastructure allows the database to facilitate a knowledge sharing, data driven and quality improvement approach to formalizing MD education, best practices and future clinical trial design to move the field forward and ultimately improve MD patient outcomes.

"We really stand at the precipice of a new era of cardiac care for patients

with muscular dystrophies and heart failure," said Deip Nandi, MD, ACTION, Nationwide Children's. "By organizing cardiologists across the country and world within the framework of ACTION, we're confident that we will be able to capture and harness the data we need to make real improvements in quantity and quality of life for our patients."

To learn more about ACTION and ACTION's Muscular Dystrophy registry and initiatives, please visit

www.actionlearningnetwork.org; or contact ACTION directly at info@actionlearningnetwork.org.

About ACTION

ACTION (the Advanced Cardiac Therapies Improving Outcomes Network) is an organization that unites and connects a global healthcare community to improve outcomes for patients with heart failure, especially children. Since 2017, ACTION has continued to grow and engage an international community by uniting key stakeholders: patients, families, clinicians, researchers, payors and industry.

Based in Cincinnati, OH, ACTION uses a quality improvement and research-based approach. ACTION works in collaboration with network sites and network members, sharing data, finding better solutions and discovering new innovations. Currently, there are over 60 network sites and 1,185 network members. Learn more at www.actionlearningnetwork.org.





JULY

28-29

CICT 2023 – CICT Controversies in Interventional Cardiovascular Therapies

Pasadena, CA, USA

<https://cictsymposium.com/>

AUGUST

25-27

3rd Annual PICS Fellows & Early Career Course

Washington, DC, USA

kimberly_ray@chdinterventions.org

27-09/01

8th World Congress of Pediatric Cardiology and Cardiac Surgery

Washington, DC, USA

<http://wcpccs2023.org/>

SEPTEMBER

08-09

2023WPC – 2023 World Pediatric Conference

Singapore

<https://pediatrics.episirus.org/>

25-26

CME HeartCare and Cardiovascular Medicine Cardiac Surgery

Paris, France

<https://heart.plenareno.com/>

OCTOBER

06-08

CSI Asia-Pacific 2023

Bangkok, Thailand

<https://www.csi-congress.org/asia-pacific>



*8th World Congress of
Pediatric Cardiology
and Cardiac Surgery*

AUGUST 27TH - SEPTEMBER 1ST, 2023
WASHINGTON D.C

www.WCPCCS2023.org

Congenital Cardiac Surgery • Ambulatory Cardiology • Cardiac Catheterization • Multimodal Imaging • Adult Congenital Heart Disease
Echocardiography • Cardiac Intensive Care • Cardiovascular Nursing • Cardiovascular Disease in the Neonate • Cardiac Anesthesia
Heart Failure and Transplantation • Administration & Leadership • Trainee Track • Neurodevelopment • Pulmonary Hypertension
Rheumatic Heart Disease • Global Cardiac Health • Electrophysiology • Advocacy • Sudden Cardiac Death in the Young
Basic Science • Fetal Cardiology • Special Sessions for Perfusionists • Big Data, New Technology & Artificial Intelligence
Quality Improvement • Cardiopulmonary Bypass, ECMO & Mechanical Circulatory Support



CORPORATE TEAM

**PUBLISHER &
EDITOR-IN-CHIEF**

Kate Baldwin
kate.f.baldwin@gmail.com

**CO-FOUNDER &
MEDICAL EDITOR**

John W. Moore, MD, MPH
jwmmoore1950@gmail.com

**FOUNDER &
SENIOR EDITOR**

Tony Carlson
tcarlsonmd@gmail.com

**STAFF EDITOR &
WRITER**

Virginia Dematatis

**EDITOR-IN-CHIEF
EMERITUS**

Richard Koulbanis

STAFF EDITOR

Loraine Watts

EDITORIAL BOARD

Aimee K. Armstrong, MD
Jacek Bialkowski, MD
Anthony C. Chang, MD, MBA
Howaida El-Said, MD, PhD
Ziyad M. Hijazi, MD, MPH
John Lamberti, MD
Tarek S. Momenah, MBBS, DCH

John W. Moore, MD, MPH
Shakeel A. Qureshi, MD
P. Syamasundar Rao, MD
Carlos E. Ruiz, MD, PhD
Hideshi Tomita, MD
Sara M. Trucco, MD
Gil Wernovsky, MD

OFFICIAL NEWS & INFORMATION PARTNER OF



Pediatric and Congenital Interventional Cardiovascular Society

Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors and are not necessarily the views of Congenital Cardiology Today.

© 2023 by Congenital Cardiology Today LLC
ISSN 1554-7787 print. ISSN 1554-0499 electronic.
Published monthly. All rights reserved.