



## Table of Contents

### 1 Transcatheter Patent Ductus Arteriosus Closure in Extremely Low Birth Weight Infants: From Catheterization Lab to Bedside

Jason N. Johnson, MD, MHS;  
Ranjit Philip, MD;  
Ronak Naik, MD, DNB, FACC;  
Vijaya Joshi, MD;  
Ashley Giannini, RCIS;  
Kaitlyn Greer, RT(R);  
Christina Tunney, RN, BSN;  
Matthew Balmut, BS, BSN;  
Lindsey L. Pierotti, RDCS;  
Nathan Stecchi, RDCS;  
Huma Abdulmajeed, RDCS;  
Mark F. Weems, MD;  
Ajay Talati, MD;  
Tim Head, DO;  
Thomas Hamilton, MD;  
Genevieve DuPuis, CRNA;  
Janet Sessums, CRNA;  
Leah Apalodimas, MSN, CPNP-AC/PC;  
Ashley Kiene, MD;  
Neil Taylor, MD;  
B. Rush Waller III, MD;  
Shyam Sathanandam, MD FSCAI

### 14 Report From the Field - Early Career Colleagues

Kamel Shibbani, MD

### 18 Medical News

- CTA and ACC Create First-Ever Industry Framework for Cardiovascular Technology
- Structured Reporting Systems are Expected to Dramatically Improve Patient Care and Data Management

### 19 Meeting Calendar

Career Opportunities Throughout

# Transcatheter Patent Ductus Arteriosus Closure in Extremely Low Birth Weight Infants: From Catheterization Lab to Bedside

Jason N. Johnson, MD, MHS; Ranjit Philip, MD; Ronak Naik, MD, DNB, FACC; Vijaya Joshi, MD; Ashley Giannini, RCIS; Kaitlyn Greer, RT(R); Christina Tunney, RN, BSN; Matthew Balmut, BS, BSN; Lindsey L. Pierotti, RDCS; Nathan Stecchi, RDCS; Huma Abdulmajeed, RDCS; Mark F. Weems, MD; Ajay Talati, MD; Tim Head, DO; Thomas Hamilton, MD; Genevieve DuPuis, CRNA; Janet Sessums, CRNA; Leah Apalodimas, MSN, CPNP-AC/PC; Ashley Kiene, MD; Neil Taylor, MD; B. Rush Waller III, MD; Shyam Sathanandam, MD FSCAI

## Introduction

Many different clinical approaches exist to the management of a Patent Ductus Arteriosus (PDA) in a premature neonate: observation, medical therapy, surgical ligation, and transcatheter PDA closure (TCPC).<sup>1</sup> TCPC in Extremely Low Birth Weight (ELBW) neonates has quickly become a common practice in many centers with expertise with several studies showing safe and successful PDA closure with this technique.<sup>2,3</sup> TCPC in ELBW infants commonly occurs in the catheterization lab requiring the patient to transfer from the neonatal intensive care unit (NICU), and likely adds to the hesitancy of neonatologists to fully support TCPC.<sup>4-6</sup> Therefore, it is imperative to bring TCPC safely and successfully “to the bedside” to provide the best possible outcomes for patients. This article will highlight our programmatic practice regarding PDA evaluation, and the evolution of the TCPC program in ELBW infants from the catheterization lab to the NICU bedside (Figure 1).



**FIGURE 1** Catheterization lab and neonatal bedside transcatheter patent ductus arteriosus closure set up. **A)** Catheterization lab. The implanter and assistant stand on the right side of the patient and the sonographer stands on the left side with left-handed scanning. **B)** Neonatal bedside. The implanter and assistant stand on the right side of the patient while the mobile digital Mini C-Arm and the echocardiographer are on the left side of the patient with right-handed scanning.

## Hemodynamically Significant PDA

A consensus definition of a hemodynamically significant PDA (hsPDA) does not exist. A combination of clinical history and echocardiographic characteristics is the best way to define PDA significance.<sup>7-10</sup> Clinical factors suggesting a hsPDA are: ventilator support due to pulmonary edema, requirement

## TABLE OF CONTENTS

### 1 **Transcatheter Patent Ductus Arteriosus Closure in Extremely Low Birth Weight Infants: From Catheterization Lab to Bedside**

Jason N. Johnson, MD, MHS; Ranjit Philip, MD; Ronak Naik, MD, DNB, FACC; Vijaya Joshi, MD; Ashley Giannini, RCIS; Kaitlyn Greer, RT(R); Christina Tunney, RN, BSN; Matthew Balmut, BS, BSN; Lindsey L. Pierotti, RDCS; Nathan Stecchi, RDCS; Huma Abdulmajeed, RDCS; Mark F. Weems, MD; Ajay Talati, MD; Tim Head, DO; Thomas Hamilton, MD; Genevieve DuPuis, CRNA; Janet Sessums CRNA; Leah Apalodimas, MSN, CPNP-AC/PC; Ashley Kiene, MD; Neil Taylor, MD; B. Rush Waller III, MD; Shyam Sathanandam, MD FSCAI

### 14 **Report From the Field - Early Career Colleagues**

Kamel Shibbani, MD

### 18 **Medical News**

- CTA and ACC Create First-Ever Industry Framework for Cardiovascular Technology
- Structured Reporting Systems are Expected to Dramatically Improve Patient Care and Data Management

### 19 **Meeting Calendar**

Career Opportunities Throughout




Interventional Systems  
**B | BRAUN**  
SHARING EXPERTISE


# COMPASS™

## Guiding Introducer Sheath

Designed for exceptional kink-resistant performance to support complex vascular access.



Distributed by:  
B. Braun Interventional Systems Inc.  
Part of the B. Braun Group of Companies  
Bethlehem, PA | USA | 877-836-2228  
www.bisusa.com




Available in  
4-8F sheath sizes  
and 15cm, 45cm,  
60cm, and  
90cm lengths

Rx only  
Refer to the Instructions for Use for complete indications, relevant warnings, precautions, complications, and contraindications.  
Compass is a trademark of B. Braun Interventional Systems Inc. ©2022 B. Braun Interventional Systems Inc. CV-9152 2/22



of inotropes, feeding intolerance, and renal insufficiency in the context of a vulnerable patient.<sup>11</sup> Neonates born <26 weeks' gestation are a vulnerable population because they do not have spontaneous PDA closure until a median of 71 days, sustaining significant complications of a hsPDA.<sup>12</sup>




**FIGURE 2** Echocardiographic assessment of a hemodynamically significant PDA. **A).** Two dimensional measurements of a large PDA. **B).** Color Doppler of a large PDA with low velocity left to right shunting. **C).** Four chamber with moderate left atrial and ventricular dilation. **D).** Left pulmonary artery spectral Doppler with diastolic continuation of flow (arrows). **E).** Abdominal aorta spectral Doppler with diastolic reversal of flow (arrows). LA=left atrium. LPA=left pulmonary artery. LV=left ventricle. PDA=patent ductus arteriosus.

There are numerous protocols to report echocardiographic parameters of a hsPDA, but none have been concurrently confirmed with cardiac catheterization hemodynamics.<sup>1,13,14</sup> Echocardiographic parameters focus on PDA shunt size, extent of volume overload, and evidence of systemic hypoperfusion.<sup>11</sup> A minimal PDA diameter >1.5 mm and low velocity left to right flow pattern suggest a large PDA shunt volume (**Figure 2**). However, PDA diameter alone may not correlate with a large shunt, and there can be interobserver variability in the measurement.<sup>15</sup> Left atrial and ventricular chamber dilation and diastolic continuation of flow in the branch pulmonary arteries suggest a large volume overload from a hsPDA (**Figure 2**). Diastolic flow reversal in the abdominal aorta suggests systemic hypoperfusion (**Figure 2**).

Accurate communication of echocardiographic findings of a hsPDA to the neonatology team is essential. Simply stating the PDA is small, moderate, or large is not sufficient information for the neonatologist to determine hemodynamic significance. At our institution a quality improvement initiative improved the echocardiographic communication of the different factors potentially defining a hsPDA. Our echocardiographic summary reports include minimal PDA diameter, PDA shunt direction with peak velocity, left atrial (LA) and ventricular chamber size, and abdominal aorta flow pattern at the minimum. Also, there is no consensus definition of a hsPDA at our institution agreed upon by all cardiologists and neonatologists. The goals of the echocardiogram report are to be consistent and provide easily understandable and reproducible data in a timely manner for the neonatologist. At some institutions, the neonatology team performs and reports the PDA echocardiograms and are experts in echocardiographic evaluation of hsPDA.<sup>16</sup>

## Timing of PDA Closure

An extensive discussion on prophylactic PDA closure, medication attempts at PDA closure, and surgical ligation are outside the scope of this review.<sup>1</sup> However, a brief review of the clinical conundrum the presence and treatment of a hsPDA poses to clinicians caring for ELBW neonates follows. Despite multiple studies outlining the morbidity caused by a hsPDA, there is a lack of decrease in morbidity and mortality in prophylactic medical or surgical PDA closure.<sup>17-19</sup> The PDA-TOLERATE trial found medical treatment after one week of life was not associated with reduced PDA ligations or PDAs present at discharge in infants born before 28 weeks gestation, and infants born between 26 and 28 weeks gestation had increased rates of late-onset sepsis and death when exposed to early medical PDA treatment.<sup>20</sup> Surgical ligation has frequently been linked to increased morbidities such as Bronchopulmonary Dysplasia (BPD), Retinopathy of Prematurity (ROP), and poor neurodevelopmental outcomes.<sup>21-23</sup> Due to no convincing studies showing clear benefit of medical and surgical PDA ligation, many neonatologists have adopted a conservative approach to PDA management.<sup>24</sup> However, spontaneous closure of PDAs and the need for rescue treatment for those in the conservative treatment group make it difficult to accurately evaluate the morbidities associated with conservative treatment alone.<sup>25</sup>



**FIGURE 3** Sequence of 1-301 patients with the procedure weight at the time of transcatheter patent ductus arteriosus closure at Le Bonheur Children's Hospital.

Initial results of TCPC in infancy showed mixed results and added to the clinical conundrum of PDA management especially in ELBW neonates.<sup>3</sup> However, a recent meta-analysis of TCPC in infants ≤1.5 kg showed a technical success of 96% with a major adverse event rate of 8%.<sup>2</sup> This increased procedural success of TCPC occurred in an era of decreasing patient weight at time of procedure.<sup>2</sup> Our institutional TCPC experience shows excellent success rate with low adverse events (AE) and suggests that earlier closure (<4 weeks) is associated with improved outcomes including quicker weaning of ventilatory support and improved growth.<sup>26,27</sup> Also, ELBW infants referred for TCPC >8 weeks may have elevated pulmonary vascular resistance suggesting a likely loss of benefit of PDA closure in patients referred late.<sup>27</sup> Over time this has led to our patient referral for TCPC to be younger and weigh less at time of procedure (**Figure 3**). There is no minimum patient weight threshold yet identified, with reports including patients as small as 490 grams, and the Amplatzer Piccolo Occluder (Abbott Medical, New Plymouth, MN, USA, APO) device is approved in infants weighing >700 grams.<sup>28-30</sup> While TCPC is technically feasible at very early ages and small weights, timing is often dictated by medical management in the absence of clear PDA treatment guidelines.<sup>31</sup> Earlier in our experience with TCPC, a core group of neonatologists and cardiologists would discuss the indications and eligibility for TCPC for every neonate <1.5 kg referred for the procedure.



# Pediatric Cardiologist

Orlando, FL, United States



## JOB DESCRIPTION

As one of the nation's leading pediatric health care systems, Nemours Health System is a not-for-profit pediatric health system dedicated to life-changing medical care and research, helping kids grow up healthy, advocating for kids nationally, and training tomorrow's pediatric experts. We offer integrated, family-centered care to more than 300,000 children each year in our pediatric hospitals, specialty clinics, and primary care practices in Florida, Georgia, Maryland, New Jersey, and Pennsylvania. Nemours strives to ensure a healthier tomorrow for all children – even those who may never enter our doors – through our world-changing research, education, and advocacy efforts. At Nemours, our Associates help us deliver on the promise we make to every family we have the privilege of serving: to treat their children as if they were our own.

Nemours Children's Hospital in Orlando, Florida is seeking a board-certified/board-eligible, full-time non-invasive pediatric cardiologist to serve as the director of pediatric cardiology for the winter garden office.

This is a full-time employed opportunity to join an expanding program dedicated to meeting the growing needs of Central Florida.

## REQUIRED QUALIFICATIONS

- Be proficient in general cardiology and noninvasive testing (EKG, Holter/ event monitoring, and echocardiography)
- Additional training in echocardiography would be welcomed and able to participate in reading in the echocardiography laboratory or perinatal cardiology clinics at Nemours
- Be eligible for an unrestricted FL license
- Be board certified or board eligible in pediatric cardiology
- Able to take calls which includes a 1:9 rotation at the hospital

Nemours cardiac center in Florida is a comprehensive program with cardiac surgery, interventional cardiology, advanced imaging program (MRI/CT), electrophysiology, heart failure/ pulmonary hypertension, adult congenital and preventative cardiology programs. The candidate would be joining a seventeen-person group that is expanding and dedicated to meeting the growing needs of children in the Central Florida region.

**For confidential consideration, please forward your formal CV to:**

Fernando Moncada, CPRP, Physician Recruiter

[Fernando.moncada@nemours.org](mailto:Fernando.moncada@nemours.org)

Our Orlando hospital is located in beautiful Lake Nona, a nationally recognized healthy community: [Lake Nona](#)

A hospital designed by families for families, Nemours Children's Hospital blends the healing power of nature with the latest in healthcare innovation to deliver world-class care to the children of Central Florida and beyond. In keeping with our goal of bringing Nemours care into the communities we serve, we also provide specialty outpatient care in several clinics located throughout the region.

Nemours offers a competitive salary and a robust benefits package that includes productivity and quality-based financial incentives, relocation, health, life, dental, CME, 403B Retirement Plan with matching, 457-retirement savings plan, licensure, and dues allowance. Florida has no state income tax, and Nemours is a not-for-profit; full-time employed physicians qualify for Public Service Loan Forgiveness. Academic appointment(s) are also available with the University of Central Florida Medical School and the Florida State University School of Medicine for qualified applicants.

## ABOUT US

As one of the nation's premier pediatric health care systems, we've made a promise to do whatever it takes to prevent and treat even the most disabling childhood conditions. It's a promise that extends beyond our nationally recognized clinical treatment to an entire integrated spectrum of research, advocacy, education, and prevention.

Equity, diversity, and inclusion guide our growth and strategy. We are looking for individuals who are passionate about, and committed to, leading efforts to provide culturally relevant care, reducing health disparities, and helping build a diverse and inclusive environment. All Nemours Associates are expected to ensure that these philosophies are embedded in their day to day work with colleagues, patients and families.

To learn more about Nemours and our commitment to treat every child as if they were our own, visit us at [www.nemours.org](http://www.nemours.org), take a [virtual tour](#) of Nemours.

[Apply Now](#)

Job Identification: 3290 -- Job Category: Physician -- Posting Date: 12/17/2021, 11:10 AM  
Degree Level: Post-Doctorate -- Job Schedule: Full time -- Location: 1717 S Orange Ave, Orlando, FL, 32806, US



## Transcatheter PDA Closure Expertise

We have created a comprehensive program in the care of preterm neonates and infants with PDA before and after TCPC.<sup>32</sup> This involves developing expertise with PDA management amongst neonatology, cardiology, anesthesiology, pulmonology, neurodevelopmental specialists, nutrition, speech therapy, social work, medical transport, and research collaborators. Our comprehensive neonatal intensive care unit provides multi-disciplinary care to all ELBW neonates and infants regardless of the presence of a PDA. Our program is unique in creating specific outpatient follow-up after patients are discharged following TCPC. This provides an opportunity for long-term follow-up of specific cardiovascular and neurodevelopmental outcomes that are currently ongoing.

TCPC is standard of care for PDA management in children and adults with PDA.<sup>33</sup> However, the technique of TCPC in ELBW neonates is vastly different than TCPC in children and adults. Therefore, creating procedural expertise amongst all individuals assisting in TCPC is essential to programmatic success.<sup>34</sup> This translates to one catheterization implanter, a core group of cardiology imagers and sonographers, a core group of pediatric anesthesiologists and certified registered nurse anesthetists (CRNA) perform and assist in a vast majority of the TCPC procedures in ELBW infants. Repeated TCPC procedural techniques, echocardiographic evaluation, and anesthetic support creates expertise among these individuals when performing TCPC in ELBW infants.<sup>4,30,35</sup>


## TCPC Program Evolution

The TCPC program at our institution has evolved as newer devices, technology, and expertise progressed over time. Initially TCPC was performed with devices, Amplatzer Vascular Plug II (Abbott, Chicago, AVPII) and Microvascular Plug (Medtronic, MVP), that were not necessarily designed for PDA closure in ELBW neonates.<sup>35-38</sup> Our experience with the AVPII led to AEs before we began using the MVP with mostly successful results and fewer AEs.<sup>26</sup> Once the APO received FDA approval for TCPC in neonates >700 grams, we utilized the APO on a vast majority of TCPC cases with excellent results.<sup>26,28</sup> As the devices available improved, the TCPC procedure in the catheterization lab became refined. Performing TCPC in the cardiac catheterization lab allows the device implanter to become comfortable with the subtle aspects of the procedure with biplane fluoroscopy and echocardiographic imaging available. Also, the device implanter has a full assortment of inventory and devices to deal with potential complications that may arise. Developing procedural expertise with multiple TCPC cases in ELBW neonates is recommended before consideration to performing the procedure outside the catheterization lab.

Despite the many benefits of performing TCPC in the catheterization lab, the best place for the procedure for the patient may be the NICU. Transfer of ELBW neonates outside the NICU is associated with increased morbidity and mortality.<sup>39,40</sup> NICU bedside TCPC (bedside-TCPC) has been previously described at other institutions but has not been performed routinely at centers performing TCPC in neonatal patients.<sup>41-44</sup> There are differences between the techniques of TCPC depending upon the location of the procedure.<sup>31</sup> Bedside-TCPC requires greater utilization of echocardiographic evaluation as there is limited fluoroscopy. With anticipation of this difference, we performed several TCPC in neonates with an attempted echocardiographic only approach. This approach was mostly successful, but there were some cases where fluoroscopy was utilized to complete the procedure. Therefore, a hybrid fluoroscopy and echocardiographic approach was adopted at the bedside. Below are the key procedural approaches to both techniques depending upon location.


## Catheterization Lab

When transporting an ELBW to the catheterization laboratory, several important modifications need to be considered and have been well described by our group.<sup>5</sup> A team including the anesthesiologist, CRNA, and two to three Cath lab staff are involved in the transport process. The goal of the transport is to minimize the risk of unplanned extubation during transport, maintain adequate temperature, and reduce the time out of the NICU. Use of a transport ventilator such as a Neopuff (NeoPuff infant resuscitator, Fisher and Paykel, Auckland, New Zealand) or a PneuPAC® BabyPAC (Smiths Medical, Minneapolis, MN) is common in patients on conventional ventilation. For patients on high frequency ventilators (high frequency jet ventilators or high frequency oscillatory ventilators), we use a Bronchotron Transport (TXP®-2D, Percussionaire Corporation, Sandpoint, ID, USA) for transport. Early in our experience, we would give a trial of conventional ventilator for two hours in patients on high frequency ventilators, and if the PCO<sub>2</sub> did not increase by more than 10mmHg from baseline, the procedure was performed on a regular ventilator. However, more recently we have found the patients are more stable when we perform the procedure on the high frequency ventilator in the catheterization lab with the assistance of the NICU respiratory therapist.



**FIGURE 4** Pre-procedure transthoracic echocardiogram and lateral angiogram of the patent ductus arteriosus. **A).** Pre-procedure transthoracic echocardiogram. **B).** Lateral angiogram. A= PDA diameter at the aortic end; B = PDA diameter at the pulmonary end; C = PDA length.

Once in the catheterization lab, the infant is positioned on the Bair Hugger (3M Maplewood, MN, USA) by the physician with a folded towel beneath the sacrum to straighten the groins and allow for easier vascular access. A temperature probe (SourceMark, Brentwood, TN, USA) is placed in the esophagus and will later serve as a landmark of the descending aorta for device deployment. Any wires or monitors that are radiopaque are moved out of the view of the frontal and lateral cameras and taped down so as not to obscure the field. During table set-up, a transthoracic echocardiogram (TTE) is performed with a sonographer at the head of the table between the frontal and lateral cameras. It is encouraged to perform the pre-procedure TTE with the lateral camera in place so the sonographer can become accustomed to the arm angle and patient windows during the procedure.<sup>45</sup> The pre-procedure TTE focuses on the details of the PDA as well as the potential structures (aortic arch, left pulmonary artery (LPA), and tricuspid valve) affected by device placement and prograde approach.<sup>31,45</sup> A TTE pre-procedure check list (**Table 1**) and is performed on every patient prior to the procedure.



**FIGURE 5** Lateral fluoroscopy angiogram after Amplatzer Piccolo Occluder (Piccolo) deployment and release. **A)** The device is between the esophageal temperature probe and upper extremity PICC line in a 10 o'clock to 4 o'clock position. **B)** The "PAC-MAN" is created by the touching of the proximal and distal discs along the lesser (inferior) curvature of the PDA and the splaying of the discs along the greater (superior) curvature of the PDA. An upper extremity PICC line can highlight the pulmonary end of the ductus. PICC = peripherally inserted central catheter; Temp = temperature.

A purely transvenous prograde approach was adopted for TCPC to avoid accessing the femoral arteries in neonates.<sup>41,43</sup> A femoral artery diameter <3 mm has been associated with loss of arterial pulse, and ELBW neonates are at high risk of arterial damage.<sup>46,47</sup> Ultrasound guided vascular access of the femoral vein is obtained, and heparinized saline flushes are utilized throughout the case instead of heparin bolus. All fluoroscopy is performed at three frames per second to limit radiation exposure to the patient. There are two methods for catheter access to the PDA from a prograde approach.<sup>4,35,36,48</sup> A balloon tip catheter can be used to access the PDA, but the preferred method at our institution is the method using a 4 Fr/65cm Terumo Glide Catheter and Wholey wire.<sup>4,31,35,43</sup> Once the catheter is in place, a biplane angiogram with 1 ml of contrast is hand-injected with the frontal plane 15° LAO and 15° Caudal and the lateral plane kept at 90° LAO. Measurements are made from the lateral angiogram and compared to the TTE measurements to choose the appropriate APO device size (**Figure 4**).<sup>49</sup> Most PDA in patients <1000 grams are the F-type described as a long tubular PDA with a bend at the pulmonary artery end much like a hockey stick.<sup>50</sup> The location of the descending aorta in relationship to the esophageal temperature probe should be clearly evaluated on the lateral angiogram as the exact location of the probe in relationship to the descending aorta is slightly different in each patient.<sup>31</sup>

Once the appropriate size device is chosen, the patient should be placed briefly on higher FiO<sub>2</sub> until the device is implanted.<sup>51</sup> Through the side-arm of the sheath, Calcium Chloride (CaCl<sub>2</sub>) 10mg/kg is administered.<sup>51</sup> These maneuvers: brief period of hyperoxygenation and CaCl<sub>2</sub> administration at our institution, have been shown to not only keep the infant stable during the device implantation process, but also to decrease the already low incidence of post-ligation syndrome after TCPC.<sup>51,52</sup> The TorqVue LP catheter (Abbott, New Plymouth, MN, USA) is then advanced over the wire to the descending aorta and a Tuohy-Borst Side arm adapter (Nordson Medical, Westlake, OH, USA) is attached. The device is then advanced through the catheter to the tip, which is 3mm beyond the radiopaque marker. For proper implantation, a single operator, two-hand technique, with the left hand on the delivery catheter and the introducer sheath, facing the ground, and the right hand on the delivery catheter and the delivery cable, but facing the ceiling, working in unison with a gentle "push-pull" method to deploy the device intra-ductal. Keeping the tip of the catheter steady with the left hand, the right hand pushes the cable to expose the distal disc of the device within the aortic end of the PDA. After this maneuver, the device can be pushed into place with the right-hand always maintaining catheter position within the PDA.

Once the device is deployed, a lateral angiogram is performed with 1 ml of contrast for verification of device position on the PA end. As stated previously, the esophageal temperature probe acts as a landmark of the descending aorta in this case. If the patient has an upper extremity PICC, this well defines the pulmonary end of the PDA. The device should sit between these two landmarks on the straight lateral position oriented in a 10-4 position on a clock face (**Figure 5**). Also, the device should be packed within the PDA to create a "PAC-MAN" sign on lateral fluoroscopy (**Figure 5**). The PAC-MAN is created by the touching of the proximal and distal discs along the lesser (inferior) curvature of the PDA and the splaying of the discs along the greater (superior) curvature of the PDA. This position is highly stable and the device is less likely to migrate or embolize.<sup>53</sup> After the device is positioned, lower extremity pulses are checked and blood pressure is obtained to rule out aortic obstruction.<sup>53</sup>


Although a levophase lateral angiogram could evaluate the aortic arch, a TTE is the preferred method to check for aortic obstruction. The first images obtained by TTE should be of the aortic arch as it is often not evaluated by post-deployment angiography. **Table 1** outlines the post-procedure echocardiogram views and techniques to ensure the device is intraductal with no residual shunting and no aortic arch or LPA stenosis. Utilization of post-procedure spectral Doppler patterns of the aortic arch and LPA compared to the pre-procedure Doppler patterns can aid in determining if device obstruction is present.<sup>31,53</sup> Trivial residual shunting through the center of the device is not uncommon immediately after device deployment, and we call this the "flame sign" due to the small width of the color jet by Doppler (**Figure 6**). Trivial residual shunting almost always resolves within 5-10 minutes and should not be used as an indication to remove the device.<sup>31,45</sup> Instruct the sonographer to obtain images continuously and be as quick as possible. Some patients will become unstable with the delivery sheath across the tricuspid valve, so timely TTE evaluation is important.<sup>53</sup> Accurate, decisive, and quick communication of the post-deployment TTE to the catheterization team is imperative.<sup>45</sup>

If device position is adequate, the device can be released, and the TorqVue catheter is removed immediately. There is no reason to perform another angiogram as a TTE is sufficient to check for device position, aortic or LPA obstruction, residual shunt, tricuspid regurgitation, left ventricle (LV) function and effusion (**Table 1**). At this point, the FiO<sub>2</sub> is weaned back down, and fluoroscopy is done to evaluate for device movement. If device is still in place, the sheath can be removed, and manual pressure is held to obtain hemostasis.

**Table 1** Transthoracic echocardiogram checklist

Structure	Techniques	PRE - PROCEDURE	POST - PROCEDURE
PDA	2D	PDA Measurements	Device Orientation
	Color Doppler	Shunt Direction	Residual Shunting
	Spectral Doppler		
Aortic Arch	2D	Rule Out Coarctation	Device Causing Obstruction
	Color Doppler		
	Spectral Doppler	Compare to Post-Procedure	Compare to Pre-Procedure
LPA	2D	Rule Out Stenosis	Device Causing Obstruction
	Color Doppler		
	Spectral Doppler	Compare to Post-Procedure	Compare to Pre-Procedure
Pulmonary Valve	2D	Valve Mobility	Valve Mobility
	Color Doppler	Evaluate for Insufficiency	Evaluate for Insufficiency Without Sheath
Tricuspid Valve	2D	Valve Mobility	Valve Mobility
	Color Doppler	Evaluate for Insufficiency	Evaluate for Insufficiency Without Sheath
LV Function	2D	Systolic Function	Systolic Function
Pericardium	2D	Effusion	Effusion

2D = Two dimensional; LPA = left pulmonary artery; LV = left ventricle; PDA = patent ductus arteriosus



**FIGURE 6** Color compare transthoracic echocardiogram post Amplatzer Piccolo Occluder deployment. A 4-2 mm Piccolo device is in good position with trivial residual shunting through the center of the device (“flame sign”).

## NICU Bedside Closure

The APO has been FDA approved for TCPC in ELBW neonates for >3 years, and our institution has closed >300 PDA in patients <4 kg. This process took many years of hard work and dedication from multiple members of the TCPC team. The immense experience and expertise in performing TCPC in ELBW neonates in the catheterization lab allowed the TCPC program at Le Bonheur Children’s Hospital to begin performing TCPC at the bedside in the NICU. There are similarities between the two procedures, but the main differences are highlighted in this section.

The catheterization team will need to anticipate all the potential needs of the TCPC procedure to bring the appropriate supplies to the NICU. The supplies routinely used for TCPC in the catheterization lab are now kept in a portable cart to mobilize for a bedside-TCPC procedure. We recommend having duplicates of all supplies routinely used available to account for supplies that may be defective or become contaminated. Ensure that the lead aprons of the key personnel to be present in the patient room are available. A discussion about the appropriate size APO device based off the most recent TTE occurs prior to the case so the different potential sizes are available. At least three catheterization lab staff are needed for a successful procedure: one to scrub and assist with the procedure, one to record and operate the mobile digital Mini C-Arm (Orthoscan, Scottsdale, AZ), and one to circulate needed equipment.

Currently, it is our practice that the dedicated pediatric cardiac anesthesiology team performs sedation for the procedure in the NICU. The anesthesiologist and CRNA assess the patient to ensure adequate IV access for sedation is present and help position the patient for the procedure. At our institution our NICU has single patient rooms that are not all designed the same. Therefore, the position of the isolette is unique to each situation. However, the same concepts apply to each case where the isolette is pulled about 1-2 feet away from the wall to allow the anesthesia team access to the head of the bed to manage the patient airway. The isolette is centered in the room to allow for the catheterization team space on the patient right side and the echocardiographic team space to the patient left side. The mobile patient bedside tray is covered with a sterile drape and situated at the foot end of the isolette to allow for room for the catheterization team to rest the distal end of the sterile catheters and wires.

Patient positioning, groin preparation, and drapes are consistent with the catheterization lab techniques. Once the patient is draped, the mobile digital Mini C-Arm is brought into the expected correct lateral position. At this time, a pre-procedural TTE is performed evaluating the same key structures described above (Table 1). However, there are key differences to the TTE evaluation in the bedside NICU compared to the catheterization lab (Figure 6). First, the echocardiogram machine is turned 180 degrees compared to the catheterization lab. Second, the cardiologist has been performing the echocardiogram as there is minimal space at the bedside for both a sonographer and a cardiologist. Third, the cardiologist is situated as to scan with the right hand during the procedure where the sonographer is situated to scan with the left hand in the catheterization lab. Finally, the pre-procedural PDA measurements can only be made by TTE, as the mobile digital Mini C-Arm does not allow for angiographic measurements.

Ultrasound-guided vascular access of the femoral vein is obtained, and the portable ultrasound machine is removed from the patient room. The mobile digital Mini C-Arm or the patient bed is moved to get the appropriate lateral fluoroscopy view. There is no anteroposterior (AP) fluoroscopy available throughout the case and this represents the biggest difference between the bedside technique and the catheterization lab technique. The implanter must feel comfortable directing the catheter from the right atrium across the tricuspid valve into the right ventricle toward the right ventricular outflow tract and ultimately across the PDA without the AP fluoroscopy. TTE guided catheter access into the PDA can supplement lateral fluoroscopy if requested. A lateral angiogram is obtained in the same manner described above, but measurements of the PDA by angiography are not available on the mobile digital Mini C-arm.

The “push-pull” technique of intraductal APO deployment remains the same as the catheterization technique. A lateral angiogram can be performed to evaluate the LPA is optional as the TTE can evaluate any potential complications. TTE is used to assess the aortic arch, LPA, and device for residual shunting as described above (Table 1). Once correct device position is confirmed the device is released per the same technique as the catheterization lab. TTE is repeated just after device release and five minutes after release to confirm no device migration or other complications. Should complications arise due to device migration or embolization the options for patient care should be discussed at that time. If biplane fluoroscopy is required to complete the necessary intervention, the patient will need to be transferred to the catheterization lab in a timely manner.

## Future Directions

Le Bonheur Children’s Hospital now performs a majority of TCPC in ELBW neonates at the bedside in the NICU. This marks a stark program shift and takes years of collaboration and expertise to ensure a smooth transition. We anticipate bedside-TCPC to be the predominate location of TCPC in ELBW infants at our institution. We are actively collecting data on procedure specifics and anticipate similar procedure times, lower radiation exposure, and similar success and complications rates with the catheterization technique.

One of the criticisms of TCPC in ELBW neonates is the technique requires transfer to a catheterization lab. Now that we show bedside-TCPC is feasible and safe, we anticipate a potential increase in device closure of PDAs in ELBW neonates. However, our institution is an all-referral children’s hospital without a delivery ward. Therefore, every patient at our institution is their second bed. If we can continue to show acceptable safety and efficacy of bedside-TCPC, we plan to perform bedside-TCPC in outlying NICUs that do not have access to pediatric catheterization labs. This will



# Pediatric Cardiologist - Interventional

## Successful Pediatric Cardiology Practice - Tropical Florida Coast

Pediatric Cardiology Associates, located in Tampa Bay on Florida's Gulf Coast, is seeking a BC Interventional Pediatric Cardiologist with advanced fellowship training and experience in Congenital Intervention.

- Ideally seeking candidates with a minimum of 5 years of experience post-fellowship
- Large, experienced, well-established team of 14 pediatric cardiologists and 3 NPs with offices and clinics located throughout the Tampa/St. Petersburg area
- Offer comprehensive congenital cardiac care from fetal life through adulthood
- The team includes members of all pediatric cardiology sub-specialties including: fetal, advanced imaging (CT, MRI, 3D echo), intervention, electrophysiology, cardiomyopathy/heart failure, prevention, and adult congenital
- The interventional team performs over 400 catheterizations per year, about 60% of which are interventions
- Recent interventional team accomplishments include:
  - Implanting their 150<sup>th</sup> transcatheter pulmonary valve, Summer 2019
  - First program in Florida to implant the Gore Cardioform ASD Occluder, Fall 2019
  - Only program in Tampa Bay currently offering PDA device closure for premature newborns, first implant, Winter 2016
- Sub-specialty clinics include general pediatric cardiology, intervention, pulmonary hypertension, cardiomyopathy, ACHD, electrophysiology, and prevention
- Our ACHD program is the ONLY certified Adult Congenital Heart Association program in central Florida
- This position also offers:
  - Full time interventional duties with expected procedural volume of 200+ catheterizations per year
  - No expectation of inpatient service coverage
  - 24/7 collaboration with our excellent pediatric cardiac surgical and pediatric cardiac intensive care teams at St Joseph's Children's Hospital
- Our center offers a unique depth of hospital infrastructure:
  - Two state of the art 1000+ square foot hybrid capable catheterization labs/ORs (one biplane, one single plane)
  - Two additional biplane catheterization labs
  - Two EP labs
- We have the added benefits of a children's hospital inside a large tertiary adult hospital simplifying care across all patient ages with easy access to consultants from all pediatric and adult specialties
- Pediatrix, as a national pediatric cardiology group with over 125 pediatric cardiologists, provides opportunities for quality initiatives that can have national impact
- We offer an attractive schedule allowing freedom to enjoy a great quality of life
- Generous compensation package offered

Tampa Bay's warm weather affords plenty of opportunities to relish the great outdoors year round. You will live in a region others only get to enjoy on vacation. Golf at one of nearly 100 courses or relax on one of the many pristine white-sand beaches. The area offers an assortment of family venues such as zoos, aquariums, theme parks, and state parks. Additionally Tampa Bay offers access to world-class museums, professional sporting events and the performing arts. There is a wide range of residential choices to fit every budget and lifestyle – whether you are looking for big city downtown living, golf course communities, waterfront lifestyle, majestic horse farms or historic neighborhoods.

Effective November 1 st , 2021 Pediatrix will require all employees and new hires to be vaccinated against COVID-19, unless they qualify for an approved medical and/or religious exemption.

### Benefits

Our clinicians enjoy a competitive compensation package with many locations offering sign on bonuses, relocation and tuition reimbursement.

\*Our benefits include:

- Health (various options), life, vision, dental and disability insurance
- 401(k) with annual matching program
- Advanced and continuing medical education
- Leadership training and advancement opportunities
- Employee stock purchase plan at a 15% discount
- Professional liability insurance
- Support and payment for mandatory license/s and hospital credentialing

\*These benefits are for full time employees, employees in other types of employment classifications may be eligible for some of these benefits.

Mednax, Inc. and its affiliated practices operating as Pediatrix® Medical Group (Pediatrix) are one of the nation's leading providers of highly specialized health care for women, babies and children. Since 1979, Pediatrix-affiliated clinicians have been committed to providing coordinated, compassionate and clinically excellent services across the continuum of care, both in hospital settings and office-based practices. Specialties including obstetrics, maternal-fetal medicine, and neonatology are complemented by 18 pediatric subspecialties and a newly expanded area of primary and urgent care clinics. The group's high-quality, evidence-based care is bolstered by investments in research, education, quality-improvement and safety initiatives.

Today through almost 8,000 employees in 38 states, dedicated teams including physicians, advanced practitioners, clinical leaders, business and operational experts work together every day to fulfill our mission to take great care of the patient®. We invite you to join the Pediatrix® family and help shape the future of health care for women, babies and children.

### Pediatrix is an Equal Opportunity Employer

All qualified applicants will receive consideration for employment without regard to race, color, religion, sex, sexual orientation, gender identity, national origin, disability or veteran status.

Apply Here: <https://www.click2apply.net/X77kbbcVVbJapu8NWl5R22>

PI171171158



truly allow TCPC to be a one-bedside procedure and prevent transfer of patients to our institution for procedural expertise. If the procedural team can travel to all referring NICUs instead of the babies being transferred, one could then truly consider TCPC performed very early on in life. Our long-term goal of the TCPC program is to become portable to provide true one-bedside-TCPC.



**FIGURE 7** The 4<sup>th</sup> PDA Symposium is scheduled for August 5<sup>th</sup>-6<sup>th</sup>, 2022, at the Disneyland Hotel, Anaheim, CA, in collaboration with NeoHeart.

Collaboration with neonatology and cardiology will continue to improve the care of ELBW neonates with a hsPDA. If you are interested in learning more about building a TCPC program, curious about outcomes following the procedure, and want to connect with the leading neonatologists and cardiologists in PDA management and TCPC, the 4<sup>th</sup> PDA symposium will help you accomplish these goals. The 4<sup>th</sup> PDA Symposium is scheduled for August 5<sup>th</sup>-6<sup>th</sup>, 2022, at the Disneyland Hotel, Anaheim, CA, in collaboration with NeoHeart. Please join us for an excellent discussion of PDA management, TCPC challenging cases, update on multicenter randomized trials, and ways to move forward with TCPC. Also, a consensus statement on the management of PDA in extremely premature infants will be formalized at the 4<sup>th</sup> PDA Symposium (**Figure 7**). To learn more please visit, <https://pdasymposium.com/>.

## Conclusion

While strategies for PDA management in ELBW neonates evolve, TCPC is a safe and effective method to close PDAs in this vulnerable population. TCPC performed before four weeks is associated with improved respiratory outcomes and better weight gain. Catheterization lab TCPC has an excellent success rate with low adverse event rate across multiple institutions. Centers beginning a TCPC program should start in the catheterization lab with patients between 2-3 kg before transitioning to 1-2 kg and, ultimately, patients <1kg, and only transition to the bedside once the technique is mastered. Based on our early experience, bedside-TCPC appears to be just as safe and effective as catheterization lab TCPC.

## References

1. Parkerson S, Philip R, Talati A and Sathanandam S. Management of Patent Ductus Arteriosus in Premature Infants in 2020. *Front Pediatr.* 2020;8:590578.
2. Bischoff AR, Jasani B, Sathanandam SK, Backes C, Weisz DE and McNamara PJ. Percutaneous Closure of Patent Ductus Arteriosus in Infants ≤1.5 kg: A Meta-Analysis. *J Pediatr.* 2020.
3. Backes CH, Rivera BK, Bridge JA, Armstrong AK, Boe BA, Berman DP, Fick T, Holzer RJ, Hijazi ZM, Abadir S, Justino H, Bergersen L, Smith CV and Kirpalani H. Percutaneous Patent Ductus Arteriosus (PDA) Closure During Infancy: A Meta-analysis. *Pediatrics.* 2017;139.

4. Sathanandam S, Balduf K, Chilakala S, Washington K, Allen K, Knott-Craig C, Rush Waller B and Philip R. Role of Transcatheter patent ductus arteriosus closure in extremely low birth weight infants. *Catheter Cardiovasc Interv.* 2019;93:89-96.
5. Willis A, Pereiras L, Head T, Dupuis G, Sessums J, Corder G, Graves K, Tipton J and Sathanandam S. Transport of extremely low birth weight neonates for persistent ductus arteriosus closure in the catheterization lab. *Congenit Heart Dis.* 2019;14:69-73.
6. Sathanandam S, Whiting S, Cunningham J, Zurakowski D, Apalodimas L, Waller BR, Philip R and Qureshi AM. Practice variation in the management of patent ductus arteriosus in extremely low birth weight infants in the United States: Survey results among cardiologists and neonatologists. *Congenit Heart Dis.* 2019;14:6-14.
7. Jain A and Shah PS. Diagnosis, Evaluation, and Management of Patent Ductus Arteriosus in Preterm Neonates. *JAMA Pediatr.* 2015;169:863-72.
8. van Laere D, van Overmeire B, Gupta S, El-Khuffash A, Savoia M, McNamara PJ, Schwarz CE and de Boode WP. Application of NPE in the assessment of a patent ductus arteriosus. *Pediatr Res.* 2018;84:46-56.
9. Alan S, Karadeniz C, Okulu E, Kilic A, Erdev O, Ucar T, Atasay B, Atalay S and Arsan S. Management of patent ductus arteriosus in preterm infants: clinical judgment might be a fair option. *J Matern Fetal Neonatal Med.* 2013;26:1850-4.
10. Bischoff AR, Giesinger RE, Bell EF and McNamara PJ. Precision medicine in neonatal hemodynamics: need for prioritization of mechanism of illness and defining population of interest. *J Perinatol.* 2020;40:1446-1449.
11. Shepherd JL and Noori S. What is a hemodynamically significant PDA in preterm infants? *Congenital Heart Disease.* 2019;14:21-26.
12. Semberova J, Sirc J, Miletin J, Kucera J, Berka I, Sebkova S, O'Sullivan S, Franklin O and Stranak Z. Spontaneous Closure of Patent Ductus Arteriosus in Infants ≤1500 g. *Pediatrics.* 2017;140.
13. Sehgal A and Menahem S. Interparametric correlation between echocardiographic markers in preterm infants with patent ductus arteriosus. *Pediatr Cardiol.* 2013;34:1212-7.
14. Sehgal A and McNamara PJ. Does echocardiography facilitate determination of hemodynamic significance attributable to the ductus arteriosus? *Eur J Pediatr.* 2009;168:907-14.
15. de Freitas Martins F, Ibarra Rios D, MH FR, Javed H, Weisz D, Jain A, de Andrade Lopes JM and McNamara PJ. Relationship of Patent Ductus Arteriosus Size to Echocardiographic Markers of Shunt Volume. *J Pediatr.* 2018;202:50-55.e3.
16. Mertens L, Seri I, Marek J, Arlettaz R, Barker P, McNamara P, Moon-Grady AJ, Coon PD, Noori S, Simpson J and Lai WW. Targeted Neonatal Echocardiography in the Neonatal Intensive Care Unit: practice guidelines and recommendations for training. Writing Group of the American Society of Echocardiography (ASE) in collaboration with the European Association of Echocardiography (EAE) and the Association for European Pediatric Cardiologists (AEPC). *J Am Soc Echocardiogr.* 2011;24:1057-78.
17. Clyman RI, Saha S, Jobe A and Oh W. Indomethacin prophylaxis for preterm infants: the impact of 2 multicentered randomized controlled trials on clinical practice. *J Pediatr.* 2007;150:46-50.e2.
18. Benitz WE and Bhombal S. The use of non-steroidal anti-inflammatory drugs for patent ductus arteriosus closure in preterm infants. *Semin Fetal Neonatal Med.* 2017;22:302-307.
19. El-Khuffash A, Weisz DE and McNamara PJ. Reflections of the changes in patent ductus arteriosus management during the last 10 years. *Arch Dis Child Fetal Neonatal Ed.* 2016;101:F474-8.
20. Clyman RI, Liebowitz M, Kaempff J, Erdev O, Bulbul A, Håkansson S, Lindqvist J, Farooqi A, Katheria A, Sauberan J, Singh J, Nelson K, Wickremasinghe A, Dong L, Hassinger DC, Aucott SW, Hayashi M, Heuchan AM, Carey WA, Derrick M, Fernandez E, Sankar M, Leone T, Perez J and Serize A. PDA-TOLERATE Trial: An Exploratory Randomized Controlled Trial of Treatment of Moderate-to-Large Patent Ductus Arteriosus at 1 Week of Age. *J Pediatr.* 2019;205:41-48.e6.

# Nicklaus Children's Hospital Proudly Welcomes Pediatric Cardiologist Dr. Nancy Dobrolet



Nancy Dobrolet, MD  
Pediatric Cardiologist

Nicklaus Children's Hospital is honored to welcome board-certified pediatric cardiologist Nancy Dobrolet, MD. Her career includes advanced training in cardiac intensive care and interventional cardiology right here at Nicklaus Children's Hospital. Her residency and fellowship training took her from New York to Boston to Pittsburgh and now she has returned to South Florida to provide compassionate care to infants, children and young adults with congenital heart disease in Plantation and Weston.

For over 70 years Nicklaus Children's Hospital has served as the region's pediatric care leader, providing a full range of services for every stage of your child's life. Through Nicklaus Children's Pediatric Specialists, the organization's medical group practice, we offer expertise in nearly every pediatric subspecialty, including our nationally ranked services for cardiology and cardiovascular surgery.

To learn more about her background or training, please visit [nicklauschildrens.org/DrDobrolet](https://nicklauschildrens.org/DrDobrolet)

---

To schedule a consultation with  
Dr. Dobrolet, please call **305-662-8301**.

---



**Nicklaus  
Children's  
Hospital**

Where Your Child Matters Most

Nicklaus Children's Dan Marino Outpatient Center | 2900 S. Commerce Pkwy., Weston, FL 33331

Nicklaus Children's Pediatric Specialists Office (located inside the Children's Medical Association office space) | 8430 W. Broward Blvd., Plantation, FL 33324



21. Raval MV, Laughon MM, Bose CL and Phillips JD. Patent ductus arteriosus ligation in premature infants: who really benefits, and at what cost? *J Pediatr Surg.* 2007;42:69-75; discussion 75.
22. Mirea L, Sankaran K, Seshia M, Ohlsson A, Allen AC, Aziz K, Lee SK and Shah PS. Treatment of patent ductus arteriosus and neonatal mortality/morbidities: adjustment for treatment selection bias. *J Pediatr.* 2012;161:689-94.e1.
23. Weisz DE, More K, McNamara PJ and Shah PS. PDA ligation and health outcomes: a meta-analysis. *Pediatrics.* 2014;133:e1024-46.
24. Benitz WE. Treatment of persistent patent ductus arteriosus in preterm infants: time to accept the null hypothesis? *J Perinatol.* 2010;30:241-52.
25. Liebowitz M, Kaempf J, Erdeva O, Bulbul A, Håkansson S, Lindqvist J, Farooqi A, Katheria A, Sauberan J, Singh J, Nelson K, Wickremasinghe A, Dong L, Hassinger DC, Aucott SW, Hayashi M, Heuchan AM, Carey WA, Derrick M, Wolf IS, Kimball A, Sankar M, Leone T, Perez J, Serize A and Clyman RI. Comparative effectiveness of drugs used to constrict the patent ductus arteriosus: a secondary analysis of the PDA-TOLERATE trial (NCT01958320). *J Perinatol.* 2019;39:599-607.
26. Philip R, Tailor N, Johnson JN, Apalodimas L, Cunningham J, Hoy J, III BRW and Sathanandam S. Single-Center Experience of 100 Consecutive Percutaneous Patent Ductus Arteriosus Closures in Infants &#x2264;1000 Grams. *Circulation: Cardiovascular Interventions.* 2021;14:e010600.
27. Philip R, Waller BR, Chilakala S, Graham B, Stecchi N, Apalodimas L, Cunningham J, Washington K and Sathanandam S. Hemodynamic and clinical consequences of early versus delayed closure of patent ductus arteriosus in extremely low birth weight infants. *J Perinatol.* 2020.
28. Sathanandam SK, Gutfinger D, O'Brien L, Forbes TJ, Gillespie MJ, Berman DP, Armstrong AK, Shahanavaz S, Jones TK, Morray BH, Rockefeller TA, Justino H, Nykanen DG and Zahn EM. Amplatzer Piccolo Occluder clinical trial for percutaneous closure of the patent ductus arteriosus in patients  $\geq 700$  grams. *Catheter Cardiovasc Interv.* 2020.
29. Tomasulo CE, Gillespie MJ, Munson D, Demkin T, O'Byrne ML, Dori Y, Smith CL, Rome JJ and Glatz AC. Incidence and fate of device-related left pulmonary artery stenosis and aortic coarctation in small infants undergoing transcatheter patent ductus arteriosus closure. *Catheter Cardiovasc Interv.* 2020;96:889-897.
30. Philip R, Towbin J, Tailor N, Joshi V, Johnson JN, Naik R, Waller BR, 3rd and Sathanandam S. Feasibility and Safety of Percutaneous Cardiac Interventions for Congenital and Acquired Heart Defects in Infants  $\leq 1000$  g. *Children (Basel).* 2021;8.
31. Garg R, Zahn E, Sathanandam S and Johnson JN. Transcatheter patent ductus arteriosus closure in extremely premature infants. *Progress in Pediatric Cardiology.* 2021;61:101366.
32. Apalodimas L, Waller Iii BR, Philip R, Crawford J, Cunningham J and Sathanandam S. A comprehensive program for preterm infants with patent ductus arteriosus. *Congenit Heart Dis.* 2019;14:90-94.
33. Schneider DJ and Moore JW. Patent ductus arteriosus. *Circulation.* 2006;114:1873-82.
34. Sathanandam S, Apalodimas L, Weems M, Waller Iii BR and Philip R. Establishing a robust transcatheter PDA closure program for extremely low birth weight infants. *Congenital Cardiology Today.* 2018.
35. Sathanandam S, Agrawal H, Chilakala S, Johnson J, Allen K, Knott-Craig C, Rush Waller B and Philip R. Can transcatheter PDA closure be performed in neonates  $\leq 1000$  grams? The Memphis experience. *Congenit Heart Dis.* 2019;14:79-84.
36. Agrawal H, Waller BR, 3rd, Surendan S and Sathanandam S. New Patent Ductus Arteriosus Closure Devices and Techniques. *Interv Cardiol Clin.* 2019;8:23-32.
37. Sathanandam S, Justino H, Waller BR, 3rd, Radtke W and Qureshi AM. Initial clinical experience with the Medtronic Micro Vascular Plug™ in transcatheter occlusion of PDAs in extremely premature infants. *Catheter Cardiovasc Interv.* 2017;89:1051-1058.
38. Nasef MA, Sullivan DO, Ng LY, Walsh KP, Oslizlok P, McCrossan B, Kenny D and Sathanandam S. Use of the Medtronic Microvascular Plug 7Q for transcatheter closure of large patent ductus arteriosus in infants weighing less than 2.5kg. *Catheter Cardiovasc Interv.* 2022.
39. Lasswell SM, Barfield WD, Rochat RW and Blackmon L. Perinatal regionalization for very low-birth-weight and very preterm infants: a meta-analysis. *Jama.* 2010;304:992-1000.
40. Chung JH, Phibbs CS, Boscardin WJ, Kominski GF, Ortega AN and Needleman J. The effect of neonatal intensive care level and hospital volume on mortality of very low birth weight infants. *Med Care.* 2010;48:635-44.
41. Zahn EM, Peck D, Phillips A, Nevin P, Basaker K, Simmons C, McRae ME, Early T and Garg R. Transcatheter Closure of Patent Ductus Arteriosus in Extremely Premature Newborns: Early Results and Midterm Follow-Up. *JACC Cardiovasc Interv.* 2016;9:2429-2437.
42. Pouladar TM, Wong R, Almeida-Jones M, Zahn E and Lubin L. Bedside Transcatheter Patent Ductus Arteriosus Device Occlusion in an Extremely Low Birth Weight Neonate: A Novel Approach in a High-Risk Population. *Case Rep Anesthesiol.* 2021;2021:4716997.
43. Zahn EM, Nevin P, Simmons C and Garg R. A novel technique for transcatheter patent ductus arteriosus closure in extremely preterm infants using commercially available technology. *Catheter Cardiovasc Interv.* 2015;85:240-8.
44. Bentham J, Meur S, Hudsmith L, Archer N and Wilson N. Echocardiographically guided catheter closure of arterial ducts in small preterm infants on the neonatal intensive care unit. *Catheter Cardiovasc Interv.* 2011;77:409-15.
45. Johnson JN, Sathanandam S, Naik R and Philip R. Echocardiographic guidance for transcatheter patent ductus arteriosus closure in extremely low birth weight infants. *Congenit Heart Dis.* 2019;14:74-78.
46. Alexander J, Yohannan T, Abuteneh I, Agrawal V, Lloyd H, Zurakowski D, Waller BR, 3rd and Sathanandam S. Ultrasound-guided femoral arterial access in pediatric cardiac catheterizations: A prospective evaluation of the prevalence, risk factors, and mechanism for acute loss of arterial pulse. *Catheter Cardiovasc Interv.* 2016;88:1098-1107.
47. Tadphale S, Yohannan T, Kauffmann T, Maller V, Agrawal V, Lloyd H, Waller BR and Sathanandam S. Accessing Femoral Arteries Less than 3 mm in Diameter is Associated with Increased Incidence of Loss of Pulse Following Cardiac Catheterization in Infants. *Pediatr Cardiol.* 2020;41:1058-1066.
48. Kim HS, Schechter MA, Manning PB, Egtesady P, Balzer DT, Shahanavaz S, Rockefeller TA and Abarbanell AM. Surgical Versus Percutaneous Closure of PDA in Preterm Infants: Procedural Charges and Outcomes. *J Surg Res.* 2019;243:41-46.
49. Paudel G, Johnson JN, Philip R, Tailor N, Fahnhorst S, Briceno-Medina M, Stecchi N, Waller BR and Sathanandam S. Echocardiographic versus Angiographic Measurement of the Patent Ductus Arteriosus in Extremely Low Birth Weight Infants and the Utility of Echo Guidance for Transcatheter Closure. *J Am Soc Echocardiogr.* 2021.
50. Philip R, Waller BR, 3rd, Agrawal V, Wright D, Arevalo A, Zurakowski D and Sathanandam S. Morphologic characterization of the patent ductus arteriosus in the premature infant and the choice of transcatheter occlusion device. *Catheter Cardiovasc Interv.* 2016;87:310-7.
51. Philip R, Spagnoli J, DuPuis G, Harvey E, Stecchi N, Tailor N, Waller B and Sathanandam S. Myocardial protection strategies during transcatheter PDA closure in ELBW infants to prevent post-ligation syndrome. *Journal of the American College of Cardiology.* 2021;77:453-453.
52. Tailor N, Philip R, Harvey E, Dupuis G, Waller Iii BR, Stecchi N and Sathanandam S. Effect of intravenous calcium administration during transcatheter PDA closure in ELBW infants. *Catheter Cardiovasc Interv.* 2020;95:1-229.
53. Sathanandam S, Gutfinger D, Morray B, Berman D, Gillespie M, Forbes T, Johnson JN, Garg R, Malekzadeh-Milani S, Fraise A, Baspinar O and Zahn EM. Consensus Guidelines for the Prevention and Management of Periprocedural Complications of Transcatheter Patent Ductus Arteriosus Closure with the Amplatzer Piccolo Occluder in Extremely Low Birth Weight Infants. *Pediatric Cardiology.* 2021.





**JASON N. JOHNSON, MD MHS**  
 Associate Chief, Pediatric Cardiology  
 Director, Cardiac MRI  
 Associate Professor of Pediatrics and Radiology  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital  
 St. Jude Children's Research Hospital  
 49 N. Dunlap Street, Faculty Office Building, 366  
 Memphis, Tennessee, USA  
 901.287.6819, [jjohn315@uthsc.edu](mailto:jjohn315@uthsc.edu)



**RANJIT PHILIP, MD**  
 Director, Pediatric Cardiology Clinic  
 Associate Professor of Pediatrics  
 The University of Tennessee Health  
 Science Center  
 Le Bonheur Children's Hospital  
 Memphis, Tennessee, USA



**MARK F. WEEMS, MD**  
 Medical Director, Neonatal Intensive Care Unit  
 Associate Professor of Pediatrics  
 The University of Tennessee Health  
 Science Center  
 Le Bonheur Children's Hospital  
 Memphis, Tennessee, USA



**AJAY TALATI, MD**  
 Chief of Neonatology  
 Vice Chair, Education  
 Sheldon B. Korones Professor of Pediatrics  
 The University of Tennessee Health  
 Science Center  
 Le Bonheur Children's Hospital  
 Memphis, Tennessee, USA



**SHYAM SATHANANDAM, MD FSCAI**  
 Medical Director of Interventional  
 Catheterization Laboratory  
 Director of The Large Animal Cardiovascular  
 Research Center  
 Associate Professor of Pediatrics  
 The University of Tennessee Health  
 Science Center  
 Le Bonheur Children's Hospital

**RONAK NAIK, MD, DNB, FACC**

Director, Fetal Cardiology Program & Associate Professor of Pediatrics  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital

**VIJAYA JOSHI, MD**

Medical Director of Non-Invasive Cardiology, Professor of Pediatrics  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital, St. Jude Children's Research Hospital

**ASHLEY GIANNINI, RCIS**

Cardiovascular Invasive Specialist  
 Heart Institute  
 Le Bonheur Children's Hospital

**KAITLYN GREER, RT(R)**

Cardiac Cath Lab Rad Tech  
 Heart Institute  
 Le Bonheur Children's Hospital

**CHRISTINA TUNNEY, RN, BSN**

Cardiac Cath Lab Nurse  
 Heart Institute  
 Le Bonheur Children's Hospital

**MATTHEW BALMUT, BS, BSN**

Cardiac Cath Lab Nurse  
 Heart Institute  
 Le Bonheur Children's Hospital

**LINDSEY L. PIEROTTI, RDCS**

Cardiac Sonographer  
 Heart Institute  
 Le Bonheur Children's Hospital

**NATHAN STECCHI, RDCS**

Cardiac Sonographer  
 Heart Institute  
 Le Bonheur Children's Hospital

**HUMA ABDULMAJEED, RDCS**

Cardiac Sonographer  
 Heart Institute  
 Le Bonheur Children's Hospital

**THOMAS HAMILTON, MD**

Pediatric Anesthesiologist  
 Le Bonheur Children's Hospital

**TIM HEAD, DO**

Medical Director, Pediatric Anesthesiology, Assistant Professor  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital

**GENEVIEVE DUPUIS, CRNA**

Certified Registered Nurse Anesthetist  
 Le Bonheur Children's Hospital

**JANET SESSUMS, CRNA**

Certified Registered Nurse Anesthetist  
 Le Bonheur Children's Hospital

**LEAH APALODIMAS, MSN, CPNP-AC/PC**

Pediatric Interventional Cardiology Nurse Practitioner, Instructor  
 College of Nursing, The University of Tennessee Health Science  
 Center, Le Bonheur Children's Hospital

**ASHLEY KIENE, MD**

Pediatric Cardiology Fellow  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital

**NEIL TAILOR, MD**

Assistant Professor of Pediatrics  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital

**B. RUSH WALLER III, MD**

Medical Director of Adult Congenital Heart Disease, Professor of Pediatrics  
 The University of Tennessee Health Science Center  
 Le Bonheur Children's Hospital



## Acute Care Cardiology Unit Pediatrician Hospitalist Opportunity

We are in search of a physician candidate for a pediatric acute care cardiology unit (“ACCU”) hospitalist. The ideal candidate is a board eligible/board certified pediatrician interested in cardiology. The hospitalist will collaborate with a team of advanced practice providers, fellows, pediatric residents, and medical students, under the supervision of general cardiology and heart failure faculty.

Clinical responsibilities include primary coverage of patients on a twenty-nine bed ACCU. Patients have a wide variety of cardiac diagnoses and complex management strategies including management of pre- and post-operative cardiac patients, single ventricle physiology, heart and lung transplant and ventricular assist devices.

The position is full time, with rotating shifts that are a combination of weekday, weeknight, weekends, and holidays.

### Qualifications:

- Doctor of Medicine OR Doctor of Osteopathic Medicine equivalent
- Unrestricted licensed physician
- Board eligible/ Board certified Pediatrician
- Approval/credentialing from CCHMC Medical Staff Services Division

There is opportunity to participate in research and quality improvement projects if interested. Candidates are encouraged to submit their apply to the position.

Apply to this position through Cincinnati Children’s Career page: <https://jobs.cincinnatichildrens.org/>

Enter Requisition # 129908 in the box provided and attach your CV, cover letter of interest and three references to your online application.



U.S. News & World Report scored our [Joint Heart Program](#), in conjunction with Kentucky Children’s Hospital, on outcomes & experience; numbers of patients & procedures; key programs, services & staff; professional recognition; quality improvement efforts; and patient support.



# Report From the Field – Early Career Colleagues

Kamel Shibbani, MD

This is the first of a series of articles from the PICS Society that focuses on early career members who have been picked at random. Keep an eye out for other “Report from the Field” articles in the future that highlight our early career members! These articles will serve as an opportunity for our colleagues to share their thoughts on their own experience and on the avenues with which PICS Society can support early career development.

Today we met up with Dr. Bassel Mohammad Nijres (BMN) from the University of Iowa and Dr. Dan McLennan (DM) from the Children’s Hospital of Wisconsin.

Dr. Nijres completed his pediatric cardiology fellowship at Rush University in Chicago and went on to do pediatric interventional cardiology training at Texas Children’s Hospital/ Baylor College of Medicine in Houston. He has been at the University of Iowa two years.

We started out the discussion with Dr. Nijres about the challenges faced by early career interventional cardiologists.

**KS:** Dr. Nijres, thank you for sitting down with us today! I’m curious, what drew you to the field?


**BMN:** A lot of things drew me to the field. The field is continuously growing - every year we have new devices and new procedures. And every year we gain the ability to treat diseases that we could not treat in the past. In addition, I feel that it’s both fun and challenging at the same time. Interpreting your angiograms with your hemodynamic data, understanding the limitation of this data, and then putting all those pieces together, it’s like solving a puzzle!



Dr. Daniel McLennan



Dr. Kamel Shibbani



Dr. Bassel Mohammad Nijres

*“I think PICS Society can play an important role in promoting and facilitating collaboration.”*

Dr. Bassel Mohammad Nijres

**KS:** I’m curious, what are some challenges you ran into as an early career interventionist?

**BMN:** I think my start was a good one because I landed in a good program. The University of Iowa has offered a lot of support; we have a collaborative surgical team, and I have a great mentor in Dr. Aldoss. I think one of the main challenges for an early career interventional cardiologist is going to be finding an appropriate first job. Finding the right environment is crucial. At the same time, I think that those jobs nationwide are rather limited.

**KS:** Given the importance of being in a supportive environment for your first job as an interventionist, what role do you see the PICS Society playing to help facilitate that?

**BMN:** It would be tremendously helpful if the PICS Society can organize a list of available jobs nationwide. Perhaps even provide details about each job like the number of interventionists, surgical and cath volumes, number of surgeons, etc... I think that would be very helpful.

**KS:** That’s a fantastic idea! Beyond that, what role do you wish to see PICS Society playing?

**BMN:** As an early career interventionist, I think it’s important to look for new research projects. Here, I think the PICS Society can play a very important role in promoting and facilitating research collaboration. Certain diseases that need to be studied are rather rare. Take pulmonary vein disease for example - PICS Society can play a vital role in establishing a multicenter study in pulmonary vein disease. Individually, programs don’t have enough volume. But engaging many programs can allow us to compare various treatment options, surgical vs cath vs medical management. To this day, we don’t know what the best treatment option is.

**KS:** You raise a great point Dr. Nijres - Recently, the PICS Society and CCISC have partnered to create the PICS/CCISC Docmatter community with this exact thought in mind, to help facilitate peer-to-peer interaction and promote collaboration. Have you had a chance to interact with the Docmatter community?

**BMN:** I did actually yes! I like the platform and find it to be very helpful.

**KS:** On a more personal note, what sort of projects have you been involved in?

**BMN:** Our program is one of the busier programs as far as transcatheter PDA device closure in premature infants. As I mentioned earlier, as an early career interventionist you want to try to look for new projects, and PDA device closure in premature infants works nicely. The procedure is a newer procedure that we have been doing for the past few years, and we have enough volume to be able to generate and share our data.

**KS:** From a clinical standpoint, what role do you see the PICS Society playing in preparing 4th year interventional cardiology fellows for life as an attending?

**BMN:** I think creating a cath simulation can be very helpful. A place where we can safely practice new techniques and familiarize ourselves with a new device, like the Harmony valve for example. I’m thinking of something like an event, perhaps, where people can meet and listen to dedicated lectures and hear from people who have done such procedures a lot. This, coupled with cath simulations, would be very helpful.

**KS:** Dr. Nijres, thank you for taking some time to chat with us!



## Medical Director of Pediatric Echocardiography Pediatric Electrophysiologist Adult Congenital Heart Disease Cardiologist General Pediatric Cardiologist Charlotte, NC

The Congenital Heart Center at Levine Children's Hospital (LCH) and Sanger Heart & Vascular Institute (SHVI), seek to recruit a **Medical Director of Pediatric Echocardiography, Pediatric Electrophysiologist, Adult Congenital Heart Disease Cardiologist, and General Pediatric Cardiologist** to join their existing faculty. All are new positions in our dynamic, rapidly growing program. Team currently includes: 12 cardiologists, 3 congenital heart surgeons, 5 cardiac intensivists, 4 pediatric cardiac anesthesiologists, 2 pediatric cardiac radiologists, 20 APPs (includes 3 surgical ACPs), 16 sonographers, 5 nurse navigators, and 9 dedicated RNs.

- **Imaging candidates** will have completed an ACGME-accredited fellowship in pediatric cardiology and be BC/BE by the American Board of Pediatrics; with expertise in echocardiography for congenital heart disease, including transthoracic, transesophageal, and fetal echocardiography. Candidate will be recruited as echocardiography lab director or will be expected to transition into that role within one year. Responsibilities will also include both outpatient and inpatient cardiology. A minimum of 5 years of experience and a 4th year imaging fellowship are preferred. Call/weekend coverage on a rotating basis including echo backup call with 5 other imaging physicians.
- **Electrophysiology candidates** will have completed ACGME-accredited fellowships in pediatric cardiology and pediatric electrophysiology, be BC/BE by the American Board of Pediatrics, and be expected to accomplish pediatric electrophysiology certification by the International Board of Heart Rhythm Examiners. Candidate should be skilled in outpatient and inpatient congenital electrophysiology with experience and interest in transcatheter ablations and device implantation/management. Days will be split between the EP lab and clinic. Responsibilities will include on-site/satellite EP outpatient clinics including pacemaker/ICD clinics, inpatient/consult service coverage, remote device management and cardiology/EP call/weekend coverage on a rotating basis. The Pediatric/Adult Congenital Electrophysiology program has grown in procedural volume over 40% in each of the last two years, and is currently staffed by a single electrophysiologist, dedicated EP APP, and dedicated nurse navigator.
- **Adult Congenital Heart Disease (ACHD) candidates** will have completed an ACGME-accredited fellowship in pediatric cardiology or internal medicine, be BC/BE by the American Board of Pediatrics or American Board of Internal Medicine, and be ABIM board certified in Adult Congenital Heart Disease. Responsibilities will also include both outpatient and inpatient pediatric or adult cardiology, depending on primary board certification, and inpatient ACHD service. Call/weekend coverage on a rotating basis including ACHD call split with two other ACHD providers.
- **General Pediatric Cardiology candidates** will have completed an ACGME-accredited fellowship in pediatric cardiology and be BC/BE by the American Board of Pediatrics. Responsibilities will primarily be in the outpatient pediatric cardiology setting with opportunities to rotate on the inpatient pediatric cardiology service. Call/weekend coverage on a rotating basis.

**The Congenital Heart Center**, established in 2010, has been ranked as one of the top-50 pediatric heart centers in the country by U.S. News and World Report for the last nine years. Our comprehensive services include cardiac imaging, diagnostic and interventional catheterization, invasive electrophysiology, dedicated cardiovascular intensive care staff, and regional referral programs in heart failure/transplantation, adult congenital heart disease, and fetal echocardiography. Surgical and cardiac catheterization volume are growing at a rate of 10-15% per year. Our state-of-the-art two lab cardiac catheterization and electrophysiology suite opened in February of 2017, with dedicated staffing and anesthesia teams. Our new outpatient office complex opened in December 2020, designed to treat all patients from fetal cardiology to ACHD. We have one of the most comprehensive Cardiac Neurodevelopment programs in the Southeast, providing a multitude of specialty services to our congenital heart patients in the same office suite. Participation in investigator initiated and multi-center industry sponsored studies is ongoing within the Heart Center, with the support of a clinical research department.

**Sanger Heart & Vascular Institute (SHVI)** is one of the Southeast's largest cardiac and vascular programs. Sanger employs more than 110 physicians in a network of more than 25 locations to provide the highest quality care available to patients with cardiovascular disease throughout North and South Carolina. SHVI has more than 50 years of experience providing world-class, comprehensive acute and chronic cardiovascular services including the region's only heart transplant center and pediatric/congenital heart surgery program.

**Levine Children's Hospital (LCH)** is a state-of-the-art facility in beautiful Charlotte, North Carolina. LCH has 11 floors and 234 inpatient beds, including on-site PICU and CVICU covered 24/7 by in-house intensivists, Progressive Care Unit, Inpatient Observation Unit, Pediatric Rehab Unit, and Pediatric Emergency Department. We are committed to being the region's leading provider of pediatric health care services.

For more information or to submit a CV, please contact:

Phillip Christofferson  
Peter.Christofferson@atriumhealth.org



We also had a chance to catch up with Dr. Daniel McLennan at the Children’s Hospital of Wisconsin. Dr. McLennan completed his pediatric cardiology training in Sydney Children’s Hospital and Bristol Children’s Hospital, before going to Colorado for his interventional training at Children’s Hospital Colorado. From there, he joined the University of Iowa for one and half years before transitioning to Children’s Hospital of Wisconsin, where he’s been for the past year.

**KS:** Dr. McLennan, thank you for your time today! I would love to hear about your experience as an early interventionist and about challenges you’ve faced out of fellowship.

**DM:** People always say that the hardest job you get is always going to be the first one. When you’re coming out of fellowship, you have to be sure to choose a good job. You want someone around you who is senior to help you make the right choices. Some people I know are stuck in jobs where they are the only interventionists straight out of fellowship, and that can be a really difficult situation to be in.

*“You’ll find that most junior [interventionists] are pretty keen to take on projects... Trying to afford more opportunities for early career people will be a great step.”*

*Dr. Daniel McLennan*

**KS:** What role do you see PICS Society playing in helping early career interventionists?

**DM:** One of the things that PICS Society can do for every interventional fellow is to create a mentor-mentee relationship outside of fellowship. This is someone who can serve as an external mentor during fellowship and also someone who can help beyond training. For example, if people are not doing a certain type of procedure that they have an interest in, this mentor-mentee relationship gives them the opportunity to go to another center and get hands on experience, or perhaps to bring the mentor to their center to teach them. Variations of this program were offered before COVID. Now that the world is starting to open up a little more, PICS Society can work with other organizations to help create a mentor-mentee program again.

Beyond just early interventionist, I think the PICS Society is starting to do a lot more. For example, the DocMatter platform that they have going on for different discussions is quite good. One thing they can consider adding to that is a “case of the month” presentation where people

can talk about cases with great learning points. I hate sitting through webinars at 7pm at night but having a pre-recorded “case of the month” that you can watch anytime would be great way of doing it!

Also, it would be nice to have more opportunities for people to be involved in the conferences, especially early career people. The PICS conference might want to separate out, a little bit more, the senior from the junior interventionists. This might create conversations with the junior interventionists that focus on where they are at in their career, which is going to be different from where the senior folks are in theirs.

**KS:** Those are great ideas Dr. McLennan! How about the flip side of that question? What are ways that early career interventionists can contribute to PICS Society?

**DM:** I think you’ll find that most of the junior guys coming out of fellowship are usually pretty keen to take on projects. Often they’re up-to-date with the latest information since they just came through fellowship. They also might have good tips and tricks that they learned during training. I think trying to afford more opportunities for the early career people will be a great step.

**KS:** On a more personal note, can you tell us more about some of the projects you’ve been working on?

**DM:** Some of the stuff I’m interested in is trying to bring in devices from international companies that are not in the US yet. Having worked in Europe and Australia, and having had access to some of the devices that we don’t have in the US, I’ve been working with the FDA on trying to get approval for some of these devices. I also have an interest in neonatal physiology and hemodynamics, for example promoting the understanding of the impact of PDA and ASD closure in neonates. I’ve been trying to improve that in our hospital.

**KS:** Dr. McLennan, you mentioned your involvement in getting approval for new devices. I’m wondering what advice you have for folks that are interested in getting involved in this?

**DM:** If there’s a device that you know of that is not in the US, it starts with simply reaching out to the company that makes it. Sometimes that company is not aware that there is a market for their device in the US, so I can help to push that device into the market. Sometimes the company might know that the market exists and are in the process of obtaining FDA approval. Then you can put your name in the hat to be one of the centers that trials their device. It comes down to seeing where there is an opportunity and seeking it out.

**KS:** Dr. McLennan, thank you very much for your time!



**PICS Society**

Pediatric and Congenital Interventional Cardiovascular Society



**SAVE THE DATE**  
**Symposium 2022**

**GO**  
**SAVE**  
**THE**  
**DATE**

**7-10 SEPT**  
**MARRIOTT**  
**CHICAGO**  
**DOWNTOWN**

Focusing on the latest interventional catheter strategies for congenital and structural heart disease in children and adults.

[www.picsymposium.com](http://www.picsymposium.com)  
[www.CHDinterventions.org](http://www.CHDinterventions.org)

**Boston Children's Hospital**

Where the world comes for answers

## Interventional Cardiologist

The Department of Cardiology at Boston Children's Hospital seeks an experienced interventional cardiologist to join a thriving, high-volume, high-complexity academic practice at a Harvard teaching hospital. The applicant will have completed advanced training in pediatric and congenital interventional cardiology, will be an active practicing pediatric interventionalist with a minimum of five years of prior experience as an independent operator, and have a record of academic productivity as evidenced by publications in medical journals, grant support, patents, or similar. The applicant must be eligible for an unrestricted medical license in the Commonwealth of Massachusetts and for appointment at the rank of Instructor, Assistant or Associate Professor at Harvard Medical School.

The catheterization facilities at Boston Children's Hospital are located in a dedicated cardiac procedural floor, which includes four state-of-the-art, biplane cardiac catheterization laboratories, a single plane lab, and a hybrid operating room in addition to four cardiac operating rooms and a cardiac MRI suite. A central, 18-bed recovery and post-anesthesia unit supports these procedural areas.

Applicants with unique areas of clinical expertise and a successful record of program development will be given priority as potential additions to a diverse existing staff of interventional cardiologists.

Robust clinical and translational resources and opportunities exist within the Department. In addition to procedural responsibilities, the position entails participation in fellowship training, quality initiatives, and limited inpatient services.

### Interested candidates should forward their CV to:

Dr. Diego Porras  
Department of Cardiology  
Boston Children's Hospital  
300 Longwood Avenue  
Boston, MA 02115  
[Diego.Porras@cardio.chboston.org](mailto:Diego.Porras@cardio.chboston.org)

*We are an equal opportunity employer. All qualified applicants will receive consideration for employment without regard to race, color, religion, sex, national origin, disability status, protected veteran status, gender identity, sexual orientation, pregnancy and pregnancy-related conditions or any other characteristic protected by law.*

**HARVARD MEDICAL SCHOOL  
TEACHING HOSPITAL**



# CTA and ACC Create First-Ever Industry Framework for Cardiovascular Technology

The Consumer Technology Association (CTA)<sup>®</sup> and the American College of Cardiology (ACC) announced the first-ever industry framework to evaluate and give guidance for consumer devices or applications designed to improve cardiovascular health. CTA and ACC have convened health and technology industry leaders to advance guidance and best practices for cardiovascular health technology developers and companies. The goal of the project is to reduce the burden of cardiovascular disease on patients and health care systems – allowing more people to access better cardiovascular care while freeing valuable public health resources.

Cardiovascular disease is the leading cause of death globally, taking an estimated 17.9 million lives every year according to the CDC. U.S. spending on cardiovascular disease has steadily increased over the last 20 years to over \$100 billion per year, mostly from costs related to Ischemic Heart Disease, heart failure, stroke and hypertension.

## Best Practices for Consumer Cardiovascular Technology Solutions

“Health tech companies will be in a better position than ever to improve cardiovascular technology with a roadmap to integration for clinicians and their practices,” said Gary Shapiro, president and CEO, CTA. “We have seen enormous breakthroughs in medical technology in the last decade alone, better cardiovascular devices are essential in improving global health and saving lives through stronger preventative care.”

The framework targets the use of consumer cardiovascular technology solutions for prevention, screening, diagnosis and health management or treatment. This guidance gives health tech companies a framework to manufacture products that allow clinicians to confidently integrate cardio technology into their practices. Consumer Cardiovascular Technology Solutions facilitate ongoing cardiovascular health promotion, disease detection and care management, instead of patients solely relying on intermittent and costly clinical visits.

“Use of digital technologies, including consumer wearables and apps for monitoring

heart health, is widespread and has the capability to transform how clinicians work with their patients to treat and prevent heart disease,” said Ritu Thamman, MD, FACC, ACC Innovation Work Group member. “By working with our colleagues on all sides of technology to ensure we are implementing these technologies safely and effectively, we’re helping clinicians use the latest technologies in a way that ensures their patients receive the highest quality care and live better, healthier lives.”

Practical uses for Consumer Cardiovascular Technology Solutions include:

**Prevention:** Consumer devices can monitor physical activity and heart rate to help promote healthy behaviors and give clinicians insights into the cardiovascular health of their patients. Consumer cardiovascular technology also allows clinicians to potentially identify areas of risk sooner.

**Screening/Diagnosis:** Atrial Fibrillation (AFib) is the most common abnormal heart rhythm. It affects about six million Americans and is a major cause of stroke. Cardiovascular Technology Solutions can be used to detect the presence of an irregular rhythm that is potentially AFib and prompt the user to confirm the diagnosis with a clinician.

**Health Management/Treatment:** For people undergoing cardiac rehabilitation (rehab), the first 6-8 months is especially critical to ensure the timely recovery. Wearables that can monitor physical activity and heart rate accurately and remotely are useful tools for home-based cardiac rehab. Leveraging these technologies can help overcome some of the barriers patients face with on-site rehab programs while expanding the breadth and depth of monitoring.

The framework also accounts for the need to protect consumer data rights and privacy. Health technology solutions need to capture comprehensive and privacy sensitive information about their users. But this data can sometimes be aggregated, analyzed, shared or sold in ways that are poorly disclosed or understood. By adhering to industry privacy guidelines, mobile technology developers can ensure that their solutions optimally protect consumers’ privacy, promote trust and mitigate perceived risks about the use and sharing of data.

The project is the first collaboration between the ACC and CTA to create best practices for evaluating Consumer Cardiovascular Technology Solutions. CTA is collaborating with the American College of Cardiology (ACC) through participation in the ACC’s Applied Health Innovation Consortium for the purpose of building a roadmap for Artificial Intelligence and digital technology in cardiology. The Consortium brings together academic, clinical, industry and technology partners and patient advocates to collaborate in the digital transformation of healthcare.

## About Consumer Technology Association

As North America’s largest technology trade association, CTA<sup>®</sup> is the tech sector. Our members are the world’s leading innovators – from startups to global brands – helping support more than 18 million American jobs. CTA owns and produces CES<sup>®</sup> – the most influential tech event in the world. Find us at CTA.tech. Follow us @CTAtech.

## About the American College of Cardiology

The American College of Cardiology envisions a world where innovation and knowledge optimize cardiovascular care and outcomes. As the professional home for the entire cardiovascular care team, the mission of the College and its 54,000 members is to transform cardiovascular care and to improve heart health. The ACC bestows credentials upon cardiovascular professionals who meet stringent qualifications and leads in the formation of health policy, standards, and guidelines. The College also provides professional medical education, disseminates cardiovascular research through its world-renowned JACC Journals, operates national registries to measure and improve care, and offers cardiovascular accreditation to hospitals and institutions. For more, visit <https://www.acc.org/>.





## MAY

06

**CiCA 2022**  
Washington, D.C., USA  
[info@cica-ep.org](mailto:info@cica-ep.org)

19-22

**SCAI 2022 Scientific Sessions**  
Atlanta, Georgia, USA  
<https://scai.org/scai2022>

24-26

**AIMED 2022 The Global Summit**  
San Francisco, California, USA  
<https://ai-med.io/all-events/global-summits/aimed-22/>

## JUNE

10-13

**ASE 2022 – 33rd Annual Scientific Sessions**  
Seattle, Washington, USA  
<https://www.asescientificsessions.org/register-now/>

22-25

**CSI Frankfurt**  
Frankfurt, Germany  
<https://www.csi-congress.org/conferences-courses/conferences/csi-frankfurt>

23-25

**ACHA's 9th National Conference:  
Leading the Way for 1 in 100**  
Bloomington, Minnesota, USA  
<https://www.achaheart.org/get-involved/events/2022/9th-national-conference/>

# Z-6™

## Atrioseptostomy Catheter

*Short distal tip*  
for easier insertion  
and improved rewrapping



NOW AVAILABLE IN THE UNITED STATES



## Transplant Cardiologist with an Interest in Rehabilitation Science

We are seeking a Board Eligible/Board Certified Pediatric Cardiologist with formal 4th year training in Cardiomyopathy, Ventricular Assist Device, and Cardiac Transplantation for a faculty position at the Cincinnati Children's Hospital Medical Center.

The Advanced Cardiomyopathy Program works closely within the Heart Institute, caring for patients with the most challenging conditions, and has been the largest pediatric heart transplant program in the Tri-State area for the last decade. Expertise includes heart, heart-liver, heart-lung, and heart-kidney transplantation. The prospective faculty member will be expected to excel in academic endeavors, have clinical and academic focus in Cardiac Transplantation, and have interests in Cardiopulmonary Exercise Testing and Cardiac Rehabilitation.

### Leaders in Pediatric Heart Transplant

The Heart Transplant Program, part of the interdisciplinary Heart Institute, is the oldest pediatric heart transplant program in Ohio. We are part of the Heart Institute's combined program linking cardiomyopathy/heart failure and mechanical circulatory support care.

Our team of cardiothoracic surgeons, cardiologists, nurse practitioners, and nurses are experts in the treatment of complex heart disease and [heart transplantation](#). We care for infants, children, adolescents, and adults with uncorrectable congenital heart disease and [cardiomyopathy](#), facilitating their candidacy for transplantation, and delivering their post-transplant care.

Our heart transplant program is approved by the United Network for Organ Sharing (UNOS) and a member of the Ohio Solid Organ Transplantation Consortium. In addition, our transplant team members are participants in the Pediatric Heart Transplant Study and the International Society for Heart and Lung Transplantation.

Our goal is to provide children facing untreatable heart disease the best chance to be active members of their families and communities, and the best chance to grow emotionally and physically with as little limitation as possible. Our areas of emphasis include:

- Care provided by a multidisciplinary team of cardiologists, surgeons, intensive care specialists, nurse practitioners, and nurses.
- Outcomes-focused clinical care and research to optimize survival and quality of life.
- Quality initiatives, such as prevention of rejection and use of cutting-edge technologies to detect for harmful antibodies and detect for transplant coronary artery disease.

Our clinical team's extensive experience and our commitment to long-term quality outcomes form the foundation of the Heart Transplant Program.

Interested candidates should apply to this position through Cincinnati Children's Career page: <https://jobs.cincinnatichildrens.org/>. Enter Requisition # 129989 in the box provided and attach your CV, cover letter of interest and three references to your online application.



U.S. News & World Report scored our [Joint Heart Program](#), in conjunction with Kentucky Children's Hospital, on outcomes & experience; numbers of patients & procedures; key programs, services & staff; professional recognition; quality improvement efforts; and patient support.



# Structured Reporting Systems are Expected to Dramatically Improve Patient Care and Data Management

Cardiology is an inherently data-dependent and data-driven practice. Patient medical history records, images, test results, and more go into every diagnosis and treatment decision. However, most organizations don't take advantage of structured reporting; most cardiovascular information systems (CVIS) cannot effectively mine the wealth of data from a facility's cardiology department. There is no debate on how crucial structured reporting is. It is considered the first step to improving patient care and enhanced data accuracy—important points raised by a coalition of 14 professional societies led by the American College of Cardiology and by the Society for Cardiovascular Angiography and Interventions.<sup>1</sup> It's estimated that only 10% of United States cardiac Cath labs use structured reporting to improve efficiency and bolster patient outcomes.<sup>1</sup>

Dr. Serge Makowski, CEO and Co-founder of leading healthcare software development firm MediReport, says, "The goal of structured reporting is to make sure there is consistency in the documentation reporting of every cardiac procedure, regardless of the physician's training, background, or style of work and that 100% of procedural data is recorded abiding by the latest clinical guidelines. The system intelligently guides the clinician on the required and important procedural data and generates near real-time reports which are automatically incorporated into the facility's existing electronic health record. Subsequently, this can be accessed by any healthcare provider, which informs evidence-based treatment decisions, improves billing, and better the patient care experience."

Structured reporting systems fully incorporate data, workflow, analytics, inventory management, billing, and process improvement in one central documentation system.<sup>1</sup> The same data may be repurposed repeatedly for clinical research or registry participation, without the need for any manual data re-entry.

## Basic Features of Structured Reporting

A growing number of healthcare organizations are exploring structured reporting. But not all structured reporting systems are created equal. Clinicians should look for several key features to ensure maximum efficiency, such as:

- The centralization of all information (dosimetry, pressures, measurements, dynamic drawing, key PACS images...) from the cardiac procedure. This means the system must seamlessly interface and operate with the facility's software packages (EMR, modalities, HIS...).
- The effortless recording and tracking of all materials and devices before, during and after the procedure using techniques like barcode scanning.
- The generation of reports complete with full sentences, which may be easily understood by the patient, or any other clinicians involved in follow-up care.
- The ability to select multiple languages for report generation, offering both convenience for patients and flexibility to providers.
- Continuous connectivity to cloud-based platforms which allows remote access to data at any time.

## A Unified System<sup>2</sup>

Two interventional cardiologists developed such a software suite, called CardioReport 360™ to help improve workflows, reduce errors, and save hospitals time and money. This platform offers several advantages compared to other CVIS; one being calculating automatic SYNTAX scoring, which saves time during procedures and reduces error rates to improve diagnostic accuracy. Additionally, the system integrates

in real-time the latest clinical guidelines and classification systems continuously available to generate reports. Another important asset are its templates, which are customizable to the physician and the practice's specific workflows and procedures, which improves clinical adoption and satisfaction during use.

However, one of the most important features of a unified reporting system is the intelligent report conclusions. The auto-generated report conclusions are based off selected key words from procedural data which provides a quick, but thorough, understanding of the lesion's description and the procedure results. Dr. Makowski says: "The algorithm behind the conclusion synthesis is based on 25-plus years of medical expertise that applies globally and is paramount in the patient's care journey and treatment. A bifurcation of a lesion can complicate an angioplasty as the physicians will make a decision between angioplasty and surgery, hence the bifurcation criteria will be selected as an important item in the conclusion. Or, if thrombosis appears in the description of the procedure, it will be selected as a determinant item in the conclusion, as it might complicate the angioplasty and will carry a tremendous impact on the patient's treatment. However, if, let's say, the lesion is concentric, or eccentric is not deemed determinant thus will not be retained in the conclusion."

Lastly, reports are finalized before the patient even leaves the Cath Lab which vastly improves timely reporting as well as valuable Cath Lab throughput time. Referring doctors, as well as those participating in follow-up care, have near instant access to all vital information about the patient and procedure. Patients benefit from no lapse in care, while hospitals and ambulatory surgical centers can boost the bottom line through quicker turnover.

Dr. Makowski says, "We, the physicians, rely on intelligent insights to be gathered from

**NEONATOLOGY TODAY**  
Peer Reviewed Research, News and Information in Neonatal and Perinatal  
Medicine

Neonatology Today is interested in publishing manuscripts from Neonatologists, Fellows, NNPs and those involved in caring for neonates on case studies, research results, hospital news, meeting announcements, and other pertinent topics.

Please submit your manuscript to: [LomaLindaPublishingCompany@gmail.com](mailto:LomaLindaPublishingCompany@gmail.com)



## Pediatric Cardiologist - Non-Invasive Imaging Service

This is an outstanding opportunity to join a vibrant and collegial academic environment and work in a children's hospital ranked as one of the best in the U.S. The Division of Pediatric Cardiology at the Spencer F. Eccles School of Medicine is seeking a Noninvasive Imaging Pediatric Cardiologist, with expertise in cardiac MRI and/or cardiac CT. The Pediatric Cardiologist's primary focus will be in the noninvasive imaging service, sharing all clinical aspects, educational efforts, program development, and quality improvement initiatives associated with the service. Noninvasive imaging physicians provide rotating coverage of night and weekend call for emergency transesophageal echocardiograms or assistance with interpreting complex transthoracic studies. Division members are expected to rotate in covering general cardiology night and weekend call with other Division members. Division members participate in general cardiology outpatient clinic and inpatient service, in addition to their duties in their areas of expertise. Clinical responsibilities may also include seeing patients in an outreach clinic setting. Clinical duties will be carried out at Primary Children Hospital and the Division's outreach sites. In addition to clinical service, there is both opportunity and expectation for academic work, including education, investigation/research and administration, as well as advocacy. There will be protected time for these efforts with a well-established support structure, expected benchmarks, and both breadth and depth in mentoring available within the Division.

Noninvasive imaging includes echocardiography (transthoracic, transesophageal, fetal, and 3D), cardiac MRI, and cardiac CT angiography. Annually, division members read over 13,000 echocardiograms, over 800 fetal echoes, over 300 cardiac MRIs, and over 200 CT angiograms.

Qualified candidates must be Board Eligible/Board Certified in Pediatric Cardiology, with focused training and expertise in noninvasive imaging, and in cardiac MRI and/or cardiac CT. The selected candidate will receive a faculty appointment in the Department of Pediatrics on the Clinical or Tenure Track at the academic level commensurate with experience and qualifications. Leadership opportunities are available for qualified candidates.

The Department and University offers a competitive salary and an unmatched benefits program, including non-contributory retirement contributions of 20.2% of annual salary that vest immediately. The Department offers a faculty development and mentoring program designed to foster success in translational or basic research, quality improvement engagement, and excellence as educators.

Salt Lake City offers an incredible quality of life with a growing economy, rich cultural scene with ballet, theatre, symphony, opera, live music, museums, outstanding restaurants, and a moderate cost of living. The city is a well-known ski destination and a gateway to the state's renowned landscapes. In addition to its 14 ski resorts, Utah boasts five scenic national parks (with five more within a day's drive), a variety of golf courses allowing for year-round play, hundreds of miles of hiking and biking trails (some leaving from adjacent to the hospital), and numerous other outdoor activities.

**Interested individuals can apply for the position at <https://utah.peopleadmin.com/postings/130651>. Cover letter and curriculum vitae are required. For additional information about the position, please contact: Antonio Cabrera, M.D., Division Chief, at [antonio.cabrera@hsc.utah.edu](mailto:antonio.cabrera@hsc.utah.edu).**

*The University of Utah Health (U of U Health) is a patient focused center distinguished by collaboration, excellence, leadership, and respect. The U of U Health values candidates who are committed to fostering and furthering the culture of compassion, collaboration, innovation, accountability, diversity, integrity, quality, and trust that is integral to our mission.*

*The University of Utah values candidates who have experience working in settings with students, staff, faculty, and patients from diverse backgrounds and possess a strong commitment to improving access to higher education, employment opportunities, and quality healthcare for historically underrepresented groups.*

*Individuals from historically underrepresented groups, such as minorities, women, qualified persons with disabilities and protected veterans are encouraged to apply. Veterans' preference is extended to qualified applicants, upon request and consistent with University policy and Utah state law. Upon request, reasonable accommodations in the application process will be provided to individuals with disabilities.*

*The University of Utah is an Affirmative Action/Equal Opportunity employer and does not discriminate based upon race, ethnicity, color, religion, national origin, age, disability, sex, sexual orientation, gender, gender identity, gender expression, pregnancy, pregnancy-related conditions, genetic information, or protected veteran's status. The University does not discriminate on the basis of sex in the education program or activity that it operates, as required by Title IX and 34 CFR part 106. The requirement not to discriminate in education programs or activities extends to admission and employment. Inquiries about the application of Title IX and its regulations may be referred to the Title IX Coordinator, to the Department of Education, Office for Civil Rights, or both.*

the reports, data that otherwise would be ignored. I no longer need to rely on multiple applications to get the complete picture of my patient. Having access to an extensive number of reporting parameters in a single unified CVIS and being able to generate a high quality and accurate report almost instantly, maximizes efficiency and accuracy.”

**About MediReport**

MediReport, a global supplier of cloud-based software for medical applications, was founded in 1995 by cardiologists Dr. Serge Makowski and Dr. Fabrice Beverelli. Through its comprehensive CVIS, CardioReport™ Suite, MediReport provides vastly improved clinical workflows that save time, improve quality, and generate high revenue for client organizations. The company is committed to delivering quality products that address the evolving needs of cardiology and IT technologies, based on its foundational principles: clinical excellence, customer satisfaction, continuous improvement of its products and innovations, and commitment and responsiveness. The system is already used by healthcare providers in more than 40 countries. For more information, visit [medireport.net](http://medireport.net).

**References**

1. Tierney, Mary C.; “Structured Reporting: The Road Map for Better Data, Workflow and Reports”; Cardiovascular Business, Accessed 08 Feb 2022; [cardiovascularbusiness.com/sponsored/1170/topics/practice-management/structured-reporting-road-map-better-data-workflow-and-0](http://cardiovascularbusiness.com/sponsored/1170/topics/practice-management/structured-reporting-road-map-better-data-workflow-and-0).
2. Landing Page; “Powering the best hospitals for 25 years”; Accessed 07 Feb 2022; MediReport; [medireport.net/cardioreport](http://medireport.net/cardioreport).



**Pediatric Cardiologist**

The Department of Pediatrics at Southern Illinois University School of Medicine is recruiting an MD/DO for a fourth pediatric cardiologist position at the Assistant or Associate Professor level. Faculty will join a rapidly expanding cardiology program at our Children’s Hospital, an 80-bed CHA affiliated pediatric referral center for Central and Southern Illinois with a referral base of almost 2 million. The current program includes state-of-the-art noninvasive imaging in TTE, TEE, fetal echocardiogram, and advanced MRI imaging. We have developed a highly successful collaborative clinical and research program with a nationally recognized pediatric cardiology center. Opportunities exist to participate in resident and medical student education and receive an advanced degree in medical education. Candidates must be board eligible in Pediatrics and Pediatric Cardiology. Illinois licensure is required prior to official start date. Travel in central Illinois to outreach clinics is required.

Applications are accepted online at: <https://www.siumed.edu/hr>

For additional information, please contact:

Ramzi Nicolas, MD  
217.545.9706  
[rnicolas@siumed.edu](mailto:rnicolas@siumed.edu)

SIU is an EO/AA employer.



**CHIP NETWORK**  
CONGENITAL HEART INTERNATIONAL PROFESSIONALS



## Pediatric Cardiac Interventionist Position

UC Davis Children's Hospital  
School of Medicine  
Department of Pediatric Cardiology

The Department of Pediatrics at the University of California, Davis School of Medicine is recruiting 1 full-time academic Pediatric Cardiac Interventionist within the Pediatric Cardiology specialty. We are recruiting at the Assistant/Associate/Full Professor level in the Clinical X or Health Science Clinical Series in the Section of Pediatric Cardiology, Department of Pediatrics and specifically trained and experienced in Pediatric Cardiac Interventions. Candidates must possess an M.D. or D.O. degree, be board certified in Pediatric Cardiology and must possess or be eligible for licensure in the State of California.

The interventional cardiology candidate must have at least 3 years of experience in interventions in congenital heart disease and have the desire and ability to build and grow a practice in interventional cardiology volume.

Work distribution will depend on the specific skillset of the candidate. It is expected that the candidate will share in the on-call and weekend/holiday coverage schedule.

In addition to the clinical responsibilities, the ideal candidate will be expected to participate in teaching of medical students, residents and fellows, research activities of the Department of Pediatrics, and serve on departmental committees.

The candidates must have the following qualifications:

- M.D. or D.O.
- Successful completion of an approved pediatric residency training program.
- Successful completion of an approved Pediatric Cardiology fellowship training program.
- Successful completion of a Pediatric Interventional Cardiology advanced fellowship training program.
- Board certification in Pediatric Cardiology.
- Eligibility for a California Medical License.
- Demonstrated proficiency in the teaching of students and housestaff.
- Demonstrated proficiency to perform clinical research.
- Ability to foster collegiality and work collaboratively in a diverse environment, including working closely with Adult Congenital Heart Disease services.
- Service including committees, leadership ability, and community outreach.
- Have Board Certification in Adult Congenital Heart Disease (preferred).

The Pediatric Heart Center at UC Davis Children's Hospital is inland Northern California's only full-service cardiac care facility for children and young adults, offering the latest tests and treatments for a range of congenital or acquired cardiovascular conditions. Our integrated multidisciplinary team of surgeons, specialists, physicians, nurses and researchers offer Northern California's most sophisticated specialized diagnostic, interventional and surgical expertise in comprehensive diagnostic, therapeutic, and surgical procedures for children with heart defects.

For full consideration applications should be received by December 27, 2021. However the position will remain open until filled. Completed applications include CV, Cover Letter, Statement of Contributions to Diversity, Equity, and Inclusion and contact information for 3-5 references.

Candidates should submit their application online at:  
<https://recruit.ucdavis.edu/JPF04580>.

# Hospital Directory 2021-2022

## Published Mid-August

- Hospitals that Offer Open Heart Surgery for Children in North America
- Contact information at each hospital for Chief of Pediatric Cardiology & Fellowship Director
- Lists each hospital's Pediatric Cardiologists & Cardiothoracic Surgeons
- Lists Pediatric Cardiology Fellowships
- Distributed to Division Chiefs by mail
- Electronic version available on CCT's website:  
[www.congenitalcardiologytoday.com](http://www.congenitalcardiologytoday.com)



## CORPORATE TEAM

**PUBLISHER &  
EDITOR-IN-CHIEF**

Kate Baldwin  
*kate.f.baldwin@gmail.com*

**CO-FOUNDER &  
MEDICAL EDITOR**

John W. Moore, MD, MPH  
*jwmmoore1950@gmail.com*

**FOUNDER &  
SENIOR EDITOR**

Tony Carlson  
*tcarlsonmd@gmail.com*

**STAFF EDITOR &  
WRITER**

Virginia Dematatis

**EDITOR-IN-CHIEF  
EMERITUS**

Richard Koulbanis

**STAFF EDITOR**

Loraine Watts

## EDITORIAL BOARD

Teiji Akagi, MD  
Zohair Al Halees, MD  
Mazeni Alwi, MD  
Fadi Bitar, MD  
Jacek Bialkowski, MD  
Mario Carminati, MD  
Anthony C. Chang, MD, MBA  
Horacio Faella, MD  
Yun-Ching Fu, MD  
Ziyad M. Hijazi, MD, MPH

Ralf Holzer, MD  
Marshall Jacobs, MD  
R. Krishna Kumar, MD, DM, MBBS  
John Lamberti, MD  
Gerald Ross Marx, MD  
Tarek S. Momenah, MBBS, DCH  
John W. Moore, MD, MPH  
Carlos A. C. Pedra, MD  
Daniel Penny, MD, PhD  
James C. Perry, MD

Shakeel A. Qureshi, MD  
P. Syamasundar Rao, MD  
Carlos E. Ruiz, MD, PhD  
Girish S. Shirali, MD  
Horst Sievert, MD  
Hideshi Tomita, MD  
Gil Wernovsky, MD  
William C. L. Yip, MD  
Carlos Zabal, MD

## OFFICIAL NEWS & INFORMATION PARTNER OF



**PICS Society**

Pediatric and Congenital Interventional Cardiovascular Society

Statements or opinions expressed in Congenital Cardiology Today reflect the views of the authors and sponsors and are not necessarily the views of Congenital Cardiology Today.

© 2022 by Congenital Cardiology Today  
ISSN 1554-7787 print. ISSN 1554-0499 electronic.  
Published monthly. All rights reserved.